CORRECTED **VERSION***

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZA International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 38/17, 48/00

(11) International Publication Number:

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

A3

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

60/034,204 60/034,205

21 January 1997 (21.01.97) US US

21 January 1997 (21.01.97)

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). GENTZ, Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).
- (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report: 5 November 1998 (05.11.98)

(54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

(57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Amenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
	Australia	GA.	Gabon	LV	Latvia	SZ	Swaziland
AU		GB	United Kingdom	MC	Monaco	TD	Chad
AZ	Azerbaijan	GE	Georgia	MD	Republic of Moldova	TG	Togo
BA	Bosnia and Herzegovina		-	MG	Madagascar	TJ	Tajikistan
BB	Barbados	GH	Ghana	MK	The former Yugoslav	TM	Turkmenistan
BE	Belgium	GN	Guinea	NIK.	Republic of Macedonia	TR	Turkey
BF	Burkina Faso	GR	Greece	241	•	TT	Trinidad and Tobago
BG	Bulgaria	HU	Hungary	ML	Mali	UA	Ukraine
BJ	Benin	IE	Ireland	MN	Mongolia		
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Јарап	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 98/31799 PCT/US98/00959

POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

FIELD OF INVENTION

5

10

15

20

25

30

35

This invention relates to newly identified polynucleotides and the polypeptides encoded by them, the use of such polynucleotides and polypeptides, and their production. More particularly, the polynucleotides and polypeptides of the present invention relate to specific receptor families described in the specification and known in the art. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

BACKGROUND OF THE INVENTION

Receptor proteins are found on the membrane of the cells and are generally involved in signal transduction. There are many types of receptor proteins, and for convenience, these proteins are grouped in families based on similarity in structure and function.

For example, the TM4SF superfamily of cell surface proteins, also known as the tetraspan receptor superfamily, is comprised of at least seventeen individual gene products (these include CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8). The TM4SF superfamily is the second largest group in the CD antigen superfamily. Each member of the TM4SF superfamily can be characterized by several putative physical features including four highly conserved transmembrane domains, two divergent extracellular loops, and two short and highly divergent cytoplasmic tails. Expression patterns for members of the TM4SF superfamily tend to be rather broad and can vary widely between members. The functional roles of TM4SF superfamily members are primarily associated with signal transduction events and pathways, but also include cell adhesion in platelets and other lymphocytic and non-lymphocytic cell lines, as well as cell motility, proliferation, and metastasis. In addition, recent evidence suggests that a subset of the members of the TM4SF superfamily may function as potassium channel molecules.

One member of the TM4SF family, CD20, is a four membrane spanning domain cell surface phosphoprotein expressed exclusively on B lymphocytes. Although the precise functional role of CD20 has yet to be determined, it is thought to function primarily as a receptor during B-cell activation. Furthermore, a large number of experimental observations suggest several additional speculative roles for the CD20 molecule. For example, CD20-specific immunoprecipitation of biochemically cross-linked plasma membrane proteins suggests that CD20 assumes a multimeric structural

10

15

20

25

30

35

conformation characteristic of other previously described membrane channel proteins. Further experimentation has revealed that expression of exogenous CD20 on the cell surface specifically increases Ca²⁺ conductance across the plasma membrane. Together, these results suggest that CD20 complexes may function as B-cell specific Ca²⁺ ion channels. In addition, monoclonal antibodies raised against CD20 have been used to stimulate resting B-cells to transition out of the G0/G1 segment of the cell cycle. It has also been demonstrated that CD20 is associated with both serine and tyrosine kinases and, more specifically, that CD20 is associated, although not directly, with the Src family of tyrosine kinases including p56/53lyn, p56lck, and p59fyn.

2

A second example of a receptor subfamily, called sialoadhesin molecules, belongs to the Ig superfamily of receptor-like molecules. The more than 100 members of the Ig superfamily are generally considered to engage in specific cell-cell interactions through which intercellular communication may occur. In addition to classical protein-protein interactions, intercellular communication may also be mediated through protein-carbohydrate interactions. In fact, all members of the sialoadhesin family of the Ig superfamily are capable of mediating protein-sialic acid binding interactions. To date, only a small number of proteins have been assigned to the sialoadhesin family including sialoadhesin, CD33, CD22, the myelin-associated glycoprotein (MAG), and the Schwann cell myelin protein (SMP). Each of these proteins is expressed in a restricted subset of cell types. For example, CD22 and CD33 are expressed exclusively by B-lymphocytes and cells of the myelomonocytic lineage, respectively.

Similarly, galectins are a family of the lectin superfamily of carbohydrate-binding proteins which have a high affinity for b-galactoside sugars. Although a large number of glycoproteins containing b-galactoside sugars are produced by the cell, only a few will bind to known galectins *in vitro*. Such apparent binding specificity suggests a highly specific functional role for the galectins. Galectin 1 (conventionally termed *LGALS1* for lectin, galactoside-binding, soluble -1) is thought to specifically bind laminin, a highly polylactosaminated cellular glycoprotein, as well as the highly polylactosaminated lysosome-associated membrane proteins (LAMPs). Galectin 1 has also been shown to bind specifically to a lactosamine-containing glycolipid found on olfactory neurons and to integrin a_7b_1 on skeletal muscle cells. Galectin 3 has also been observed to bind specifically to laminin, immunoglobulin E and its receptor, and bacterial lipopolysaccharides.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1-mediated disruption of laminin-integrin a_7b_1 interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined

WO 98/31799 PCT/US98/00959

presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively.

A relatively new family of cell-surface proteins has been identified and termed the Ly6 superfamily. The members of this family include murine and human SCA-2, rat Ly-6 (also termed ThB), human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)], and E48 antigen. The determination of an initial functional role for SCA-2 may lie in an analysis of its expression profile with regard to the complex process of hematopoiesis. SCA-2 is highly expressed in early thymic precusor cells. In turn, progeny of the intrathymic precusor population continue to express SCA-2, but only until the point of transition occurs from blast cell to small cell. Further experimental evidence demonstrates that mature thymocytes and peripheral T-cells do not express detectable levels of SCA-2, whereas mature, peripheral B-cells do continue to express SCA-2. As a result, it seems very likely that SCA-2 plays an important role in thymocyte maturation and differentiation. A plausible explanation for this functional hypothesis is that SCA-2 may act as a receptor for a unknown cytokine which regulates thymocyte maturation and differentiation.

In addition, CD59 is a recently identified integral membrane protein which appears to be involved in the regulation of complement. Recent studies show that the CD59 antigen may prevent damage from complement C5b-9 and protect astrocytes during inflammatory and infectious disorders of the nervous system. Expression of recombinant human CD59 on porcine donor organs have been shown to prevent complement-mediated lysis and activation of endothelial cells that leads to hyperacute rejection. Recently, researchers at Alexion Pharmaceuticals (New Haven, CT) reported on the production of transgenic pigs which expressed human CD59. In these animals, xenogeneic organs were resistant to hyperacute rejection. (Fodor, et al., "Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection," Proc. Natl. Acad. Sci., 91:1153-11157 (1994).) The same company also reported that expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria (PNH) B-

5

10

15

20

25

30

10

15

20

25

30

35

cells confers resistance to human complement. (Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B-cells confers resistance to human complement," Blood, 84:2604-2611 (1994).) PNH is an acquired hematopoietic disorder characterized by complement-mediated hemolytic anemia, pancytopenia, and venous thrombosis. It is thought that retroviral gene therapy with this molecule could provide a treatment for PNH patients.

A final Ly6 superfamily member, the E48 antigen, is involved in intercellular adhesion between keratinocyte cells of the squamous epithelium. Such keratinocytes are attached to adjoining cells by large numbers of desmosomes, which are thought to play a role in the transition of transformed keratinocytes to metastatic tumor cells. Treatment with a monoclonal antibody raised against the E48 antigen has been successful in the eradication of residual, postoperative squamous cell carcinoma cells of the upper aerodigestive tract in several *in vivo* models and, to some degree, in humans. (van Dongen, et al., "Progress in radioimmunotherapy of head and neck cancer," Oncol. Rep. 1:259-264 (1994).) The gene encoding the E48 antigen has been mapped to the q24-qter region of human chromosome 8. Interestingly, a number of human diseases have been mapped to this region of chromosome 8 including Langer-Giedion syndrome, brachio-otorhinolaryngeal syndrome, trichorhinolaryngeal syndrome, and epidermolysis bullosa simplex.

A further example of a receptor family includes the prohibitin receptors. The prohibitin gene product is expressed in a wide variety of tissues and has been implicated as a component of a number of anti-proliferative mechanisms. The prohibitin gene encodes a 30 kD postsynthetically modified polypeptide located primarily in the mitochondria, but also may be associated with the IgM receptor on the B-cell plasma membrane. The protein functionally inhibits DNA synthesis and entry into S phase of the cell cycle by an unknown mechanism. Interestingly, although the prohibitin gene product is hypothesized to be involved in the maintenance of senescence and the prevention of cancer, one study found that, although somatic mutations in the prohibitin gene were present in a small number of breast cancers, no mutations were identified in any other breast, ovary, liver, and lung cancers examined. (Sato et al., Genomics 17:762-764 (1993).) However, the prohibitin gene has been mapped to human chromosome 17q12-21, the same region thought to contain the gene involved in sporadic breast cancer. Furthermore, DNA sequence analysis of the prohibitin gene identified somatic mutation in 4 of 23 cases of sporadic breast cancer examined. Thus, prohibitin family members may be involved in the development of cancer.

Moreover, the EGFR family of plasma membrane proteins are an integral component of normal cellular proliferation and in the pathogenesis of the cancerous state. The family is relatively small and includes the EGFR, c-erbB-2, c-erbB-3, and others. Various cancers are correlated with aberrant expression of one or more of these

10

15

20

25

30

35

genes. A number of ligands have been identified which bind to the EGFR-like receptors listed above including TGF-a, heparin-binding EGF, amphiregulin, criptoregulin, heregulin, and others. A large fraction of adenocarcinomas examined to date, especially those of the breast, colon, and pancreas, are typified by the amplification or overexpression of the c-erbB-2 gene. EGF, or an analogous ligand, initiates the cellular growth factor response by binding to the EGFR, or EGFR-related, receptor. Following the binding event, the receptor molecule dimerizes activating its intracellular tyrosine kinase domain. This event results in the phosphorylation of specific tyrosine residues near the carboxy terminus of the receptor. The diversity of signals able to be transduced through the relatively small number of EGFR-related receptor molecules is amplified considerably by the recent finding that EGFR-like receptor molecules can function when dimerized with other EGFR family members forming heterodimers.

Members of the EGFR-related family of integral membrane proteins have been implicated in the pathogenesis of a number of human disease-states. For example, a mutation in the EGFR itself appears to play an important role in the development of glioblastomas. (Sang et al., J. Neurosurg 82:841-846 (1995).) The EGFR gene is amplified or overexpressed in the majority of primary human glioblastomas. Although not conferring a distinct advantage on cell growth, an increase in EGFR expression was found to confer an increase in the ability of glioma cells to maintain anchorage-independent growth in soft agar especially in response to EGF and retinoic acid. Anchorage-independent growth *in vitro* correlates highly with tumorigenicity *in vivo*, therefore, it is likely that cells which express abnormally high levels of EGFR in human glioblastoma cells may be involved in the high potential for these cells to cause tumors *in vivo*.

Moreover, overexpression or amplification of c-erbB-2 has been reported to be involved in a high number adenocarcinomas, particularly of the breast, colon, and pancreas, and in a small proportion of ovarian carcinomas.

Thus, there is a clear need for identifying and exploiting novel members of the receptor families, such as those described above. Although structurally related, these receptors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such receptors may prove useful as therapeutics in an antitumor, diagnostic, or other capacity. Furthermore, receptors described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

SUMMARY OF THE INVENTION

5

10

15

20

25

30

35

In one aspect, the invention relates to receptor polypeptides and polynucleotides, as well as the methods for their production. Another aspect of the invention relates to methods for using such receptor polypeptides and polynucleotides. Such uses include the treatment of the specified diseases, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate receptor activity or levels.

DESCRIPTION OF THE INVENTION Definitions

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Receptor" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:Y, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said receptor including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said receptor.

"Receptor gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:X or allelic variants thereof and/or their complements.

"SEQ ID NO:X" comprises all or a substantial portion of the polynucleotide encoding each receptor of the invention. The value X for the nucleotide sequence is an integer specified in Table 1. This nucleotide sequence was translated into the receptor polypeptide identified in Table 1 as "SEQ ID NO:Y," where the value of Y for each receptor polypeptide is an integer defined in Table 1.

The invention further provides a composition of matter comprising a nucleic acid molecule which comprises a human cDNA clone identified by a cDNA Clone ID (Identifier) in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection ("ATCC") and given the ATCC Deposit Number shown in Table 1 for that cDNA clone. The ATCC is located at American Type Culture Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852, USA. The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that

WO 98/31799

required under 35 U.S.C. §112. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

PCT/US98/00959

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can

5

10

15

20

25

30

occur anywhere in a polypeptide, including the peptide backbone, the amino acid sidechains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphotidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis:

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to

Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.)

5

10

15

20

25

30

10

15

20

25

30

35

occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991.) While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073.) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., et al., Nucleic Acids Research (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. et al., J Molec Biol (1990) 215:403.)

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO:X is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: X. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence or anywhere between those terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:Y is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO:Y. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

20

25

10

5

Polypeptides of the Invention

In one aspect, the present invention relates to receptor polypeptides (or receptor proteins). The receptor polypeptides include the polypeptide of SEQ ID NO:Y; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:Y; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% identity to SEQ ID NO:Y are highly preferred. Also included within receptor polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% are highly preferred. Preferably receptor polypeptides exhibit at least one biological activity of the receptor.

30

The receptor polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

35

Fragments of the receptor polypeptides are also included in the invention. A "fragment" is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned receptor polypeptides. As with receptor polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most

10

15

20

25

30

35

preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of receptor polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of receptor polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus.

Also preferred are fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. The "domains" of each receptor polypeptide are illustrated in the Figures. The Figures compare SEQ ID NO:Y to the closest know homologue. Identical amino acids shared between the two polypeptides are shaded, while conservative amino acid changes are boxed. By examining the regions or amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. The amino acids sequences of SEQ ID NO:Y falling within these conserved domains are "fragments" and are specifically contemplated by the present invention. Especially preferred is the extracellular domains of a receptor of the invention. Soluble extracellular domains have antagonist activity mediated by competition with a receptor ligand.

Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain a biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala. Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

10

15

20

25

30

35

The receptor polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

Polynucleotides of the Invention

Another aspect of the invention relates to receptor polynucleotides. Receptor polynucleotides include isolated polynucleotides which encode the receptor polypeptides and fragments, and polynucleotides closely related thereto. More specifically, a receptor polynucleotide of the invention includes a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:X encoding a receptor polypeptide of SEQ ID NO:Y, and polynucleotide having the particular sequence of SEQ ID NO:X.

Receptor polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under receptor polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:X, or contained in the cDNA insert in the plasmid deposited with ATCC, to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, the receptor polynucleotide includes a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above receptor polynucleotides.

The receptors of the invention are structurally related to other proteins of specified receptor families, as shown by the results in the Figures. The cDNA sequence of SEQ ID NO:X encodes a polypeptide as described in Table 1 as SEQ ID NO:Y. Because the receptor polypeptides contain domains similar in structure to other receptor family members, the receptors of the present invention are expected to have,

inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1

5

Clone ID Name	SEQ ID NO:X	SEQ ID NO:Y	ATCC Deposit No.	ATCC Deposit Date	Receptor Family	Homology
HMACR70	11O.A	110.1	209054	05/16/97	Ig	Sialoadhesin
INVIACIO/O	1	18	#####	01/21/98	- 8	OB-1
HTEDK48			209054	05/16/97	TM4SF	MRC-OX44 PETA-3
1-1849 bp	2			·		
160-900 bp	3	19				
HTPED39 HPWAE25	4	20	209054 #####	05/16/97 1/21/98	TM4SF	NAG-2 TALLA-1
HTPEF86	5	21	209053	05/16/97	TM4SF	CD20 B1 Antigen
HSBBF02	6	22	209054	05/16/97	TM4SF	TALLA-1
HLTAH80	7	23	97242 209054	08/02/95 05/16/97	TM4SF	TALLA-1
HTPBA27	8	24	97242 209054	08/02/95 05/16/97	TM4SF	NAG-2
HAIDQ59			209054	05/16/97	TM4SF	CD9 Antigen
5' Sequence	9	25				
3' Sequence	10					
HHFEK40	11	26	209054	05/16/97	TM4SF	PETA-3
HGBGV89	12	27	209125 209054	06/09/97 05/16/97	TM4SF	L6H
HUVBB80	13	28	209054	05/16/97	TM4SF	L6
HJACE54	14	29	209053	05/16/97	Lectin	Galectin-3 Galectin-5 Galectin-8
HROAD63	15	30	209053	05/16/97	Ly6	E48 splice variant
HMWGS46	16	31	209053	05/16/97	Prohibitin	BAP-37
HNFGW06	17	32	209053	05/16/97	EGFR	EGFR

The novel full-length cDNA clone designated HMACR70 may be a member of the sialoadhesin family of the Ig superfamily of receptor-like molecules and a CD33 homologue. HMACR70 contains a 1497 nucleotide cDNA insert encoding a 315 amino acid ORF and was cloned from a GM-CSF-treated human macrophage cDNA library. The only additional cDNA libraries in the HGS database which include this clone are human eosinophils and possibly human gall bladder. A BLAST analysis of the amino acid sequence of HMACR70 demonstrates that this clone exhibits approximately 50% identity and 69% similarity over a 300 amino acids stretch of a gene termed human

10

differentiation antigen, and 38% identity and 62% similarity of the human myelin-associated glycoprotein precursor CD33 gene.

A more recent BLAST analysis confirms HMACR70's designation as a sialoadhesin family member. HMACR70 is homologous to two recently identified sialoadhesin family members, human OB binding protein (OB) 1 and 2. (See, Genbank Accession No. U71382; see Figure 1.) It is thought that OB-1 and OB-2 may bind leptin. Thus, HMACR70, as a sialoadhesin family member, may act to attenuate or even amplify intercellular routes of communication, including binding to leptin or modulating the activity of immune cells, such as macrophages. Clearly, any diseases affected by these processes could be treated by the polypeptide or fragment of HMACR70.

The full-length nucleotide sequences of ten novel human cDNA clones which potentially belong to the TM4SF superfamily are disclosed in the table above and will be addressed sequentially.

The cDNA clone HTEDK48 contains a 1849 nucleotide cDNA insert encoding a 245 amino acid ORF that was cloned from a human testes cDNA library. The coding sequence of HTEDK48 (SEQ ID NO: 3) may be fused to other human proteins, such as 3-hydroxyacyl-CoA dehydrogenase. BLAST analysis of the amino acid sequence of HTEDK48 demonstrates that this clone exhibits approximately 30% identity and 51% similarity over a 245 amino acid stretch of the CD82 molecule. Recent studies have shown that CD82 can associate with CD4 or CD8 and deliver costimulatory signals for the TCR/CD3 pathway. CD82 has also been found to be involved in syncytium formation in HTLV-I-infected T-cells. And finally, in a recently published study in which the expression of the CD82 gene by tumors of the lung was examined retrospectively, it was reported that CD82 may be linked to the suppression of tumor metastasis of prostate cancer. The study also reported that decreased CD82 expression may be involved in malignant progression of such cancers. Thus, HTEDK48 may also be involved in the development of cancer.

A more recent BLAST analysis shows that HTEDK48 is homologous the rat leukocyte antigen, MRC OX-44, and the platelet endothelial tetraspan antigen -3 (PETA-3). (See Figure 2X.) MRC OX-44, a member of a new family of cell surface proteins, appears to be involved in growth regulation. (See, Bellacosa, A., et al., "The Rat Leukocyte antigen MRC OX-44 is a Member of a New Family of Cell Surface Proteins which Appear to be Involved in Growth Regulation," Mol. Cell. Bio. 11: 2864-2872 (1991).) Similarly, PETA-3 has been located to platelet endothelial cells, and an anti-PETA-3 antigen monoclonal antibody can stimulate platelet aggregation and mediator release. (See, Fitter, S., "Molecular Cloning of cDNA Encoding a Novel Platelet-Endothelial Cell Tetra-Span Antigen, PETA-3," Blood, 86(4):1348-1355 (1995).) Thus, HTEDK48 may function similar to MRC OX-44 or PETA-3 to affect

5

10

15

20

25

30

WO 98/31799 PCT/US98/00959

growth of blood cells. Administering polypeptides or fragments of HTEDK48 may be an effective treatment of blood disorders.

The cDNA clone **HPWAE25** contains a 1288 nucleotide cDNA insert encoding a 273 amino acid ORF that was cloned from a human pancreas tumor cDNA library, while clone **HTPED39** represents a truncated cDNA sequence. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including keratinocytes, ulcerative colitis, striatum depression, lymph node breast cancer, ovarian cancer, stage B2 prostate cancer, kidney medulla, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human cell lines including U937, MM96, WM115, and MDAMB231. A BLAST analysis of the amino acid sequence of HTPED39 demonstrates that this clone exhibits approximately 35% identity and 50% similarity over the entire length of the CD37 molecule. The CD37 antigen is expressed on B cells and on a subpopulation of T cells, but not on pre-B or plasma cells. It has been reported that CD37 expression is downregulated in conjunction with B-cell activation, suggesting that CD37 may be involved in the processes which dictate the activation state of the B-cell.

Moreover, HPWAE25 is also homologous to recently identified TM4SF members, NAG-2 and TALLA-1. (See Figure 3.) NAG-2 is thought to complex with integrins and other TM4SF proteins, while TALLA-1 is a highly specific marker of T-cell acute lymphoblastic leukemia and neuroblastoma. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that Complexs with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997); Takagi, S., "Identification of a Higly Specific Surface Marker of T-cell Acute Lymphoblastic Leukemia and Neuroblastoma as a New Member of the Transmembrane 4 Superfamily," Int. J. Cancer 61(5):706-715 (1995).) Thus, HPWAE25 may be involved the development of cancer, particularly leukemia, lymphoma, and neuroblastoma. HPWAE25 may be used as an effective treatment of these cancers, as well as a diagnostic marker.

A subfamily of TM4SF receptors include CD20 proteins. A CD20-like cDNA clone was obtained from a human pancreas tumor cDNA library and contains a 1236 nucleotide insert which encodes a 250 amino acid ORF. A BLAST analysis of the deduced amino acid sequence of HTPEF86 exhibits approximately 41% identity and 61% similarity to the CD20 gene, also known as B1 antigen. (See Figure 4.) Expression of this gene is detected in only two additional HGS human cDNA libraries; amygdala depression and 9 week early stage human. Although the precise functional role of CD20 has yet to be determined, it is clear that CD20 plays a key role in the regulation of B-cell activation. Based primarily on sequence identity, the novel CD20-like molecule presented herein may also be involved in cell cycle activation. Potential therapeutic and/or diagnostic applications for HTPEF86 may include such clinical

5

10

15

20

25

30

presentations as juvenile rheumatoid arthritis, Graves' Disease, and a number of B-cell lymphomas or other lymphoid tumors.

PCT/US98/00959

The clone **HSBBF02** contains a 1115 nucleotide cDNA insert encoding a 245 amino acid ORF and was cloned from an HSC 172 cell line cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including brain amygdala depression, endothelial cells, fetal liver and heart, osteoblasts, testes, and others. A BLAST analysis of the amino acid sequence of HSBBF02 demonstrates that this clone exhibits approximately 64% identity and 80% similarity with the A15 molecule over a 131 amino acid stretch (A15 is composed of 244 amino acids). A more recent BLAST search shows that HSBBF02 is similar to the TALLA-1 protein and may in fact be a closely related family member. (See Figure 5.)

In addition, a second cDNA clone, designated HLTAH80, exhibits sequence similarity to the A15 molecule and TALLA-1. (See Figure 6.) This clone contains a 1662 nucleotide cDNA insert encoding a 253 amino acid ORF and was cloned from a human T-cell lymphoma cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including B-cell lymphoma, corpus collosum, endometrial tumor, osteosarcoma, testes, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human tissues including spleen, lymph node, thymus, PBLs, heart, and a particularly strong signal in skeletal muscle and pancreas. A BLAST analysis of the amino acid sequence of HLTAH80 demonstrates that this clone exhibits approximately 35% identity and 55% similarity over the entire length of the A15 molecule.

Since expression of A15 drops to undetectable levels when comparing immature T-cells to peripheral blood lymphocytes, it is thought that A15 may play a role in the development of T-cells. Furthermore, the MXS1(CCG-B7) gene which codes for A15 contains a number of triplet nucleotide repeats which have been associated with neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome, and myotonic dystrophy. In addition, A15 appears to be expressed exclusively on T-cell acute lymphoblastic leukemia cell lines, including several derived from adult T-cell leukemia and those established by immortalization with human T-cell leukemia virus type 1 or Herpesvirus saimiri. Thus, clones HLTAH80 and/or HSBBF02 may also be involved in diseases caused by the expansion of repeats or chromosomal instability.

The cDNA clone HTPBA27 contains a 1345 nucleotide cDNA insert encoding a 238 amino acid ORF and was cloned from a human tumor pancreas cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including cerebellum, breast lymph node, osteosarcoma, adult testes, RS4;11 bone marrow cell line, microvascular endothelial cells, and others. A BLAST analysis of the amino acid sequence of HTPBA27 demonstrates that this

5

10

15

20

25

30

clone exhibits approximately 40% identity and 64% similarity with a glycoprotein termed CD53 over its entire length. CD53 is thought to be involved in thymopoiesis, since rat CD53 can be detected on immature CD4-8-thymocytes and the functionally mature single-positive subset, but cannot be detected on the intermediate CD4+8+ The CD53 molecule has also been implicated as a thymocytic subset of cells. component of signal transduction pathways in B cells, monocytes and granulocytes, rat macrophages, NK, and T cells. Moreover, as illustrated in Figure 7, HTPBA27 was recently confirmed as a TM4SF receptor. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997).) Calling the HTPBA27 polypeptide NAG-2, this group confirmed HTPBA27's status as a TM4SF receptor by showing that NAG-2 complexes with integrin and other TM4SF receptors. diseases caused by the failure of HTPBA27 to complex with integrin and other TM4SF receptors can be treated by administering HTPBA27. HTPBA27 can also be used to diagnose these diseases.

PCT/US98/00959

The cDNA clone **HAIDQ59** contains cDNA insert encoding a 221 amino acid ORF that was cloned from a human epithelial cell induced with TNFa and INF cDNA library. The 5' end of HAIDQ59 is represented by the SEQ ID NO: 9, while the 3' end is represented by SEQ ID NO: 10. This clone appears in only two additional cDNA libraries in the HGS database. These two libraries were constructed from the human Jurkat T-cell line and human microvascular endothelial cells. A BLAST analysis of the amino acid sequence of HAIDQ59 demonstrates that this clone exhibits approximately 53% identity and 69% similarity over 226 amino acids of the CD9 TM4SF molecule. (See Figure 8.) It has been demonstrated that the CD9 molecule is involved in signal transduction pathways in platelets, as well as in cell adhesion in both platelets and pre-B-cell lines. Intriguingly, a monoclonal antibody (vpg15), which recognizes the feline homologue of CD9, has been shown to block infection by feline immunodeficiency virus (FIV). Furthermore, a recent study shows that cells expressing high levels of CD9 exhibited suppressed cell motility. Thus, HAIDQ59 may also be involved in signal transduction of blood cells.

The cDNA clone **HHFEK40** contains a 936 nucleotide cDNA insert encoding a 252 amino acid ORF and was cloned from a human fetal heart cDNA library. This clone appears once in the human fetal heart cDNA library and possibly in a hemangiopericytoma cDNA library. A BLAST analysis of the amino acid sequence of HHFEK40 demonstrated that this clone exhibits approximately 60% identity and 75% similarity over the entire length of a molecule designated PETA-3. (See Figure 9.) PETA-3 was originally identified as a novel human platelet surface glycoprotein termed gp27. Although PETA-3 is present in low abundance on the platelet surface, an anti-PETA-3 monoclonal antibody can stimulate platelet aggregation and mediator release.

5

10

15

20

25

30

10

15

20

25

30

35

Thus, HHFEK40 may function similar to PETA-3 to affect growth of blood cells. Administering polypeptides or fragments of HHFEK40 may be an effective treatment of blood disorders.

The cDNA clone HGBGV89 contains a 738 nucleotide cDNA insert encoding a 197 amino acid ORF and was cloned from a human gall bladder cDNA library. The only two additional appearances of this clone in the HGS database are in a normalized fetal liver cDNA library and in a fetal liver/spleen cDNA library. The cDNA clone HUVBB80 contains a 1071 nucleotide cDNA insert encoding a 201 amino acid ORF and was cloned from a human umbilical vein cDNA library. This clone appears in several additional cDNA libraries in the HGS database including prostate BPH, thyroid, and fetal liver/spleen. BLAST analyses of the amino acid sequences of HGBGV89 and HUVBB80 demonstrate that these clones exhibit approximately 49% identity and 65% similarity and 47% identity and 68% similarity, respectively, over the entire length of a molecule designated L6 surface protein or human tumor-associated antigen L6. (See Figures 10 & 11.) Moreover, another group has confirmed the TM4SF receptor homology of HGBGV89 by describing the protein as a putative transmembrane protein L6H. (See Genbank Accession No 2587054; see Figure 10.) The L6 cell surface antigen is highly expressed on lung, breast, colon, and ovarian carcinomas. Promising results of phase 1 clinical studies have been reported with an anti-L6 monoclonal antibody, or its humanized counterpart, suggesting that the L6 antigen may be an attractive target for monoclonal antibody-based cancer therapy.

In summary, there is a clear need for identifying and exploiting novel members of the TM4SF superfamily such as those described herein. Although structurally related, these factors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules, such as the novel potential members of the TM4SF superfamily detailed here, should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such factors may prove useful as therapeutics in an anti-tumor, diagnostic, or other capacity. Furthermore, factors such as the nine novel TM4SF superfamily-like molecules described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

Besides TM4SF receptors, receptors from other families are also described. For example, clone **HJACE54**, also called galectin 11, exhibits significant sequence identity to the rat galectin 5, the chicken galectin 3 gene, and the human galectin 8 genes. (See Figure 12.) The galectin 11 cDNA clone contains an 865 nucleotide insert which encodes a 133 amino acid ORF. The clone was obtained from a Jurkat T-cell G1 phase cDNA library. A BLAST analysis of the deduced amino acid sequence of HJACE54 demonstrates approximately 35% identity and 57% similarity to the amino

WO 98/31799 PCT/US98/00959

acid sequence of the rat galectin 5 gene. Expression of galectin 11 is quite limited in the HGS database. In fact, the only two additional ESTs in the HGS database which contain the HJACE54 sequence were found in human neutrophil and human infant adrenal gland cDNA libraries. Northern blot analyses have not been performed to examine expression patterns of the galectin 11 gene.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1mediated disruption of laminin-integrin a₇b₁ interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively. As a result, the discovery of a novel galectin, such as that encoded by HJACE54, is likely to be a valuable asset both diagnostically and therapeutically.

Additionally, a full-length nucleotide sequence of a novel human cDNA clone which encodes an apparent splice variant of the previously described human E48 antigen has recently been determined. (See Figure 13.) Clone HROAD63 contains a 441 nucleotide cDNA which encodes a 70 amino acid polypeptide. This novel clone exhibits significant sequence identity to several members of a relatively new family of cell-surface proteins termed the Ly6 superfamily. These members include murine and human SCA-2, rat Ly-6 (also termed ThB), and human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)]. The novel E48 splice variant was obtained from the HGS human stomach cDNA library. The clone is present in only a limited number of other HGS cDNA libraries including kidney cancer, keratinocyte, and tongue. An alignment of the nucleotide sequences of the human E48 and HROAD63 cDNAs demonstrates that the initial 168 and 178 nucleotides of E48 and HROAD63, respectively, are identical, with the exception of an additional 10 nucleotides of sequence at the extreme 5' end of the HROAD63 sequence. The

5

10

15

20

25

30

10

15

20

25

30

35

sequence of the two clones is also identical for an additional 229 nucleotides including the 3' end of the coding sequences and the entire 3' untranslated regions. The only divergence of nucleotide sequence in this region of the clones is the deletion of a single thymidine residue in the 3' UTR of the E48 cDNA. The major difference between the two nucleotide sequences is a 329 nucleotide deletion from the HROAD63 sequence. This deletion causes a shift in the HROAD63 reading frame and encompasses the translational stop signal used in the E48 clone. As a result, the carboxy terminal sequence of HROAD63 is radically altered with regard to that of E48 (as illustrated in Figure 13 by the obvious differences between amino acids 56-128 of E48 and 56-70 of HROAD63 in the amino acid alignment). The clinical presentation of disorders, including abnormal skin and hair phenotypes, may be attributed, at least in part, to a non-functional Ly6 superfamily member such as E48 or HROAD63. HROAD63 may also be involved in blood disorders, as seen with its homologues SCA-2 and CD59.

20

A novel prohibitin cDNA clone presented herein was originally identified in a human bone marrow cell line (RS4;11) cDNA library. The clone contains a 1066 nucleotide insert which encodes a 299 amino acid polypeptide. BLAST and BestFit analyses of the predicted amino acid sequence of HMWGS46 demonstrate a highly significant sequence identity to a murine protein termed IgM B-cell receptor associated protein (BAP)-37 (Genbank accession number X78683). The HMWGS46 amino acid sequence exhibits nearly perfect identity and similarity over the entire length of the murine BAP-37 sequence. (See Figure 14.) In addition, the full-length nucleotide sequences of HMWGS46 and BAP-37 exhibit at least 87% identical. The HMWGS46 clone also exhibits approximately 49% sequence identity and 85% sequence similarity to a human gene designated prohibitin. Finally, the HMWGS46 cDNA appears in a substantial number of HGS human cDNA libraries in addition to the bone marrow cell line cDNA library from which it was cloned. Some of the cDNA libraries in which this clone appears include keratinocytes, induced endothelial cells, activated neutrophils, synovial sarcoma, colon carcinoma cell line, Jurkat cell line membrane bound polysomes, epileptic frontal cortex, primary dendritic cells, and a number of others. The novel gene related to prohibitin and BAP-37 may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event. Thus, although the precise functional role of the prohibitin family members are less than clear, it is quite likely that such homologues are involved in such complex processes as development, senescence, and tumor suppression. Therefore a novel gene, such as HMWGS46, may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event.

A human cDNA clone encoding a novel epidermal growth factor receptor (EGFR)-like molecule is also disclosed. The novel EGFR-like cDNA clone presented herein was originally identified in an activated human neutrophil cDNA library. The

WO 98/31799

clone contains a 704 nucleotide insert which encodes a 168 amino acid polypeptide. A BLAST analysis of the predicted amino acid sequence of HNFGW06 demonstrates that this novel clone exhibits approximately 85% identity and 90% similarity to a protein designated epidermal growth factor receptor-related protein [Homo sapiens]. (See Figure 15.) The expression profile of the HNFGW06 clone in the HGS database indicates the existence of a fairly highly restricted expression pattern. In addition to the activated neutrophil library from which this clone was obtained, it also appears in the following HGS human cDNA libraries: synovial sarcoma, smooth muscle, placenta, and possibly primary dendritic cells.

PCT/US98/00959

The novel EGFR-like cDNA clone HNFGW06 may lead to a number of exciting possibilities for therapeutic and/or diagnostic treatments or reagents. For example, HNFGW06 may be involved in the onset of human breast cancers as well. In addition, due to the fact that TGF-a acts through binding to the EGFR, it is possible that HNFGW06 may also play a role in a variety of gastric processes including regulation of acid secretion, regulation of mucous cell growth, and protection against ethanol- and aspirin-induced injury to gastric tissues.

GENERATING POLYNUCLEOTIDES

Polynucleotides of the present invention encoding a receptor may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells specified in Table 1 using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174.) Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding a receptor polypeptide of SEQ ID NO:Y may be identical to the polynucleotide encoding SEQ ID NO:Y, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:Y.

When the polynucleotides of the invention are used for the recombinant production of a receptor polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in

5

10

15

20

25

30

10


15

20

25

30

35



Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding receptor variants comprising the amino acid sequence of receptor polypeptide of Table 1 (SEQ ID NO:Y) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:X or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC, or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding the receptor and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs) that have a high sequence similarity to the receptor gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding the receptor polypeptide, including homologs and orthologs from other species, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO:X or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

WO 98/31799 PCT/US98/00959

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

5 Vectors, Host Cells, Expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (supra).

10

15

20

25

30

10

15

20

25

30

35

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the receptor polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the receptor polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Receptor polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

This invention also relates to the use of receptor polynucleotides or polypeptides for use as diagnostic reagents. Detection of a mutated form of the receptor gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from underexpression, over-expression or altered expression of the receptor. Individuals carrying mutations in the receptor gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled receptor nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. (See, e.g., Myers et al., Science (1985) 230:1242.) Sequence changes at specific locations may also be revealed by

10

15

20

25

35

nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. (See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401.) In another embodiment, an array of oligonucleotides probes comprising receptor nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996).)

The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutation in the receptor gene by the methods described.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of receptor polypeptide or receptor mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises:

- (a) a receptor polynucleotide, preferably the nucleotide sequence of SEQ ID NO:X, or a fragment thereof;
 - (b) a nucleotide sequence complementary to that of (a);
- (c) a receptor polypeptide, preferably the polypeptide of SEQ ID NO:Y, or a fragment thereof; or
- (d) an antibody to a receptor polypeptide, preferably to the polypeptide of SEQ ID NO: Y.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the

sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10

15

20

25

5

Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the receptor polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the receptor polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole et al., MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

30

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against receptor polypeptides may also be employed to treat diseases.

Vaccines

35

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a receptor polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from a disease. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which

10

15

20

25

30

35

comprises, delivering a receptor polypeptide via a vector directing expression of the receptor polynucleotide in vivo in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a receptor polypeptide wherein the composition comprises a receptor polypeptide or receptor gene. The vaccine formulation may further comprise a suitable carrier. Since a receptor polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation instonic with the blood of the recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Screening Assays

The receptor polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).

The receptor polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate the receptor on the one hand and which can inhibit the function of the receptor on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions and diseases. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions and diseases.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof.

10

15

20

25

30

35

Such cells include cells from mammals, yeast, Drosophila or E. coli. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a receptor polypeptide to form a mixture, measuring receptor activity in the mixture, and comparing the receptor activity of the mixture to a standard.

The receptor cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of receptor mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of receptor protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of the receptor (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential receptor antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the receptor, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for receptor polypeptides; or compounds which decrease or enhance the production of receptor, which comprises:

- (a) a receptor polypeptide, preferably that of SEQ ID NO:Y;
- (b) a recombinant cell expressing a receptor polypeptide, preferably that of SEQID NO:Y:
 - (c) a cell membrane expressing a receptor polypeptide; preferably that of SEQ ID NO: Y; or
 - (d) antibody to a receptor polypeptide, preferably that of SEQ ID NO: Y.

SUBSTITUTE SHEET (RULE 26)

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

PCT/US98/00959

Prophylactic and Therapeutic Methods

5

10

15

20

25

30

35

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of receptor activity.

If the activity of the receptor is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking the binding of ligands to the receptor or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of the receptor polypeptides still capable of binding the ligand in competition with endogenous receptor may be administered. Typical embodiments of such competitors comprise fragments of the receptor polypeptide.

In still another approach, expression of the gene encoding endogenous receptor can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. (See, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Alternatively, oligonucleotides which form triple helices with the gene can be supplied. (See, for example, Lee et al., Nucleic Acids Res (1979) 6:3073; Cooney et al., Science (1988) 241:456; Dervan et al., Science (1991) 251:1360.) These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

For treating abnormal conditions related to an under-expression of the receptor and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates the receptor, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of the receptor by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and

15

20

25

30

35

other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

5 Formulation and Administration

Peptides, such as the soluble form of receptor polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of $0.1\text{-}100~\mu\text{g/kg}$ of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

SUBSTITUTE SHEET (RULE 26)

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

SUBSTITUTE SHEET (RULE 26)

Page 1

HMACR 70	
GCAGTTCCTGAGAGAAGAACCCTGAGGAACAGACGTTCCCTCGCGGCCCTGGCACCTCCAACCCCAGATATGCTGCTGCTGCTGCTGCTG	~
CGTCAAGGACTCTCTTCTTGGGACTCCTTGTCTGCAAGGGAGCGCCGGGACCGTGGAGGTTGGGGTCTATACGACGACGACGACGACGAC	90
MILLLL	
	180
GGGGACGAGACCCCCTCCCCTCCCACCTTCCTGTCTTCTCATTGGCCTTCCTAATGAGCGACTGCTACGTCTCAAGGCACTGGCACGTT	
PLLWGRERVEGOKSNRK DYSLTMOSSV TV O	
GAGGGCATGTGTGCCATGTGCGCTGCTCCTTCTCCTACCCAGTGGACAGCCAGACTGACT	270
CTCCCGTACACACAGGTACACGCGACGAGGAAGAGGATGGGTCACCTGTCGGTCTGACTGA	170
E G M C V H V R C S F S Y P V D S Q T D S D P V H G Y W F R	
GCAGGGAATGATATAAGCTGGAAGGCTCCAGTGGCCACAAACAA	360
CGTCCCTTACTATATTCGACCTTCCGAGGTCACCGGTGTTTGTT	
A G N D I S W K A P V A T N N P A W A V Q E E T R D R F H L	
CTTGGGGACCCACAGACCAAAAATTGCACCCTGAGCATCAGAGATGCCAGAATGAGTGATGCGGGGGAGATACTTCTTTCGTATGGAGAAA	450
GAACCCCTGGGTGTCTGGTTTTTAACGTGGGACTCGTAGTCTCTACGGTCTTACTCACTACGCCCCTCTATGAAGAAAGCATACCTCTTT	
L G D P O T K N C T L S I R D A R M S D A G R Y F F R M E K	
GGAAATATAAAATGGAATTATAAATATGACCAGCTCTCTGTGAACGTGACATACCCTCCTCAGAACTTGACTGTGACTGTCTTCCAAGGA	540
CCTTTATATTTTACCTTAATATTTATACTGGTCGAGAGACACTTGCACTGTATGGGAGGAGTCTTGAACTGACACTGACAGAAGGTTCCT	
G N 1 K W N Y K Y D Q L S V N V T Y P P Q N L T V T V F Q G	
GAAGGCACAGCATCCACAGCTCTGGGGAACAGCTCATCTCTTTCAGTCCTAGAGGGCCAGTCTCTGCGCTTGGTCTGTGCTGTTGACAGC	630
CTTCCGTGTCGTAGGTGTCGAGACCCCTTGTCGAGTAGAGAAAGTCAGGATCTCCCGGTCAGAGACGCGAACCAGACACCAGACAACTGTCG	
EGŢASTALGNSSSLSVLEGOSLRLVCAVDS	
AATCCCCCTGCCAGGCTGAGCTGGACCTGGAGGAGTCTGACCCTGTACCCCTCACAGCCCTCAAACCCTCTGGTACTGGAGCTGCAAGTG	720
TTAGGGGGACGGTCCGACCTGGACCTCCTCAGACTGGGACATGGGGAGTGTCGGGAGTTTGGGAGACCATGACCTCGACGTTCAC	720
N P P A R L S W T W R S L T L Y P S Q P S N P L V L E L Q V	
CACCTGGGGGATGAAGGGGAATTCACCTGTCGAGCTCAGAACTCTCTGGGTTCCCAGCACGTTTCCCTGAACCTCTCCCTGCAACAGGAG	810
GTGGACCCCTACTTCCCCTTAAGTGGACAGCTCGAGTCTTGAGAGACCCAAGGGTCGTGCAAAGGGACTTGGAGAGGGACGTTGTCCTC	,,,

HMACRIC Y T G K M K P V S G V L L G A V G G T G A T A L V F L S F C. GTCATCTTCATTGTAGTGAGGTCCTGCAGGAAGAATCGGCAAGACCAGCAGCGGACGTGGGAGACATAGGCATGAAGGATGCAAACACC CAGTAGAAGTAACATCACTCCAGGACGTCCTTCTTTAGCCGTTCTGGTCGTCGCCTGCACCCTCTGTATCCGTACTTCCTACGTTTGTGG VIFIVVRSCRKKSARPAADVGDIGMKDANT ATTCAGGGGCTCAGCCTCTCAGGGTAACTGGATGAGTCCTGGGCAGATGATAACCCCCGACACCATGGCCTGGCTGCCCACTCCCTCAGG TAAGTCCCCGAGTCGGAGAGTCCCATTGACCTACTCAGGACCCGTCTACTATTGGGGGCTGTGGTACCGGACCGACGGGTGAGGGAGTCC 106 L S L S G (SEGIDNO:18) GGAGGAAAGAGAGATCCCAGTATGCACCCCTCAGCTTTCATAAGGGGGAGCCTCAGGACCTATCCAGGTCAAGAAGCCACCAACAA**TGAG** CCTCCTTCTCTCTAGGGTCATACGTGGGGAGTCGAAAGTATTCCCCCTCGGAGTCCTGGATAGGTCCAGTTCTTCGGTGGTTGTTACTC TACTCAGAGATCAAGATCCCCAAGTAAGAAAATGCAGAGGCTCGGGCTTGTTTGAGGGTTCACGACCCCTCCAGCAAAGGAGTCTGAGGC 1260 ATGAGTCTCTAGTTCTAGGGGTTCATTCTTTTACGTCTCCGAGCCCGAACAACTCCCAAGTGCTGGGGAGGTCGTTTCCTCAGACTCCG CCAAACTCTCCCTTTCCCCATCCAATCGGTCCACACTCCCCGCCCTGGCCTCTGTACCCACCATTCTCCTCTGTACTTCTCTAAGGATGA (SeQ JONO:1)

Page 1

																									\mathcal{C}					
CG	CAC	GAG	ŢGG	ACA	ACC	ATC	AGG	GAG	CCA	GGA	CAC	AGA	GGG	GCA	GAG	CĄA	GTC	AGC	ATT	GGC	GCC	CCT	ŢCC	TCA	GAT	ccc	TAT	CAT	CTT	
GCCGTGCTCACCTGTTGGTAGTCCCTCGGTCCTGTGTCTCCCCGTCTCGTTCAGTCGTAACCGCGGGGAAGGAGTCTAGGGATAGTAGAA												GAA	90																	
$oldsymbol{igwedge}$																														
			-i		ÀGA	- +		+	-+				+			\div			+									+		180
CCI	TTT	GTC	ATC	GGG	TCT	CCA	AGT	CCT	TCT	ACA	ATT	GAA	TTT	ACA	AGC	ccc	ACG	GGG	TCA	GAC	AAG	TCG	TAC	CGA	CTT	TAG	GTG	TGA	GGC	
																							М	А	Ε	i	н	T	Р	
TA.	TCT	TCC	TTG	AAG	AAA	CTG	TTA	тст	TTA	стс	AAT	GGC	TTC	GTG	GCT	GTG	TCT	GGC.	ATC.	ATC	CTA	GTT	GGC	CTG	GGC.	ATT	GGT	GGT	AAA	
					TTT								+					+	-+				+	-+					-+	270
														0,10			,,,,,,		.,,,		U 7111			u c					• • •	
Y	S	S	L	K	K	L	L	S	L	L	N	G	F	٧	Α	٧	S	G	1	I	L	٧	G	L	G	į	G	G	K	
TG	TGGA	GGG	ĠCC	TCT	CTG	AÇG.	AAT	GTC	CTC	GGG	CTG	TCC	TCC	GCA	TAC	CTC	стт	CACI	GTT	GGC.	AAC	CTG	TGC	CTG	GTG.	ATG	GGA	TGC	ATC	
ACA	CCT	ccc	CGG	AGA	GAC	TGC	TTA	CAG	GAG	CCC	GAC	AGG	AGG	CGT	ATG	GAG	GAA	GTG	CAA	CCG	TTG	GAC.	ACG	GAC	CAC	TAC	ССТ	ACG1	ΓAG	360
С	G	G	А	S	L	T	N	٧	L	G	L	S	S	А	Y	L	L	н	٧	G	N	L	С	L	٧	м	G	С	ī	
		C T C	c + +	000		000	200																							
			÷		TGT	-+-			+				+			-			+				;							450
160	LAI	GAC	GAA	CCG	ACA	CGG	CCC.	ACC.	AIA	CCT	CGC	TGA	TTT	CTE	TCG	TCT	CCG	TGC	GAG	AAC.	AAA	ACG	TAG	GAC.	AGT	TAC	CAA'	TAAC	AG	
T	٧	L	L	G	C	Α	G	W	Υ	G	A	T	Κ	Ε	S	R	G	T	L	L	F	С	I	L	S	M	٧	ı	٧	
сто	ATC	ATG	GAA	GTT	ACA	GCT	GCC.	ACA	GTG	GTC	CTT	стт	TTC	TTT	CCA	ATT	GTT	GGAG	SAT	STG	GCCI	rtge	GAA	CAC	ACC.	TTC	STG	4000	TG	
			+		TGT								+			+		 -	+				!							540
	,	м	_	v	τ.	٨		7	v				_	_				^	_				_		+	_		·	•	
-	•	11	C	٧	1	A	A	•	٧	٧	L	L	r	۲	٢	1	٧	G	U.	٧	А	L	=	н	ŀ	r	٧	ı	L	
AGG	AAG	AAT	TAC	AGA	GGT	TAC	AAC	GAG	CCA	GAC	GAC	TAT		ACA				TTG	GTC	ATG	GAGA	AGG	TA.	AAG	TGC	TGT	GGG	GTG/	AAT	630
TCC	TTC	TTA	ATG	TCT	CCA	ATG	FTG	CTC	GGT	CTG	CTG.	ATA.						AAC	CAG	TAC	CTCI	TTC	GAT	TTC	ACG/	ACA	ccc	CACT	TÀ	-
R	K	N	Y	R	G	Υ	N	Ε	Р	D	D	Υ	S	T	Q	W	N	L	٧	M	Ε	ĸ	L	κ	С	С	G	٧	N	
4 A C	TAC	A C A	CAT	.	TCT	200	· C T	T.C.C.	TTC/	~														***		201				•
			+			 -			-+							+													+	720
116	АІЬ	161	LIA	ААД.	AGA	. .	46A/	4667	ады	- 1 1	IAL	161	IGU	LLGI	ا ا	1667	A 1 G (افاقاة	ICL	ICAA	AUGA	ICA I	1 1 1 1	AGG	IAGO	LLI	ICAL	.ALA	166	
N	Υ	Ţ	D	F	S	G	S	S	F	Ε	M	T	1	G	Н	T	Y	Ρ	R	S	С	С	K	S	l	G	S	٧	S	
TGT	GAC	GGA	CGC	GAT	GTG.	тсто	CA	AAC	GTC	ATC	CAC	CAG							TCC	TAA	AAAA	TCA	ACE/	AAG	ACTO	CAGA	AGC1	TCA		
ACA	CTG	CCT	GCG	CTA	CACA	AGAG	GT	TTG	CAG	rago	STG	GTC	TTC	CCG	ACA.	AAG(STAT	TTC	AGO	SAT	TTT	AGI	rgg'	TTC.	r G A C	STCT	CGA	AGT	GG	810
С	D	G	R	D	٧	S	Р	N	٧	ı	н	0	ĸ	G	c	F	н	K	,	,	ĸ	ı	ī	ĸ	т	ß	S	F	т	
																													150	CID NO:3
					CTGC																									900
					GAC																									
L	s	G	s	s	L	G	Α	A	٧	i	a	R	W	G	s	R	Y	٧	Α	Q	Α	G	L	Ε	L	L	A	. 6	EG	ID
																													NO	D: 19/

HIEDK TO	
GATCCCCCCGCCTAGGCCTCCCAAAGTGCTGGGTTTACCAGCGTGAGCCACCACGCTGGGCTTCCTGCATCCTTTTAAGGTTCCTGAGGG	
CTAGGGGGGGGGATCCGGAGGGTTTCACGACCCAAATGGTCGCACTCGGTGGTGCGACCCGAAGGACGTAGGAAAATTCCAAGGACTCCC	990
TCTGCCTGAGAGGÀGCTGTCCCTGAATCTCCATGCAGCCCCACCTGCCACATCACCAAGACATCTTTGCCAGCAACACTTCCTCC	· 1080
AGACGGACTCTCCTCGACAGGGACTTAGAGGTACGTCGGGGTGGACGGTGTAGTGGTTCTGTATGTTAGAAACGGTCGTTGTGAAGGAGG	
TTGCAGATTACAAGCATAGCTAATGCCACCACCAGACAAGACCGATTCGCTGGCCTCCATTTCTTCAACCCAGTGCCTGTCATGAAACTT	. 1170
AACGTCTAATGTTCGTATCGATTACGGTGGTCGGTCTGTTCTGGCTAAGCGACCGGAGGTAAAGAAGTTGGGTCACGGACAGTACTTTGAA	
GTGGAGGTCATTAAAACACCAATGACCAGCCAGAAGACATTTGAATCTTTGGTAGACTTTAGCAAAACCCTAGGAAAGCATCCTGTTTCT	1260
CACCTCCAGTAATTTTGTGGTTACTGGTCGGTCTTCTGTAAACTTAGAAACCATCTGAAATCGTTTTGGGATCCTTTCGTAGGACAAAGA	_
TGCAAGGACACTCCTGGGTTTÁTTGTGAACCGCCTCCTGGTTCCATACCTCATGGAAGCAATCAGGCTGTATGAACGAGGGCCTCCTGGC	1350
ACGTTCCTGTGAGGACCCAAATAACACTTGGCGGAGGACCAAGGTATGGAGTACCTTCGTTAGTCCGACATACTTGCTCCCGGAGGACCG	
TTTCCCTGTGGGCTTCTGAGAAAGGTTTCTGGAACTCCCACCACCACCACCACCACCAGGCCAGAGCAATTGCATGGCCGGCC	- 1440
AAAGGGACACCCGAAGACTCTTTCCAAAGACCTTGAGGGTGGTGGGGGGTGATGTCAGGGTCGGTC	
GATATCCTGGATCTCTGCTTTTGATTAAAAGGTGACGCATCCAAAGAAGACATTGACACTGCTATGAAATTAGGAGCCGGTTACCCCATG	; - 1530
CTATAGGACCTAGAGACGAAAACTAATTTTCCACTGCGTAGGTTTCTTCTGTAACTGTGACGATACTTTAATCCTCGGCCAATGGGGTAC	
GGCCCATTTGAGCTTCTAGATTATGTCGGACTGGATACTACGAAGTTCATCGTGGATGGGTGGCATGAAATGGATGCAGAGAACCCATTA	- 1620
CCGGGTAAACTCGAAGATCTAATACAGCCTGACCTATGATGCTTCAAGTAGCACCTACCCACCGTACTTTACCTACGTCTCTTGGGTAAT	
CATCAGCCCAGCCCATCCTTAAATAAGCTGGTAGCAGAGAACAAGTTCGGCAAGAAGACTGGAGAAGGATTTTACAAATACAAGTGATGT	- 1710
GTAGTCGGGTCGGGTAGGAATTTATTCGACCATCGTCTCTTGTTCAAGCCGTTCTTCTGACCTCTTCCTAAAATGTTTATGTTCACTACA	ı
GCAGCTTCTCCGGTTCTGAGAAGAACACCTGAGAGCGCTTTCCAGCCAG	- 1800
CGTCGAAGAGGCCAAGACTCTTCTTGTGGACTCTCGCGAAAGGTCGGTC	
CCTCACACAGTACAGTTTAATAAATGTGCATTTTGATTGTAAAAAAAA	
GGAGTGTCATGTCAAATTATTTACACGTAAAACTAACATTTTTTTT	

	15
GGCGTCCCTCTGCCTGCCCACTCAGTGGCAACACCCGGGAGCTGTTTTGTCCTTTGTGGAGCCTCAGCAGTTCCCTCTTCAGA	
CCGCAGGGAGACGGACGGTGAGTCACCGTTGTGGGCCCTCGACAAAACAGGAAACACCTCGGAGTCGTCAAGGGAGAAAGTCT	TGAGTG
•	
TOCCAACACCCCTCAACACCACCCACCATCCACTACTACTAC	
TGCCAAGAGCCCTGAACAGGAGCCACCATGCAGTGCTTCAGCTTCATTAAGACCATGATGATCCTCTTCAATTTGCTCATCTTT	190
ACGGTTCTCGGGGACTTGTCCTCGGTGGTACGTCACGAAGTCGAAGTAATTCTGGTACTACTAGGAGAAGTTAAACGAGTAGAAA	GACACA
MOCESEIKIMMITENTIE	
MULFSFIKIMMILFNLLIF	L C
GGTGCAGCCCTGTTGGCAGTGGGCATCTGGGTGTCAATCGATGGGGCATCCTTTCTGAAGATCTTCGGGCCACTGTCGTCCAGT	SCCATG
CCACGTCGGGACAACCGTCACCCGTAGACCCACAGTTAGCTACCCCGTAGGAAAGACTTCTAGAAGCCCGGTGACAGGTCA	+ 270
	,dd / AC
G A A L L A V G I W V S I D G A S F L K I F G P L S S S	A M
CAGTTTGTCAACGTGGGCTACTTCCTCATCGCAGCCGGCGTTGTGGTCTTTGCTCTTGGTTTCCTGGGCTGCT	
	360
GTCAAACAGTTGCACCCGATGAAGGAGTAGCGTCGGCCGCAACACCAGAAACGAGAACCAAAGGACCCGACGA	GACTC
Q F V N V G Y F L I A A G V V V F A L G F L G C Y G A K	T F
	, <u>-</u>
AGCAAGTGTGCCCTCGTGACGTTCTTCTTCATCCTCCTCCTCATCTTCATTGCTGAGGTTGCAGCTGCTGTGGTCGCCTTGGTG	
TCGTTCACACGGGAGCACTGCAAGAAGAAGTAGGAGGAGGAGTAGAAGTAACGACTCCAACGTCGACGACACCAGCGGAACCAC	TGTGG 450
	YT
ACAATGGCTGAGCACTTCCTGACGTTGCTGGTAGTGCCTGCC	GGAAC
TGTTACCGACTCGTGAAGGACTGCAACGACCATCACGGACGG	+ 540
The state of the s	CCITG
T.M. A E H F L T L L V V P A ! K K D Y G S Q E D F T Q V	W N
ACCACCATCAAACCCCTCAACTCCTCTCACCAACTATACCAATTTTCACCAACTATACACAACTACACAACTACACAACTATACACAACTACACAACTACACAACTACACACACTACACACACTACACACACACTAC	
ACCACCATGAAAGGGCTCAAGTGCTGTGGCTTCACCAACTATACGGATTTTGAGGACTCACCCTACTTCAAAGAGAACAGTGCC	111CCC 630
TGGTGGTACTTTCCCGAGTTCACGACACCGAAGTGGTTGATATGCCTAAAACTCCTGAGTGGGATGAAGTTTCTCTTGTCACGG.	AAGGG
TTMKGLKCCGETNYTDEEDSPYEKENSA	F P
CCATTCTGTTGCAATGACAACGTCACCAACACAGCCAATGAAACCTGCACCAAGCAAAAAGGCTCACGACCAAAAAGTAGAGGGT	GCTTC
GGTAAGACAACGITACTGTTGCAGTGGTTGTGTCGGTTACTTTGGACGTGGTTCGTTTTCCGAGTGCTGGTTTTTCATCTCCCA	+ 720 CGAAG
P F C C N D N V T N T A N E T C T K Q K A H D Q K V E G	C F
AATCAGCTTTTGTATGACATCCGAACTAATGCAGTCACCGTGGGTGG	TCAAT
	
TTAGTCGAAAACATACTGTAGGCTTGATTACGTCAGTGGCACCCACC	ACTTA
NOLLYDIRTNAVTVGGVAAGIGGLELAA	M N
FGTGTCCATGTATCTGTACTGCAATCTACAATAAGTCCACTTCTGCCTCTGCCACTACTGCTGCCACATGGGAACTGTGAAGAG	
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	CACCC
ACACAGG TACATAGACATGACGTTAGATGTTATTCAGGTGAAGACGGAGACGGTGATGACGACGGTGTACCCTTGACACTTCTCC	+ 900

HPWAE25

TGGCAAGCAGCAGTGATTGGGGGAGGGGACAGGATCTAACAATGTCACTTGGGCCAGAATGGACCTGCCCTTTCTGCTCCAGACTTGGGGACCGGTCTCGTCGTCGTCGTCAGACTCGAACCCCCTGCCCTGTCCTAGATTGTTACAGTGAACCCGGTCTTACCTGGACGGGAAAGACGAGGTCTGAACCCC DAAYIGGGDRI. (SCOIDNO: 20) 

GCCAGTTCTGTTGCCCATTCCCCCAGTCTATTAAACCCTTGATATGCCCCCTAGGCCTAGTGGTGATCCCAGTGCTCTACTGGGGGATGA CGGTCAAGACAACGGGTAAGGGGGTCAGATAATTTGGGAACTATACGGGGGATCCGGATCACCACTAGGGTCACGAGATGACCCCCTACT

GAGAAAGGCATTTTATAGCCTGGGCATAAGTGAAATCAGCAGAGCCTCTGGGTGGATGTGTAGAAGGCACTTCAAAATGCATAAACCTGT CTCTTTCCGTAAAATATCGGACCCGTATTCACTTTAGTCGTCTCGGAGACCCACCTACACATCTTCCGTGAAGTTTTACGTATTTGGACA

Päge 1

HTPE	F86	
AAAAAAACAAGGTCCCCACAGCAAAGAAAAGGAATAGGATCAAGAGATACGTGGCTGCTGGCAGAGCAAGCA	GATGACTTCA 90	
TTTTTTTTGTTCCAGGGGTGTCGTTTCTTTTCCTTATCCTAGTTCTCTATGCACCGACGACCGTCTCGTTCGT		•
M N S	S M T S .	
CAGTTCCGGTGGCCAATTCTGTGTTGGTGGTGGCACCCCACAATGGTTATCCTGTGACCCCAGGAATTATGTCTCACGT	TGCCCCTGTAT	30
COTCAAGGCCACCGGTTAAGACACCACCACCGTGGGGTGTTACCAATAGGACACTGGGGTCCTTAATACAGAGTGCA		,0
A V P V A N S V L V V A P H N G Y P V T P G I M S H V	V P L Y	
CCAAACAGCCAGCCGCAAGTCCACCTAGTTCCTGGGAACCCACCTAGTTTGGTGTCGAATGTGAATGGGCAGCCTGTGCA	AGAAAGCTCTG	
GGTTTGTCGGTCGGCGTTCAGGTGGATCAAGGACCCTTGGGTGGATCAAACCACAGCTTACACTTACCCGTCGGACACG	<del></del>	70
PNSOPOVHLVPGNPPSLVSNVNGQPV(	Q K A L	
	******	
AAAGAAGGCAAAACCTTGGGGGCCATCCAGATCATCATTGGCCTGGCTCACATCGGCCTCGGCTCCATCATGGCGACGG	<del></del>	60
TTTCTTCCGTTTTGGAACCCCCGGTAGGTCTAGTAGTAACCGGACCGAGTGTAGCCGGAGCCGAGGTAGTACCGCTGCCA	AAGAGCATCCC	
KEGKTLGAIQIIIGLAHIGLGSIMAT	V L V G	
GAATACCTGTCTATTTCATTCTACGGAGGCTTTCCCTTCTGGGGAGGCTTGTGGTTTATCATTTCAGGATCTCTCTC	TGGCAGCAGAA	50
CTTATGGACAGATAAAGTAAGATGCCTCCGAAAGGGAAGACCCCTCCGAACACCAAATAGTAAAGTCCTAGAGAGAG	ACCGTCGTCTT	
EYLSISFYGGFPFWGGLWFI!SGSLS	V A A E	
AATCAGCCATATTCTTATTGCCTGCTGTCTGGCAGTTTGGGCTTGAACATCGTCAGTGCAATCTGCTCTGCAGTTGGAGT	TCATACTCTTC	
TTAGTCGGTATAAGAATAACGGACGACAGACCGTCAAACCCGAACTTGTAGCAGTCACGTTAGACGAGACGTCAACCTC.	<del></del>	40
	v I I F	
N Q P Y S Y C L L S G S L G L N I V S A I C S A V G	, , , , ,	
ATCACAGATCTAAGTATTCCCCACCCATATGCCTACCCCGACTATTATCCTTACGCCTGGGGTGTGAACCCTGGAATGG	<del></del>	30
TAGTGTCTAGATTCATAAGGGGTGGGTATACGGATGGGGCTGATAATAGGAATGCGGACCCCACACTTGGGACCTTACC	GCTAAAGACCG	
ITDLSIPHPYAYPDYYPYAWGVNPGM	A I S G	
GTGCTGCTGGTCTTCTGCCTCCTGGAGTTTGGCATCGCATCGCATCTTCCCACTTTGGCTGCCAGTTGGTCTGCTGTC	AATCAAGCAAT	20
CACGACGACCAGAAGACGGAGGACCTCAAACCGTAGCGTAGCGTAGAAGGGTGAAACCGACGGTCAACCAGACGACGACAG	TTAGTTCGTTA	20
V L L V F C L L E F G I A C A S S H F G C Q L V C C	a s s N	
GTGAGTGTCATCTATCCAAACATCTATGCAGCAAACCCAGTGATCACCCCAGAACCGGTGACCTCACCACCAAGTTATT	CCAGTGAGATC	
CACTCACAGTAGATAGGTTTGTAGATACGTCGTTTGGGTCACTAGTGGGGTCTTGGCCACTGGAGTGGTGGTTCAATAA	+	10
V S V I Y P N I Y A A N P V I T P E P V T S P P S Y	S S E I	

Päge 1

																								H?	5P	R	FC	2د		
CACG	AG	CAG	GGT	CTC	GGG	CTA	GTC	ATG	GCG	TCC	CCG.	TCT	CGG/	AGAG	CTG	CAG	ACT	AAA	CCAI	GTC	ATT	ACT	GT	TTC	AAGA	AGC	STTO	CTG	CTA	00
GTGC	TC	GTC	CCA	GAG	CCC	GAT	CAG	TAC	CGC	AGG	GGC,	AGA	3CC	TCT	GAC	GTC	TGA	TTT	GGT	CAG	TAA	TGA	CAA	AAG	rtci	rcg	CAAC	GAC		90
				ķ	_			M	A	S	Р	S	Ŕ	R	L	Q	T	κ	Р	٧	I	T	С	F	K	s	٧	L	ι.	
ATCT																		<del></del>								+				180
TAGA	TG	TGA	AAA	TAA	AAG	ACC	TAG	TGA	CCĠ	CAA	TAG	GAA	GAA	CGT	CAA	CCG	TAA.	ACC	CCG	TTC	CAC	TCG	GAC	CTC	TTA	ATG.	444	AGA	GAA	
I	Υ	T	F	i	F	W	I	T	G	٧	i	L	L	A	V	G	i	w	G	ĸ	٧	S	L	Ε	N	Y	F	S	L	
TTAA	ΑT	GAG	AAG	GCC.	ACC.	ΑĄΤ	GTC	ccc	TTC	GTG	стс	ATT	GC T	ACT	GGT	ACC	GTC.	ATT	ATT	стт	TTG	GGC.	ACE.	TTT	GGT.	TGT	777	GC T	ACC	270
AATT	ΤA	стс	TTC	CGG	TGG	TTA	CAG	GGG.	AAG	CAC	GAG	TAA	CGA	TGA	CCA	TGG	CAG	TAA	TAA	GAA.	AAC	CCG	TGG	AAA	CCA	ACA.	AAA	CGA	TGG	
L	N	Ε	Κ	Α	T	N	٧	Ρ	F	٧	Ł	I	A	T	G	T	٧	ı	I	L	L	G	T	F	G	С	F	A	T	
TGCC	GA	GCT	TCT	GCA	TGG	ATG	CTA.	AAA	CTG	TAT	GCA	ATG	TTT	CTG	ACT	CTC	GTT	TTT	TTG	GTC	GAA	CTG	GTC	GCT	GCC	ATC	GTA	GGA	TTT	360
ACG	CT	CGA	AGA	CGT	ACC	TAC	GAT	TTT	GAC	ATA	CGT	TAC	AAA	GAC	TGA	GAG	CAA	AAA	AAC	CAG	CTT	GAC	CAG	CGA	CGG	TAG	CAT	CCT	AAA	
С	R	Α	S	Α	W	М	L	K	L	Y	Α	M	F	L	T	L	٧	F	L	٧	Ε	L	٧	Α	Α	í	٧	G	F	
GTTI	TC	AGA	CAT	GAG	ATT	AAG	AAC	AGC	<b>T</b> T T	AAG	AAT	AAT	TAT	GAG	AAG	GCT	TTG	AAG	CAG	TAT	AAC	TCT	ACA	GGA	GAT	TAT	AGA.	AGC	CAT	450
CAAA	AAG	TCT	GTA	стс	TAA	тіс	TTG	TCG	AAA	TTC	TTA	TTA	ATA	CTC	TTC	CGA	AAC	TTC	GTC	ATA	TTG	AGA	TGT	CCT	CTA.	ΑТА	TCT	TCG	GTA	
٧	F	R	Н	Ε	I	Κ	N	S	F	κ	N	N	Y	Ε	ĸ	Α	L	ĸ	a	Y	N	s	T	G	O	Y	R	s	н	
GCAC	STA	GAC	AAG	ATC	CAA	AAT	ACG	TTG	CAT	TGT	TGT	GGT	GTC	ACC	GAT	TAT	AGA	GAT	TGG	ACA	GAT	ACT	AAT	TAT	TAC	TÇA	GAA	AAA	GGA	E#0
CGT	AT	CTG	TTC	TAG	GTT	TTA	TGC	AAC	GTA	ACA	ACA	CCA	CAG	TGG	CTA	ATA	TCT	CTA	ACC	TGT	CTA	TGA	TTA	ATA	ATG	AGT	CTT	TTT	cct	340
A	٧	ם	κ	I	a	N	T	L	н	С	С	G	٧	T	D	Y	R	D	W	T	D	T	N	Υ	Y	S	Ε	K	G	
TTT	ст	AAG	AGT	TGC	TGT	AAA	CTT	GAA	GAT	TGT	ACT	CCA	CAG	AGA	GAT	GCA	GAC	AAA	GTA	AAC	AAT	GAA	GGT	TGT	TTT	AŢA	AAG	GTG	ATĢ	630
AAA	GGA	TTC	TCA	ACG	ACA	TTT	GAA	CTT	CTA	ACA	TGA	GGT	GTC	TCT	СТА	CGT	CTG	TTT	CAT	TTG	TTA	CTT	CCA	ACA	AAA	TAT	TTC	CAC	TAC	030
F	Р	ĸ	s	С	С	κ	L	Ε	٥	c	τ	ρ	۵	R	D	Α	D	Κ	٧	N	N	ε	G	С	F	I	κ	٧	M	
ACC	ATT	ATA	GAG	TCA	GAA	ATG	GGA	GTC	GTT	GCA	GGA	ATT	TCC	TIT	GGA	GTT	GCT	TGC	TTC	CAA	CTG	ATT	GGA	ATC	TTT	CTC	GCC	TAC	TGC	720
TGG	TAA	TAT	стс	AGT	CTT	TAC	CCT	CAG	CAA	CGT	CCT	TAA	ÁGG	AAA	CCT	CAA	CGA	ACG	AAG	GTT	GAC	TAA	ĊCT	TAG	AAA	GAG	CGG	ATG	ACG	
T	į	ī	Ε	s	Ε	M	G	٧	٧	A	G	i	s	F	G	٧	Α	С	F	Q	L	ı	G	I	F	L	A	Y	С	
стс	TCT	CGT	rgcc	ΑΤΑ	ACA	AAT	AAC	CAG	TAT	GAG	ATA	GTG	TAA	ccc	AAT	GŢA	тст	GTG	GGC	CTA	TTC	стс	TCT	ACC	TTT	AAG	GAC	ATT	TAG	810
GAG	AGA	GCA	VC G G	TAT	TGT	TTA	TTG	GTC	ATA:	CTC	TAT	CAC	ATT	GGG	TTA	CAT	AGA	CAC	CCG	GAT	AAG	GAG	AGA	TGG	AAA	TTC	CTG	TAA	ATC	0,0
L	s	R	A	1	т	N	N	0	Υ	ε	i	٧	. (	LS	eG	, J	D	N	D: (	2.	2)									
CCT	רבו				. T T A	CAA	ACT	TCC	TTO	ימרו	cc.	CA4	ירדמ		ΔCΔ	.r T Δ	CTI	ר א ר ז	:CAT	AGA	CCA	AAA	AAC	TAC	ACC	AGT	AGG	TTG	ATT	900
CCA	GGG	GGG	SACA	LC T 1	TAAT	CTI	TTCA	ACG	SAAC	CGA	CCI	CTI	GAC	TGT	TGT	GAT	GAA	TGA	CTA	TCT	GGT	TTT	TTG	ATG	TGG	TCA	TCC	AAC	TAA	

HLTAH8O	
CACGAGCATTGCCGCTCTCTCGGTGAGCGCAGCCCCGCTCTCCGGGCCGGGCCTTCGCGGGCCACCGGCGCCATGGGCCAGTGCGGCAT	 C + 90
STGCTCGTAACGGCGAGAGAGCCACTCGCGTCGGGGCGAGAGGCCCGGCAGCGCCCGGTAGCCCGGTACCCGGTCACGCCGTA	
м в о с в 1	•
ACCTCCTCCAAGACCGTGCTGGTCTTTCTCAACCTCATCTTCTGGGGGGGCAGCTGGCATTTTATGCTATGTGGGAGCCTATGTCTTCAT	C + 180
TGGAGGAGGTTCTGGCACGACCAGAAAGAGTTGGAGTAGAAGACCCCCCGTCGACCGTAAAATACGATACACCCTCGGATACAGAAGTA	
T S S K T V L V F L N L I F W G A A G I L C Y V G A Y V F I	
ACTTATGATGACTATGACCACTTCTTTGAAGATGTGTACACGCTCATCCCTGCTGTAGTGATCATAGCTGTAGGAGCCCTGCTTTTCAT	۲
TGAATACTACTGATACTGGTGAAGAAACTTCTACACATGTGCGAGTAGGGGACGACATCACTAGTATCGACATCCTCGGGACGAAAAGTA	+ 2/0
	u
TYDDYDHFFEDVYTLIPAVVIIAVGALLFI	
ATTGGGCTAATTGGCTGCTGTGCCACAATCCGGGAAAGTCGCTGTGGACTTGCCACGTTTGTCATCATCCTGCTCTTGGTTTTTGTCAC	A + 360
TAACCCGATTAACCGACGACACGGTGTTAGGCCCTTTCAGCGACACCTGAACGGTGCAAACAGTAGTAGGACGAGAACCAAAAAACAGTG	T
I G L I G C C A T I R E S R C G L A T F V I I L L V F V T	
GAAGTTGTTGTAGTGGTTTTGGGATATGTTTACAGAGCAAAGGTGGAAAATGAGGTTGATCGCAGCATTCAGAAAGTGTATAAGACCTA	Ç , ,, 50
CTTCAACAACATCACCAAAACCCTATACAAATGTCTCGTTTCCACCTTTTACTCCAACTAGCGTCGTAAGTCTTTCACATATTCTGGAT	+ 450 G
EVVVVLGYVYRAKVENEVDRSI QKVYKTY	
	c
AATGGAACCAACCCTGATGCTGCTAGCCGGGCTATTGATTATGTACAGAGACAGCTGCATTGTTGTGGAATTCACAACTACTCAGACTG TTACCTTGGTTGGGACTACGACGATCGGCCCGATAACTAATACATGTCTCTGTCGACGTAACAACACCTTAAGTGTTGATGATGACTAC	+ 540
TIALCI IGGI IGGGACIACGALGAI CGGCCCGAIAACI AAIACAIGICIC IGICGACGI AACAACACCI IAAGIGI IGAIGAAGA	
NGTNPDAASRAIDYVOROLHCCGIHNYSDV	,
GAAAATACAGATTGGTTCAAAGAAACCAAAAACCAGAGTGTCCCTCTTAGCTGCTGCAGAGAGACTGCCAGCAATTGTAATGGCAGCCI	G + 630
CTTTTATGTCTAACCAAGTTTCTTTGGTTTTTGGTCTCACAGGGAGAATCGACGACGTCTCTCTGACGGTCGTTAACATTACCGTCGGA	Ċ
ENT D'WFKETKNQS VPLS CCRETAS NCNGS L	
GCCCACCCTTCCGACCTCTATGCTGAGGGGTGTGAGGCTCTAGTTGTGAAGAAGCTACAAGAAATCATGATGCATGTGATCTGGGCCGC	A + 720
CGGGTGGGAAGGCTGGAGATACGACTCCCACACTCCGAGATCAACACTTCTTCGATGTTCTTTAGTACTACGTACACTAGACCCGGCC	
A H P S D L Y A E G C E A L V V K K L Q E I M M H V I W A A	ı
CTGGCATTTGCAGCTATTCAGCTGCTGGGCATGCTGTGTGCTTGCATCGTGTTGTGCAGAAGGAGTAGAGATCCTGCTTACGAGCTCT	.c
GACCGTAAACGTCGATAAGTCGACGACCCGTACGACACACGACGTAGCACAACACGTCTTCCTCATCTCTAGGACGAATGCTCGAGAA	7 010
LAFAAIOLLGMLCACIVLCRRSRDPAYEL [§]	

							_																ŀ		PΈ	A	$\mathcal{I}\mathcal{I}$			
CA	GA	CG	CAG	AGC	TTG	GGG	CTT	CCT	TGG	TCG	CAC	CCA	CCA	CCT	CCC	TGC	CCA	CTG	GTC.	AGC	CTT	CAG	3GAI	ccc.	GAG	CAC	CGC	СТС	GTC	90
GT	SCT	CGC	GTC	TCG	AACI	ccc	GAA	GGA	ACC.	AGC	GTG	GGT	GGT	GGA	CGG.	ACG	GGT	GAC	CAG	TCG	GAA	STC	CCT	GGG	ACTO	ccto	GCO	GAC	CAG	90
					AGČ(								<del></del>										<del></del>							180
AG	444	GGA	CAC	CGG	TCG	GGT	CTT	GAC	TTC	GCG.	ACG	CCG	TAC	CGC	GCG	CGG	ACG	GAG	GTC:	CGG	CAG	TTC	ATG:	GAG	rac,	AAG(	GG	AG1	TGG	
													п	А	ĸ	А	L	L	u	д	٧	^	,	_	m	г	А	r	N	
TG	ттс	TTC	TĢG	CTG	GGA	GGC	TGT	GGC	GTG	ÇTG	GGT	GTC	GGC	ATC	TGG	CTG	GCC	GCC	ACA	ÇAG	GGG	AGC	TTC	GCC	ACG	CTG.	TCT:	rc T	rcct	270
AC.	AAG,	AAG.	ACC	GAC	CCT	ccg	ACA	CCG	CAC	GAC	CCA	CAG	ccg	TAG	ACC	GAC	CGG	CGG	TGT	GTC	ccc.	TCG	AAG	CGG	TGC	GAC,	4GA/	4GA/	AGGA	270
L	F	F	W	Ĺ	G	G	С	G	٧	L	G	٧	G	ı	W	L	A	Α	T	Q	G	S	F	A	τ	L	S	S	S	
TC	ccg	тсс	стс	TCG	GET	GCC	AAC	CTG	стс	ATC.	ATC	ACC	GGC	GCC	TTT	GTÇ	ATG	GCC.	ATC	GGC.	TTC	GTG	GGC	TGC	CTG	GGŢ	GCC.	ÁTC	AAGĢ	200
AG	GGC	AGG	GAC	AGC	CGA	CGG	TTG	GAC	GAG	TAG	TAG	TGG	CCG	CGG	AAA	CAG	TAC	cgg	TAG	CCG.	AAG	CAC	CCG	ACG	GAC	CCA	CGG	TAG	TTCC	360
F	ρ	s	L	S	Α	A	N	L	L	I	I	T	G	А	F	٧	м	A	I	G	F	٧	G	С	L	G	А	I	К	
AG.	AAC.	AAG	TGC	СТС	CTG	стс	ACT	TTC	TTC	CTG	CTG	CTG	CTG	CTG	GTG	TTC	CTG	CTG	GAG	GCC.	ACC.	ATC	GCC	ATC	стс	TTC	TTC	BCC.	TACA	
			-+-				<del></del>			+			-+-		+	+				<del></del>			-		+				ATGT	450
Ε	N	K	С	L	L	L	т	F	F	Ļ	L	L	L	L	٧	F	L	L	ε	A	T	I	A	ŀ	L	F	F	Α	Υ	
CG	GAC.	AAG	ATT	GAC	AGG	TAT	GCC	CAG	CAA	GAC	CTG	AAG	ΑΑΑ	GGC	TTG	CAC	CTG	TAC	GGC	ACG	CAG	GGC.	AAC	GTG	GGC	стс.	ACC.	AAÇ	GCCT	
			-+-		<del></del>			-+-		+			-+-		+	+		-+		<del></del>			- 1		+	<del>i</del>			CGGA	540
т	0	v	1	Ω	D	¥	٨	Λ.	0	n		v	r	c	1	ч	,	Y	G	T	n	ı:	N	v	G	ı	т	N	A	
•	U	•	•	Ü	IX.	,	^	· ·	•	Ü	_		``	ŭ	_	••	_	•		•	_		,,	-		<u>-</u>	· 		<b></b>	
			-+-		+	+				+			-+-			$\rightarrow$				<del></del>	+		-+-		+	<del></del> -		-+-	GACT	630
CC	TCG	TAG	TAG	GTC	TGG	CTG	AAG	GCG	ACG	ACA	CCG	CAG	AGG	TTG	ATG	TGA	CTG	ACC	AAG	CTC	CAC	ATG	TTG	CGG	TGC	GCC	CAT	GGA	CTGA	
W	s	I	Ī	Q	T	D	F	R	С	С	G	٧	s	N	Y	T	D	W	F	ε	٧	Y	N	A	T	R	٧	Р	D	
CC	TGC	TGC	TTG	GAG	TTC.	AGT	GAG	AGC	TGT	GGG	CTG	CAC	GCC	ССС	GGC	ACC	TGG	TGG	AAG	GCG	ccg	TGC	TAC	GAG	ACG	GTG	AAG	GTG	TGGÇ	720
GG	ACG.	ACG	AAC	стс	AAG	TCA	CTC	TCG	ACA	ccc	GAC	GTG	CGG	GGG	CCG	TGG	ACC	ACC	TTC	CGC	GGC	ACG	ATG	стс	TGC	CAC	TTC	CAC	ACCG	720
s	С	С	L	ε	F	s	Ε	s	С	G	L	н	A	Р	G	T	W	w	κ	A	Ρ	С	Y	Ε	T	٧	K	٧	W	
ŦT	CAG	GAG	AAC	CTG	CTG	GCT	GTG	GGC	ATC	TTT	GGG	CTG	TGC	ACG	GCG	CTG	GTG	CAG	ATC	CTG	GGC	CTG	ACC	TTC	GCC	ATG	ACC	ATG	TACT	
AA	GTC	CTC	TTG	GAC	GAC	CGA	CAC	CCG	TAG	AAA	ccc	GAC	ACG	TGC	CGC	GAC	CAC	GTC	TAG	GAC	CCG	GAC	TGG	AAG	CGG	TAC	TGG	TAC	ATGA	810
		Ε							ı									•		L								М		
GC	CAA	GTG	GTC	AAG	GCA	GAC	ACC	TAC	TGC	GCG	TAG	GCC	GCC	CAC	כפר	CGG	CTT	CTC	TGC	CAA	AAG	GAC	GCC	CAC	GGG	GAG	ATG	GCC	GCAC	
																										<del></del>			CGTG	900
С	۵	٧	٧	K	A	D	т	Υ	С	A		<b>(</b> S	eG	J	D.	110	ე : .	24	Ð											

GCGCCGCCGGGCCGCAGCATGGGGCGCTTCCGCGGGGGCCTGCGGTGCATCAAGTACCTGCTGCTTGGCTTCAACCTGCTCTTCTG	GCTG 90
CGCGGCGGCCCGGCGTCGTACCCCGCGAAGGCGCCCCCGGACGCCACGTAGTTCATGGACGAACCGAAGTTGGACGAGAAGAC	
M G R F R G G L R C I K Y L L L G F N L L F V	, L.
	. <b>-</b>
GCTGGATCGGCCGTCATTGCTTTTGGACTATGGTTTCGGTTCGGAGGTGCCATAAAGGAGTTATCATCAGAGGACAAGTCCCCAGA	+ 180
CGACCTAGCCGGCAGTAACGAAAACCTGATACCAAAGCCAAGCCTCCACGGTATTTCCTCAATAGTAGTCTCCTGTTCAGGGGTCT	CATA
A G S A V I A F G L W F R F G G A I K E L S S E D K S P E	. Y
TTCTATCTCCCCCTCTATATCTTCTCCTTCAACCCCCCCC	CTCC
TTCTATGTGGGGCTGTATGTTCTGGTTGGAGCCGGGGCCCTGATGATGGCCGTGGGGTTCTTCGGATGCTGCGGAGCCATGCGGGA	<del>+</del> 270
AAGATACACCCCGACATACAAGACCAACCTCGGCCCCGGGACTACTACCGGCACCCCAAGAAGCCTACGACGCCTCGGTACGCCCT	CAGL
FYVGLYVL V G A G A L M M A V G F F G C C G A M R E	: s
CAATGTGTGCTTGGATCATTTTTTACCTGCCTCCTGGTGATATTTGCTGCTGAAGTAACCACTGGAGTATTTGCTTTTATAGGCAA	GGGG
GTTACACACGAACCTAGTAAAAAATGGACGGAGGACCACTATAAACGACGACTTCATTGGTGACCTCATAAACGAAAATATCCGTT	<del>+</del> 360
Q C V L G S F F T C L L V I F A A E V T T G V F A F I G K	, G
GTAGCTATCCGACATGTTCAGACCATGTATGAAGAGGCTTACAATGATTACCTTAAAGACAGGGGAAAAGGCAATGGGACACTCAT	CACC
CATCGATAGGCTGTACAAGTCTGGTACATACTTCTCCGAATGTTACTAATGGAATTTCTGTCCCCTTTTCCGTTACCCTGTGAGTA	GTGG
V A I R H V Q T M Y E E A Y N D Y L K D R G K G N G T L I	Т
TTCCACTCAACATTTCACTCTCTCTCCAAAAAAAAAAA	TICC
TTCCACTCAACATTTCAGTGCTGTGGAAAAGAAAGCTCCGAACAGGTCCAACCTACATGCCCAAAGGAGCTTCTAGGACACAAAGAA	<del>+</del> 540
AAGGTGAGTTGTAAAGTCACGACACCTTTTCTTTCGAGGCTTGTCCAGGTTGGATGTACGGGTTTCCTCGAAGATCCTGTGTTCTT	AACG
F H S T F Q C C G K E S S E Q V Q P T C P K E L L G H K M	i C
ATCGATGAAATTGAGACCATAATCAGTGTTAAGCTCCAGCTCATTGGAATTGTCGGTATTGGAATTGCAGGTCTGACGATCTTTGG	CATG
TAGCTACTTTAACTCTGGTATTAGTCACAATTCGAGGTCGAGTAACCTTAACAGCCATAACCTTAACGTCCAGACTGCTAGAAACC	GTAC
	. м
	1 <b>11</b>
ATATTCAGCATGGTCCTCTGCTGTGCGATACGAAACTCACGAGATGTGATATGAAGCTACTTCTACATGAAAATTGCAATCTAAAG	720
TATAAGTEGTACEAGGAGAEGAEACGETATGETTTGAGTGETETACAETATAETTEGATGAAGATGTAETTTTAAEGTTAGATTTE	GAAA
IFSMVLCCAIRNSROVI. (SCOIDNO:25)	
$\supset$	
GTATGGTTTACAAG 734 (SCO ID NO'.9)	
GTATGGTTTACAAG LOCK LOCKS, J	

Päge 1

HAI DQ59	
AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC	90
TCACAAATACCCTGATTTTTTGAAAATTGTGGAAAAATCCCCTTTATAAAACCAGGATATGTTTTGTACATTTATACGAAATAATGAAAG	
ATTTTCTGACCCTGCTGTAAACTACTGCAACCCTCACATCCCTCAAAGGGACTTTTATGTCAAACTCTTCTGTTTCTCCAAATATAAGGA	180
TAAAAGACTGGGACGACATTTGATGACGTTGGGAGTGTAGGGAGTTTCCCTGAAAATACAGTTTGAGAAAGACAAAGAGGTTTATATTCCT	
AAAAAGACTAAAGCAAGAGATCTGGCAGTTGAAAATTGTGGGGAAAGAGAATTTGTATGGGCACTGTATCTATGAAATACCTCATACTTAC	270
TTTTCTGATTCGTTCTCTAGACCGTCAACTTTTAACACCCTTTCTCTTAAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	
GTTTACATGTTTTCCTAACTTTTTGTATTTTCTTGTATAGCCACCTAGAGAATTCTTCATAGATTAAGAACTACAGTTTTCACCACTTA	360
CAAATGTACAAAAGGATTGAAAAAACATAAAAAGAACATATCGGTGGATCTCTTAAGAAGTATCTAATTCTTGATGTCAAAAGTGGTGAAT	
ACATAAGTAAAACAAAGTCCTTCATAATTTAACCATTAGCATCTTTGGCCAAACCAAAATAAAGAAAAGCATCTTCTCCTAGTTGTGTG	- 450
TGTATTCATTTTGTTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGGATCAACACACA	i
GGGCAACAGAACAAGTTAAGGAAACAAAAATACTTATATACACAGAACAAAAATAATGTTCTTTTTATGCAAATCCCCTGTGAAAAT	540
CCCGTTGTCTTTGTTCAATTCCTTTGTTTTTATGAATATATGTGTCTTGTTTTTATTACAAGAAAAATACGTTTAGGGGACACTTTTA	•
AAAAATTTTCAATGTTTAAAAAAAAAAAAAAAAAAAAAA	
TTTTAAAAGTTACAAATTTTTTTTTTTTTTTTTTTTTTT	

																									口	止	<u> </u>	<u>54</u>	$\bigcirc$	
TTC	GGC	ACG	AGC	TGC	GGG	CGG	TGG	GCG	GCT	GGG	CGG	CCC	ÇGG	GAG	CCG	CGC	TCT	CAG	TCT	CTC	TAG	GCG	ÇAG	TCC	CTT	cgc	CGC	TTC	CGG	00
AAG	CCG	TGC	TCG	ACG	ccc	GCC	ACC	CGC	CGA	ccc	GCC	GGG	GCC	стс	GGC	GCG	AGA	GTC	AGA	GAG	ATC	CGC	GTC	AGG	GAA	GCG	GCG.	AAG	GCC	90
				,																										
AGC	ccc	TGG	CAG	GGC	` ECA	GAA	GCC	ATG	GCC	CAC	TAT.	AAG	ACT	GAG	CAG	GAC	GAC	TGG	CTG	ATC	ATC	TAC	TTG	AAG	TAT	TTA	стс	TTT	GTC	
TCG	ccc	۸۲۲	GTC		GGT	CIT	cee	TAC		GTG	ΔΤΔ	TTC	TGA	r T C	GTC	r T G	CIG	4 C C	——÷	TAG	TAG	ΔΙΩ	<del> </del>	TTC	ΔΤΔ	ΔΑΤ	GAG	ΔΔΔ		180
1 C G	300	ALC	6,6	CCG	331	C 1 1	CGG	IAC	cuu	<b>G</b> 1 G	A 1 A		IGA	<b>C</b> 1 C	010	CIG	,	700	مرد			~, 0	AAC	, , ,	^'^	AA 1	GAG.	~~~	CAG	
								М	Α	Н	Y	K	T	Ε	Q	D	D	W	L	I	I	Y	L	K	Y	L	L	F	٧	
TTC	AAC	TTC	TIC	TTC	TGG	GTC	GGG	GGA	GCA	ecc.	GTC	CTG	GCT	GTG	GGC	ATC	TGG	ACC	CTG	GTG	GAG	AAG	AGT	GGC	TAC	стс	AGC	GTC	CTG	
			<del></del>	+		<del></del>		+			<del></del>		+	+				<del></del>	+	CAC			<del> </del>			<del></del>			$\rightarrow$	270
AAG	116	ААС	ААС	ААС	ACC	CAG		<b>CC</b> 1	CGI	666	CAG	GAC	CGA	CAC		1 40	ACC	100	GAC				, , ,	cca	~ ' ' '	GAG	166	CAG	GAL	
F	Ν	F	F	F	W	٧	G	G	A	Α	٧	L	Α	٧	G	I	W	T	L	٧	ε	K	S	G	Υ	L	S	٧	L	
ec c	TCC	AGC	ACE	TTT	GCC	GCC	TCC	GCC	TAC	ATC	CTC	ATC	TTT	GCG	GGC	GTA	CTT	GTC	ATG	GTG	ACC	GGC	ттс	CTG	GGC	TTC	GGT	GCC	ATC	
	ACC	TCC	TCC	^ ^ ^	CCC	CGG	AGG.	cee	4 T.G	TAG	GAG	TAG	1	CGC	ccc	_ <del> </del>	CAA	CAG	TAC	CAC	TGG	cce	A A G	GAC	LLE	ΔAG	CCA	CGG	TAG	360
LGG	AGG	100	100	~~~	CGG	CGG	AGG	CGG	~ 1 0	1 7 0	470			CGC	cca	CAI	UAA	CAG	170				~~0	ممد			CCA	cuo		
A	S	S	T	F	A	Α	S	Α	Y	I	L	I	F	A	G	٧	Ł	٧	М	٧	Ţ	G	F	Ĺ	G	F	G	A	I	
стс	TGG	GAG	CGG	AAG	GGC	TGC	стс	TCC	ACG	TAT	TTC	TGC	CTG	TTG	стс	GTC	ATC	TTC	CTG	GTT	GAG	CTG	GTG	GCG	GGA	GTC	CTG	GCC	CAT	
GAG	ACC	CTC	ecc.	TTC	cce	ACG	GAG	AGG	TGC	ATA	AAG	ACG	GAC	AAC	GAG	CAG	TAG	AAG	GAC	CAA	CTC	GAC	CAC	CGC	CCT	CAG	GAC	CGG	GTA	450
<i>0</i> /1 <b>0</b>			000			,	٠,٠٠						•,	, ., .		•														
L	W	Ε	R	K	G	С	L	S	T	Υ	F	С	L	L	L	٧	ŧ	F	L	٧	Ε	Ĺ	٧	Α	G	٧	. L	Α	Н	
GTG	TAT	TAC	CAG	AGG	CTG	AGT	GAT	GAA	CTG	AAG	CAG	CAC	TTG	AAC	CGG	AC T	CTG	GCT	GAG	AAC	TAC	GGG	CAG	CCG	GAG	CAC	GCA	GAT	CAC	
CAC	ATA	ATG	GTC	TCC	GAC	TCA	CTA	CTT	<del>+</del> GAC	TTC	GTC	GTG	+ AAC	TTG	GCC	TGA	GAC	CGA	C T C	TTG	ATG	CCC	G T C	GGC	CTC	GTG	CGT	CTA	GTG	540
																												_		
٧	Υ	Y	Q	R	L	S	D	Ε	L	K	Q	Н	L	N	R	T	L	Α	Ε.	N	Y	G	٥	Ρ.	Ε	н	Α	D	н	
GCC	TCA	GTG	GAC	CGA	стс	CAG	CAG	GAT	TTC	AAG	TGC									TGG	CAG	CAC	AGC	ACG	TAC	ATC	CTG	TTG	CGG	
CGG	AGT	CAC	CTG	GCT	GAG	GTC	GTC	CTA	AAG	TTC	ACG	ACG	CCT	TCG	TTG	TCG	AGT	CGG	CTG	ACC	GTC	GTG	TCG	TGC	ATG	TAG	GAC	AAC	GCC	630
	_			_			_		_		_	_		_		_	_		_		_		_	_					_	
Α	S	٧	D	R	L	Q	Q	D	F	K	С	ε	G	S	N	5	S	Α	D	W	a	н	S	F	Y	ì	L	L	к	
GAG	GCC	GAG	GGC	cgc	CAG	GTG	ссс	GAC	AGC	TGC	TGC	AAG	ACA	GTG	GTG	GCG	CGC	TGC	GGC	CAG	cgg	GCC	CAC	ссс	TCC	AAC	ATC	TAT	AAĢ	700
CTC	CGG	CTC	CCG	GCG	GTC	CAC	GGG	CTG	TCG	ACG	ACG	TTC	TGT	CAC	CAC	<del>-:-</del> CGC	GCG	ACG	CCG	GTC	GCC	CGG	GTG	GGG	AGG	TTG	TAG	ATA	TTC	720
_		_		_	_		_	_	_	_	_		_				_			_	_			_	_			.,	.,	
Ε	A	Ę	G	R	Q	٧	Р	D	S	С	С	K	T	٧	٧	Α	R	С	G	Q	R	А	н	Р	S	N	i	Y	K	
GTG	GAG	GGA	GGC	TGC	стс	ACC	AAG	CTG	GAG	CAG	TTC	CTG	GCC	GAC	CAC	стс	ста	CTT	ATG	GGG	GCA	GTG	GGC	ATC	GGG	GTG	GCC	TGC	CTĢ	010
CAC	стс	CCT	CCG	ACG	GAG	TGG	TTC	GAC	CTC	GTC	AAG	GAC	CGG	СТС	GTG	<del>G</del> AC	GAC	GAA	TAC	ccc	CGT	CAC	CCG	TAG	ccc	CAC	EGG	ACG	GAC	810
						_												•					_							
٧	Ε	G	G	С	Ĺ	T	K	L	Ε	Q	F	L	Α	D	Н	L	L	Ĺ	M	G	Α	٧	G	ı	G	٧	А	L	L	
CAG	ATC	TGC	GGG	ATG	GTT	стс	ACC	TGC	TGC	TTG	CAC	CAG	AGG	сто	CAG	CGG	CAT	ттт	TAC	TAA	TGG	CAA	CCA	ССТ	CCT	стт	CCA	ACT	GCĆ	000
GTC	TAG	ACG	CCC	TAC	CAA	GAG	TGG	ACG	ACG	AAC	GTG	GTC	TCC	GAG	GTC	GCC	GTA	AAA	ATG	ATT	ACC	GTT	GGT	GGA	GGA	GAA	GGT	TGA	CGG	900
						_																						_		
Q	Ī	С	G	М	٧	L	T	С	С	L	H	Q	R	L	a	R	Н	F	Υ	. (	' <b>\</b>	P(\)	7	$\perp$	$\sim$	$\cap$	(به ۱	6	)	

GGAGTICTGTTGTACACCGTGTACGGTAGACGTTCC 936 SCO IDNO: 11

																								Н	G	P)(	<del>}</del>	89	ı	
AGC	TT	ACTI	TCA	CTO	ACC	GÇC	TGT	CCT	TCC	TGA	CAC	CTC	ACC	ATG		ACG				GCC	CGC	TGT	GTG	GGG	СТС	TCC	CTC	ATT	ACC	
TCG	AAI	rga.	AGT	GAG	STGG	CGG	ACA	GGA	AGG	ACT	GTG	GAG	TGG	TAC							GCG	SACA	CAC	ccc	GAG	SAGO	GAG	TAA		90
				į.	1									M	С	T	G	K	С	A	R	С	٧	G	L	S	L	I	Τ.	
сто	TGC	сто	GTC	TGC	ATT	GTG	GCC	AAC	GCC	стс	СТС	CTG	GTA	CCT	AAT	GGG	GAG	ACC	TCC	TGG	ACC	AAC	ACC	AAC	CAT	гстс	:AGC	TTG	CAA	
GAG	ACC	GAG	CAG	ACC	TAA	CAC	CGG	TTG	CGG	GAG	GAC	GAC	CAT	GGA	TTA	ссс	стс	TGG	AGG	ACC	TGG	TTG	TGG	TTG	GTA	1GAC	TCG	AAC	GTT	180
L	С	L	٧	С	Ī	٧	A	N	A	L	L	L	٧	Ρ	N	G	Ε	Ţ	s	W	T	N	T	N	н	L	s	L	۵	
GTC	TGO	CTC	ATG	GGC	GGC	TTC	ATT	GGC	GGG	GGC	CTA	ATG	GTA	C T G	TGT	CCA	GGG	ATT	GCA	GCC	GTI	CGG	GCA	.GGG	GGC	CAAG	GGC	TGC	TGT	
CAG	ACC	GAG	TAC	ccc	CCG	AAG	TAA	CCG	ccc	CCG	GAT	TAC	CAT	GAC	ACA	GGT	CCC	TAA	CGT	CGG	CAA	GCC	CGT	CCC	CCG	3770	:ccg	ACG	ACA	270
٧	W	Ł	М	G	G	F	I	G	G	G	L	м	٧	L	С	٩	G	I	A	Α	٧	R	A	G	G	κ	G	С	С	
GGT	GCT	rggg	TGC	TGT	GGA	AAC	CGC	TGC	AGG	ATG	CTG	CGC	TCG	GTC	TTC	TCC	TCG	GCG	TTC	GGG	GTG	CTT	GGT	GCC	ATC	TAC	TG			
CCA	CGA	ACCC	ACG	ACA	CCT	TTG	GCG	ACG	TCC	TAC	GAC	GCG	AGC	CAG	AAG	AGG	AGC	CGC	AAG	CCC	CAC	GAA	CCA	CGG	TAC	ATG	ACG	G		
G	Α	G	С	С	G	N	R	С	R	М	L	R	S	٧	F	s	s	A	F	G	٧	L	G	Α	ı	Y	С	L		
GTG	TCI	rgg.	GCT	GGG	стс	CGA	AAT	GGA	ccc	AGA	TGC	TTA	ATG	AAC	GGC	GAG	TGG	GGC	TAC	CAC	TTC	GAA	GAC	ACC	GCG	GGA	GCT	TAC		
CAC	AGA	CCT	CGA	ccc	GAG	GCT	TTA	CCT	GGG	TCT	ACG	AAT	TAC	TTG	CCG	CTC	ACC	CCG	ATG	GTG	AAG	CTT	CTG	TGG	CGC	CCT	CGA	ATG		450
٧	s	G	A	G	L	R	N	G	P	R	С	L	M	N	G	Ε	W	G	Υ	н	F	Ε	D	Т	А	G	A	Υ	L	
стс	AAC	CGC	ACT	CTA	TGG	GAT	CGG	TGC	GAG	GCG	ccc	сст					TGG	AAT	GTG	ACG	стс	TTC	TCG	CTG	сте	GTG	GCC	GCC.	TCC	
GAG	TTO	GCG	TGA	GAT	ACC	CTA	GCC	ACG	СТС	CGC	GGG	GGA			CAG		ACC	TTA	CAC	TGC	GAG	AAG	AGC	GAC	GAC	CAC	CGG	CGG	AGG	540
L	N	R	Т	L	W	D	R	С	Ε	A	Р	Р	R	٧	٧	P	W	N	<b>V</b>	Т	L	F	s	L	L	. <b>v</b>	A	A	s	
TGC	CTO	GAG	ATA	GTA	CTG	TĢT	GGG	ATC	CAG	CTG	GTG	AAC	GCG	ACC	ATT	GGT	GTC	TTC	TGC	GGC	GAT	TGC	AGG	AAA	AAA	CAG	GAC	ACA	CCT	
ACG	GAC	стс	TAT	CAT	GAC	ACA	ССС	TAG	GTC	GAC	CAC	TTG	CGC	TGG	TAA	CCA	CAG	AAG	ACG	CCG	CTA	ACG	TCC	TTT	TTT	GTC	CTG	TGT	GGA	630
С	L	Ε	I	٧	L	С	G	I	Q	L	٧	N	A	T	i	G	٧	F	С	G	٥	С	R	K	K	Q	D	T	P	
CAC	TGA	GGC	TCC	AC T	GAC	CGC	CGG	GTT	ACA	CCT	GCT	ccr	TCC	TGG	ACG	стс	ACT	CCE	TTG	стс	GCT	AGA	ATA	AAC	TGC	TTT	GCG	CTC		
GTG	AC T	CCG	AGG	TGA	CTG	GCG	GCC	CAA	TGT	GGA	CGA	GGA	AGG	ACC	TGC	GAG	TGA	GGG	AAC	GAG	CGA	TCT	TAT	TTG	ACG	AAA	CGC	GAG		720
н	· (	(5	eG	J	D	Ν	0:	2	7	)																				
AAA											7.	ンィ	SC	) ,	12	7														
TTT	TTI	TIT	TTT	TTT	TTG	/ /3	ರ	ت	×	<i>.</i>	ルし	ノ i `	<b>V</b>	<b>'</b> .	104	J														

																									HI	۷ر	Bi	186	0_		
GĢCAC	GΑ	GAG	SAT	TGT	CGG	CŢ	GCG	GG	TAI	[AT]	TCC.	AAT	TCC	CCG	TCT	CCT	CAT	GAA	TAT	GAAC	TGA	AGO	GC	CTO	SACO	СТС	GAA	GTG	GTT	CT	90
CCGTG	CT	CTO	TA	ACA	GCC	GA	CGC	:cc	ATA	ATA	AGG	TTA.	AGG	GGC	AGA	GGA	GTA	CTT	ATA	CTTO	CACT	TTC	CGA	GAC	CTGC	GGA	CTT	CAC	:CAA	IGA	
				;	١,																										•
AAGCA																															180
TTCGT	CC	C G	TTT	TAC	ccc	AG	AGC	CT	TC	4CA	CCT	CCG	ACG	GAT	TCA	ACA.	AAC	GAC	TAA	GGC	GAA(	CGT(	SAA	ACC.	rca:	TAT	rago	CACT	TGT	TAT	
				M	G	S	F	?	Κ	С	G	G	С	L	S	С	L	L	I	Ρ	L	Α	L	W	S	Ī	I	٧	N	1	
TTATI																			+					,							270
AATAA	CA	TA.	AAG	GGC	:11/	AC C	CGT	TTT	GA	AGG.	ATA	CGT	AGG	TCG	TTA	TTT	GAG	TGG	TTG	ATG	CAC	ACC	ΔTΑ	AAAI	CTT	CCT	TAGA	4C A	4AGA	AGT	
L l	-	Y	F	ρ	N	G	(	3	T	S	Y	A	s	s	N	κ	Ł	Ţ	N	Υ	٧	W	Y	F	ε	G	i	C	F	S	
GGCAT	TC.A	TG	ATG	CTI	ΓΑΤΑ	AGT	AA(	CAA	AC A	GTŢ	CTT	CTG	GTA	ÇTG	GAG	AAT	AAT	AAC	AAC	TAT	AAA	TGT	TGC	CAG.	AG:T	GAA	AAC	TGC	AGC	AAA	360
CCGT	GT	AC	TAC	GA	ATA	TCA	TTO	STI	GT	CAA	GAA	GAC	CAT	GAC	CTC	TTA	TTA	TTG	TTG	ATA	TTT.	ACA.	ACG	GTC	TCA	CTT	TTG	ACG:	TCGT	TTT	
G	ſ	M	М	L	i	٧	, .	Т	T	٧	L	L	٧	L	Ε	N	N	N	N	Y	K	С	С	۵	S	ε	N	С	S	K	
AAAT	<b>4</b> T G	TG	A C A	AC T (	GCT	GTÇ	AA'	T T #	ATC	TTŢ	тст	TÇC	CTC	GGA	ATT	GCT	TTT	TCT	GGA	TAC	TGC	CTG	GTC	ATC	TCT	GCC	TTG	GGT	CTT	GTC	450
TTTA	TAC	AC	TGI	GA	CGA	CAG	TT	AAI	TAG	AAA	AGA	AGG	GAG	CCT	TAA	CGA	ΑΑΑ	AGA	CCT	ATG	ACG	GAC	CAG	TAG	AGA	CGG	AAC	CCA	GAA	CAG	
ĸ	Y	٧	Ţ	L	L	5	;	I	I	F	s	s	L	G	ı	Α	F	s	G	Υ	¢	L	٧	i	s	A	L	G	L	٧	
CAAG	GGO	:CA	TAI	rtgi	CCG	CAC	:cc	TTO	GAT	GGC	TGG	GAG	STAT	rgc 1	TTT	GAA	GGC	ACI	GCT	GGĄ	CGT	TTC	стт	ĄСА	GAT	TCT	AGC	ATA	TGG	ATT	540
GTTC										1_							_		<del></del> -	$\overline{}$				_						•	340
Q	G	Р	Υ	c	R	1	Γ I	L	Ð	G	W	Ε	Y	A	F	Ε	G	Т	A	G _.	R	F	Ļ	T	D	s	S	I	W	l	
CAGT	GCI	:TG	GA	ACC	TGC	AC/	ATG	TTI	GTG	GAG	TGC	GAAC	CATO	CAT	TTT#	TTT	TCC	ATI	сто	ATA	ACC	стс	AGT	GGG	CTT	CAA	GŢG	ATC	ATC	TGC	630
GTCA																															630
a	С	L	Ε	Р	А	. }	4	٧	٧	Ε	W	N	I	ı	L	F	s	I	L	1	Ŧ	L	s	G	L	a	٧	ı	I	С	
CTCA	TC.	A C: A	CT.	ACT	CAT	GC (	ΔΔΓ	ТΔ	TCC	ΔΔ6	ATA	ACTO	GTG.	TGG	AAGO	TAT	rtca	AGT	SATO	TTC	CAG	ССТ	GGA	ATC	ATT	TGA	ATA	AGG	ACA	AAA	700
GAGT																															720
L	I	R	٧	٧	М	i (	a	L	S	K	1	L	С	G	s	Y	S	٧	1	F	Q	Р	G	i	i	. (	SE	D	IJ	> <i>V</i>	٥١٤
TGTT	TT	CCA	TT.	ATC	AAG	AC.	ATG	GC	CAT	CTA	ATC.	TAA	ATA	TTA	TAT	CAAC	TG.	TGT	TAGA	ACTI	GAG	GGC	AAT	ATT	GAA	AAT	GAT	GGT	GCT	TTC	810
ACAA																															
																			•												
TGCA																															
ACGT	AA	AC	CAC	AAA	TAA	AC.	ATT	TT	TTA	AAA	CGT	CAG	GAG	TGA	CGT	GTA	CGT	TCA	TAT(	GGT	SGGA	AGG	TAA	ATC	ATA.	AU A	IAAA	AII	ĻΑΙ	IAI	

HUVBB80

AGAAACCTTTATGCAATATGTATATTGCAACATTATTTAATATTCTGGAAAATTGGAAACCCCCAAAATTCTAACCTCAAA

TCTTTGGAAATACGTTATACATATAACGTTGTAATAAAATTATAAGACCTTTTAACCTTTGTGGGGTTTTAAGATTGAGTTT

1071

Page 1

53

HJACE54

TITGTGGAGGGCAGCAGAGAGTACCCAGCTGGACATCCTTTCCTGCTGATGAGCCCCAGGCTGGAGGTGCCCTGCTCACATGCT	CTTCCC	90
AAACACCTCCCGTCGTCTCTCATGGGTCGACCTGTAGGAAAGGACGACTACTCGGGGTCCGACCTCCACGGGACGAGTGTACGA	GAAGGG	
M S P R L E V P C S H A	L P	•
·	<b></b>	
CAGGGTCTCTCGCCTGGGCAGGTCATCATAGTACGGGGACTGGTCTTGCAAGAGCCGAAGCATTTTACTGTGAGCCTGAGGGAC	<del></del>	180
GTCCCAGAGAGCGGACCCGTCCAGTAGTATCATGCCCCTGACCAGAACGTTCTCGGCTTCGTAAAATGACACTCGGACTCCCTG	GTCCGA	
QGLSPGOVIIVRGLVLQEPKHFTVSLRD	A D	
GCCCATGCTCCTGTGACACTCAGGGCCTCCTTCGCAGACAGA	TCAGCC	270
CGGGTACGAGGACACTGTGAGTCCCGGAGGAAGCGTCTGTCT	SAGTEGG	
A H A P V T L R A S F A D R T L A W I S R W G Q K K L I	S A	
CCCTTCCTCTTTTACCCCCAGAGATTCTTTGAGGTGCTGCTCCTGTTCCAGGAGGGAG	GGGCTG	360
GGGAAGGAGAAATGGGGGTCTCTAAGAAACTCCACGACGAGGACAAGGTCCTCCCTC	CCCGAC	
PFLFYPORFFEVLLLFOEGGLKLALNGO	G L	
GGGGCCACCAGCATGAACCAGCAGGCCCTGGAGCAGCTGCGGGAGCTCCGGATCAGTGGAAGTGTCCAGCTCTACTGTGTCCAC	TCCTGA	450
CCCCGGTGGTCGTACTTGGTCGTCCGGGACCTCGTCGACGCCCTCGAGGCCTAGTCACCTTCACAGGTCGAGATGACACAGGTC	AGGACT	- TC
G A T S M N Q Q A L E Q L R E L R I S G S V Q L Y C V H	S .	NO:
AGGATGGTTCCAGGAAATACCGCAGAAAACAAGAGTCAGCCACTCCCCAGGGCCCCACTCTCCTCCCCTCATTAAACCATCCAC	CTGAAC	540
TCCTACCAAGGTCCTTTATGGCGTCTTTTGTTCTCAGTCGGTGAGGGGTCCCGGGGTGAGAGGAGGGGGAGTAATTTGGTAGGTC	GACTTG	
ACCAGCACATCAGGGCCTGGTTCACCTCTGGGGTCACGAGACTGAGTCTACAGGAGCTTTGGGCCTGAGGGAAGGCACAAGAG	FGCAAAG	630
TGGTCGTGTAGTCCCGGACCAAGTGGAGACCCCAGTGCTCTGACTCAGATGTCCTCGAAACCCGGACTCCCTTCCGTGTTCTCA	ACGTTTC	
GTTCCTCGAACTCTGCACCTTCCTCCACCAGGAGCCTGGGATATGGCTCCATCTGCCTTCAGGGCCTGGACTGCACTCACAGAC	3GC AAGT	- 720
CAAGGAGCTTGAGACGTGGAAGGAGGTGGTCCTCGGACCCTATACCGAGGTAGACGGAAGTCCCGGACCTGACGTGAGTGTCTC	CGTTCA	
	~ A A T A A	
GTTGTAGACTAACAAAGATACTCCAAAATACAATGGCTTAAAGAATGTGGTCATTTATTCTTTATTTA	<del>+</del>	810
CAACATCTGATTGTTTCTATGAGGTTTTATGTTACCGAATTTCTTACACCAGTAAATAAGAAATAAAT	2111A11	
ATAAATAAGGTTATTTAAAAAAAAAAAAAAAAAAAAAA	}	
TATITATICCAATAAATAAATITTTTTTTTTTTTTTTTT		

HROAD63

Päge 1

GCACGAGAGACGACATCAGAGATGAGGACAGCATTGCTGCTCCTTGCAGCCCTGGCTGTGGCTACAGGGCCAGCCCTTACCCTGCGCTGC

90
CGTGCTCTCTGCTGTAGTCCTACTCCTGTCGTAACGACGAGGAACGTCGGGACCCGACACCGATGTCCCGGTCGGGAATGGGACGCGACG

MRTALLLAALAVATGPALTLRC'

H V C T S S S N C K H S V V C P A S S R F C K T T N T V E P

CTGAGGGCTTCCCCGAAAGTCTGGGACCAGGTCCAGGTGGGCATGGAATGCTGATGACTTGGAGCAGGCCCCACAGACCCCACAGAGGAT
GACTCCCGAAGGGGCTTTCAGACCCTGGTCCAGGTCCACCCGTACCTTACGACTACTGAACCTCGTCCGGGGTGTCTGGGGTGTCTCCTA

LRASPKVWDQVQVGMEC. (SEQIDNO: 30)

GAAGCCACCCCACAGAGGATGCAGCCCCCAGCTGCATGGAAGGTGGAGGACAGAAGCCCTGTGGATCCCCGGATTTCACACTCCTTCTGT
CTTCGGTGGGGTGTCTCCTACGTCGGGGGTCGACGTACCTTCCACCTCCTGTCTTCGGGACACCTAGGGGCCTAAAGTGTGAGGAAGACA

AI CAGOT1500 - DW. - DO21700A2 IA

HMW6546

AGG	GG	GCC	CGA.	ACC		افالفا	IGAA	1666	IG	AG	TACC	TA	AGC	CGG	AGC	SGGC	STA	GAG	GCGC	GGCC	GG	ACC	CCC	:110	TG	ACC:	TCCA	AGTO	SCCG	~~
TC	CC	CGG	GCT	TGG	GAG	CACA	ACTI	rccc	ACC	TC/	ATGO	SAT	rcc	CC.	TCGC	ccc	CAT	CTC	GCC	cccc	CCC	GTGC	seco	AAG	AC	TGG	4GG1	CAC	GGC	90
				ţ																										
cce	GC	CTC	AAG	ATC)	AGA	CATO	GCC	CAG	SAAC	TTO	SAAG	GAG	: TT(	SGC	GGGA	ACG6	SCT	GCC	GCC	GGG	scco	cac	aggo	ATG	1881	CACC	יפכר	CTO	SAAG	
	-+				+			<del></del>		<del></del>	<del></del>		+			-+				<del></del>			<del></del>			GTGG				18
000	CG	SAG	116	IAG		JAC		3010	, .	3AAL			SAAL		1	666	.GA	. 666	3666	3666	.661	1000		HAL		3   (s)		SGAL	.110	
						М	Α	Q	N	L	Κ	D	L	A	G	R	L	Ρ	Α	G	Ρ	R	G	М	G	T	A	L	κ	
ĊŦĠ	TŤſ	SC FO	ağ g (	GCC	i gğı	isce	Gig	GCC	TAC	:661	ete.	ירנו	· C A A	ı TC	rete	:דדר	. A C (	re Ti	GA A	reer	.cc	: C A C	.VCV	crr	ΔΤί	CTTO	`TTC	· A A T	ררר	
			-+					-					-+					<del></del>		<del></del>			+	+		GAAG				27
GAC	^^	JOA			3000	3000	3CAC			300	·CAC		) L I I	AGA	4CAL	AAU	3166	3 L A					1161	CGG	HAU	JAAC	JAAG	SIIA	1666	
L	L	L	G	Α	G	A	٧	Α	Υ	G	٧	R	Ε	S	٧	F	T	٧	Ε	G	G	Н	R	A	I	۴	F	N	R	
ATC	GG1	rgg.	AGTO	SCAG	GCAC	GGAC	CACT	ATC	СТО	seco	GAG	GGG	CTI	CAC	:TTC	AGG	ATO	2001	rtgo	STTC	CAG	TAC	ccc	ATT	ATI	CTAT	GAC	ATT	'cgg	
			-+-					+	<del></del>				+			-+-							+			GATA		<del></del>	+	36
.,,,	••,						,,,,,				,		10//		,,,,			300,							' ^ \	3016				
I	G	G	٧	a	Q	D	Т	I	L	Α	Ē	G	L	Н	F	R	I	Ρ	W	F	Q	Υ	Р	I	1	Y	D	i	R	
GCC	AGA	ACCI	rcg/	AAA	AATO	тсс	TCC	ССТ	ACA	GGC	TCC	AAA	GAC	CTA	CAG	ATG	GTO	SAAI	ATO	TCC	сто	CGA	GTG	TTG	TC1	TCGA	ccc	AAT	GCT	
			-+					<del></del>			<del></del>		+								-	-	+			AGC T		<del></del>	-+	45
										-	,,,,,						. •		. , , , ,	,,,,,,				,,,,	,,,,,		-			
Α	R	Р	R	K	I	S	S	Ρ	T	G	S	K	D	L	Q	М	٧	N	i	S	L	R	٧	L	S	R	Р	N	Α	
CAG	GAG	CTI	rcci	AGC	ATO	STAC	CAG	CGC	CTA	GGG	CTG	GAC	TAC	GAG	GAA	CGA	GTO	STTO	CCG	TCC	ATT	GTC	AAC	GAG	GT(	SCTO	AAG	AGT	GTG	
GTC	CTO	GAA	AGGA	TCG	TAC	ATG	GTC	GCG	GAT	CCC	GAC	CTG	ATG	CTO	CTT	GCT	CAC	AAC	GGC	AGG	TAA	CAG	TTG	CTC	CAC	GAG	TTC	TCA	CAC	54
	_																													
a	Ε	L	Р	S	М	Y	a	R	L.	G	L	D	Y	Ε	Ε	R	٧	L	Р	S	I	٧	N	Ε	٧	L	K	S	٧	
GTG	GCC	AAG	STTE	AAT	GCC	TCA	CAG	CTG	ATC	ACC	CAG	CGG	GCC	CAG	GTA	TCC	сто	STT	ATC	CGC	CGG	GAG	CTG	ACA	GA(	SAGG	GCC	AAG	GAC	
CAC	CGG	TTC	AAG	TTA	CGG	AGT	GTC	GAC	TAG	TGG	GTC	GCC	CGG	GTO	CAT	AGG	GAC	AAC	TAG	GCG	GCC	CTC	GAC	TGT	CTO	TCC	CGG	TTC	CTG	63
٧	A	K	F	N	A	S	a	L	I	Ŧ	a	R	A	Q	٧	S	L	٤	I	R	R	Ε	L	T	Ε	R	Α	K	D	
TTC	AGC	стс	ATC	CTG	GAT	GAT	GTG	GCC	ATC	ACA	GAG	CTG	AGC	TTT	AGC	CGA	GAG	TAC	ACA	GCT	GCT	GTA	GAA	GCC.	AAA	CAA	GTG	GCC	CAG	
AAG	TCG	GAG	TAG	GAC	CTA	CTA	CAC	CGG	TAG	TGT	CTC	GAC	TCG	AAA	TCG	GCT	CTC	ATG	TGT	CGA	CGA	CAT	CTT	CGG	TTT	GTT	CAC	CGG	GTC	72
			_							-																				
F	S	L	I	Ļ	D	D	٧	A	i	T	Ε	L	S	F	S	R	Ε	Y	T	A	Α	٧	Ε	A	K	Q	٧	Α	Q	
CAG	GAG	GCC	CAG	CGG	GCC	CAA	TTC	TTG	GTA	GAA	AAA	GCA	AAG	CAG	GAA	CAG	CGG	CAG	AAA	ATT	GTG	CAG	GCC	GAG	GGT	GAG	GCC	GAG	GCT	
	<del></del>		+	$\longrightarrow$				+					+	<del></del> +		<del></del>			-+		<del></del>		+			CTC		+	<del>+</del>	810
۵	Ε	A	0	R	Α	Q	F	L	٧.	Ε	K	Α	K	0	Ę	Q	R	Q	K	I	٧	a	A	Ε	G	Ε	A	E	Α	
GCC.	AAG	ATG	CTT	GGA	GAA	GCA	CTG.	AGC.	AAG	AAC	сст	GGC	TAC	ATC	AAA	стт	CGC	AAG	ATT	CGA	GCA	GCC	CAG	AAT	ATC	TCC	AAG.	ACG.	ATC	
			+-			-+-		<del></del>					+	+		-+-		<del></del>	-+				+			AGG		+	<del>+</del>	900
				'	-,,		,		. , .		-J-J-A	0	~ r Q	<i>-</i> ~ u		<b>J</b> AA	<b>3</b> C C		144	361			J , C			,,,,,,			•	
Δ	×	м		C	F	Δ	4	S	X	N	P	C	Y	1	ĸ		0	V		D	٨	Δ	Λ	N		ς	ĸ	T	ı	

HMWF546

GCCACATCACAGAATCGTATCTATCTCACAGCTGACAACCTTGTGCTGAACCTACAGGATGAAAGTTTCACCAGGGGAAGTGACAGCCTC
CGGTGTAGTGTCTTAGCATAGATAGAGTGTCGACTGTTGGAACACGACTTGGATGTCCTACTTTCAAAGTGGTCCCCTTCACTGTCGGAG

ATSQNRIYLTADNLVLNLQDESFTRGSDSL.

(SEQIDNO:16 ATCAAGGGTAAGAAATGAGCCTAGTCACCAAGAACTCCACCCCCACAAGAAGTGGATCTGCTTCTCCAGTTTTTGA TAGTTCCCATTCTTTACTCGGATCAGTGGTTCTTGAGGTGGGGGTGTTCTTCACCTAGACGAAGAGGTCAAAAACT

IKGKK (SCOIDNO: 31)

HAF. WOL

GGCA	CGA	AGAT	GAC	ATC	ACT	ΓĂΑG	TGG	CCG	AŢC	TGC	AÇA	GAG	CAG	GCC	AGG	AGC	AAC	CÁC	ACA	GGC	TTC	CTO	CAC	ATG	GAC	TGC	GA	GAT	CAA	00
CCGT	GCT	rc T/	CTG	TAG	TGA	T.T.C	ACC	GGC	TAG	ACG	TGT	сто	GTC	CGG	TCC	TCG	TTG	GTG	TGT	CCG	AAG	GAC	GTG	TAC	сте	ACG	СТ	CTA	GTT	90
				1																				М	D	С	Ε	ſ	K	•
GGGC	CGC	ccc	TGC	TGC	ATC	GGC	ACÇ	AAG	GĢC	AGC	TGT	GAG	ATC	ACC	ACC	ÇGG	GAA	TAC	TGT	GAG	TTC	ATO	CAC	GGC	TAI	TTC	CA	TGA	GGĄ	180
cccc	GCC	GGG	ACG	ACG	TAG	CCC	TGG	TTC	CCG	TCG	ACA	CTC	TAG	TGG	TGG	GCC	CTT	ATG	ACA	стс	AAG	TAC	GTG	CCG	ATA	AAG	GT	ACT		100
G	R	Р	С	С	ı	G	Т	κ	G	s	С	Ε	j	T	T	R	Ε	Y	С	Ε	F	M	Н	G	Y	F	Н	Ε	Ε	
AGCA	ACA	ACTO	TGC	TCC	CAG	GTO	AGG																				:cc	GGG	стс	070
TCGT	TGT	GAG	ACG	AGG	GTC	CAC	TCC	GCT	CCG	TCC	GGA	CCI	CAT	CAC	стс	CTC	TCC	TGC	GAC	ccc	TAC	CGI	CGG	ACG	ACC	CCC	GC	CCC	GAG	270
Α	T	L	С	s	Q	٧	R	R	G	R	Р	G	v	V	Ε	Ε	R	τ	L	G	M	A	A	С	W	G	R	G	S	
ACGO						+			-					<del></del>					-+-		+			<del></del>		$\overline{}$		+	+	360
TGC	TGA	GGG	AGG	GTA	CAG	1333	CGG	AGT	CTG	AGT	CCC	ACC	SAAG	ACC	CCG	CGA	CTC	GTG	GTA	TAC	GGG	TAA	GGG	TCC	ACC	STGA	IC A	AAA	ССТ	
R	ĭ	P	S	Н	٧	G	A	S	D	S	G	С	F	W	G	A	Ε	Н	Н	M	P	l	Ρ	R	С	T	٧	L	0	
CAAG	GTO	STG	TGG	GCT	GCT	rgcc	TTC	CTC	AĄC	сст	GAG	GTO	CCA	GAT	CAG	;; <b>;;</b>	TAC	AGG	TÇT	GGC	TGT	CTC	111	тсс	TAC	ATO	TT	GGG	TAA	450
GTTC	CAC	ACA	ACC	CGA	CGA	CGG	AAG	GAG	TTG	GGA	CTC	CAG	GGT	CTA	GTC	ÀAA	ATG	TCC	AGA	CCC	ACA	GAG	AAA	AGG	ATO	TAC	AA	CCC		430
κ	٧	С	w	A	A	Ά	F	L	N	P	Ε	٧	Ρ	D	Q	F	Y	R	s	G	С	L	F	s	Y	М	L	G	κ	
GAGG	TCC	TCA	ATG	ccc	CCG	SAAC	CCG	ACC	ССТ	GTG	ATG	GAC	ACC	CAG	GCG	GAC	ccc	TGG	GĢA	AAC	GT1	rcci	GGG	CCA	GGG	TAT	GG	TCG	GTÇ	540
стсс	AG	SAG	TAC	GGG	GGC	TTG	GGC	TGG	GGA	CAC	TAC	CTC	TGG	GTC	CGC	CTG	GGG	ACC	ССТ	TTC	ĊA/	\GG/	ccc	GGT	CCC	ATA	CC	AGC		540
R	s	S	м	Р	Р	N	₽	T	Р	٧	м	D	T	Q	A	D	P	W	G	K	٧	Ρ	G	Ρ	G	Y	G	R	s	
CAAC	ст	seco	AAG	ACT	ACT	GCT	ССТ	GAA	GTG	тст	GGA	TG	AGG	cce	CTG	CCT	GGT	GTG	TCC	сто	ccc	CAG	TGT	GGG	TGO	ACT	GC	CCT	CGG	630
GTT	GAC	GGG	TTC	TGA	TGA	\CGA	GGA	CTT	CAC	AGA	CCT	AC1	TCC	GGC	GAC	GGA	CCA	CAE	AĞG	GAG	GGG	GTO	ACA	CCC	ACC	TGA	CG	GGA	GCC	
N	L	ρ	κ	τ	T	A	Р	E	٧	s	G		Œ	<b>(</b>	IC	7 1	<b>W</b> .	36	2)	)										
TGTO	TTO	STG	GTC	TTT	TCA	AAAT	GAC	ATC	CÇT	GAA	GGG	GAÇ	сто	GAG	GAA	GGT	GGT	CCG	GÇT	GGC	ACC	CTA	TCG	ic ,	704	<u> (</u> Sc	G	I	ر ت	NO;
ACAG	AAC	CAC	CAG	AAA	AGT	TTA	CTG	TAG	GGA	CTT	ccc	ст	GAC	сто	стт	CCA	CCA	GGC	CGA	ccc	TGC	GAT	AGC		<b></b>					

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism refer	red to in the description  Table 1
on page 13 , tine	Further deposits are identified on an additional sheet
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution  American Type Culture Collection	
Address of depositary institution (including postal code and count 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	try)
Date of deposit May 16, 1997	Accession Number 209053
C. ADDITIONAL INDICATIONS (leave blank if not applicab	This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATIONS A	
E. SEPARATE FURNISHING OF INDICATIONS (leave bla The indications listed below will be submitted to the International Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized officer Yvetts Singer  Paralegal Specialis:	Authorized officer
IAPD-PCT Operation:	

Form PCT/RO/134 (July 1992)

REC'D

2 3 FED 1998

### INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

	WIPO PCT									
A. The indications made below relate to the microorganism referr on page13, lime	ed to in the description able 1									
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet									
Name of depositary institution										
American Type Culture Collection										
Address of depositary institution (including postal code and country 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	אי									
Date of deposit	Accession Number									
May 16, 1997	209053									
C. ADDITIONAL INDICATIONS (leave blank if not applicable	e) This information is continued on an additional sheet									
In respect of those designations in which of the deposited microorganism will be mathemention of the grant of the European application has been refused of withdrawn the issue of such a sample to an expert resample (Rule 28(4) EPC).	nde available until the publication of patent or until the date on which the or is deemed to be withdrawn, only by									
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)										
E. SEPARATE FURNISHING OF INDICATIONS (leave blan	k if not applicable)									
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)  The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g "Accession Number of Deposit")										
For receiving Office use only	For International Bureau use only									
This sheet was received with the international application	This sheet was received by the International Bureau on:									
Authorized office verte simple Particial Specialist IAFO-POT Operations	Authorized officer									

Form PCT/RO/134 (July 1992)

### INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 13
B. IDENTIFICATION OF DEPOSIT  Further deposits are identified on an additional sheet  Name of depositary institution American Type Culture Collection  Address of depositary institution (including postal code and country)  12301 Parklawn Drive Rockville, Maryland 20852 United States of America  Date of deposit  May 16, 1997  Accession Number  209054  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
Name of depositary institution American Type Culture Collection  Address of depositary institution (including postal code and country)  12301 Parklawn Drive Rockville, Maryland 20852 United States of America  Date of deposit May 16, 1997  Accession Number 209054  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
Address of depositary institution (including postal code and country)  12301 Parklawn Drive Rockville, Maryland 20852 United States of America  Date of deposit May 16, 1997  Accession Number 209054  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
Address of depositary institution (including postal code and country)  12301 Parklawn Drive Rockville, Maryland 20852 United States of America  Date of deposit May 16, 1997  Accession Number 209054  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
12301 Parklawn Drive Rockville, Maryland 20852 United States of America  Date of deposit May 16, 1997  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
12301 Parklawn Drive Rockville, Maryland 20852 United States of America  Date of deposit May 16, 1997  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
Rockville, Maryland 20852 United States of America  Date of deposit  May 16, 1997  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
Date of deposit  May 16, 1997  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  Accession Number  209054  This information is continued on an additional sheet
Date of deposit  May 16, 1997  C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
C. ADDITIONAL INDICATIONS (leave blank if not applicable)  This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (1) the indications are not for all designated states)
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")
For receiving Office use only ————————————————————————————————————
For receiving Office use only  This sheet was received with the international application  This sheet was received by the International Bureau on:
This sheet was received with the international application  This sheet was received by the International Bureau on:
This sheet was received with the international application  This sheet was received by the International Bureau on:

#### 59/2

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

REC'D	2 3	FEB	1998
WIPO	PC	T	

A. The indications made below relate to the microorganism	referred to in the description									
	Table l									
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet									
Name of depositary institution										
American Type Culture Collection										
Address of depositary institution (including postal code and	(country)									
12301 Parklawn Drive										
Rockville, Maryland 20852 United States of America										
United States of America										
Dur of descript	Accession Number									
Date of deposit May 16, 1997	209054									
C. ADDITIONAL INDICATIONS (leave blank if not app	olicable) This information is continued on an additional sheet									
of the deposited microorganism will the mention of the grant of the Euro application has been refused or with	which a European Patent is sought a sample be made available until the publication opean patent or until the date on which addrawn or is deemed to be withdrawn, only expert nominated by the person requesting									
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)										
E. SEPARATE FURNISHING OF INDICATIONS (lea	ve blank if not applicable)									
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")										
For receiving Office use only	For International Bureau use only									
This sheet was received with the international applicat	ion This sheet was received by the International Bureau on:									
l X										
Authorized officer Visite Simmis	Authorized officer									

Form PCT/RO/134 (July 1992)
BNSDCCID: <WO__9831799A3_IA>

10

15

20

25

30

35

What is claimed is:

- 1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y; or a nucleotide sequence complementary to said isolated polynucleotide.
- 2. The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:X encoding the receptor polypeptide of SEQ ID NO:Y.
- 3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length.
- 4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO:  $\mathbf{X}$ .
  - 5. The polynucleotide of claim 1 which is DNA or RNA.
- 6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a receptor polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:Y when said expression system is present in a compatible host cell.
  - 7. A host cell comprising the expression system of claim 6.
- 8. A process for producing a receptor polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
  - 9. A process for producing a cell which produces a receptor polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a receptor polypeptide.
  - 10. A receptor polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:Y over its entire length.

- 11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:Y.
  - 12. An antibody immunospecific for the receptor polypeptide of claim 10.
- 13. A method for the treatment of a subject in need of enhanced activity or expression of receptor polypeptide of claim 10 comprising:
- (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
  - (b) providing to the subject polynucleotide of claim 1 in a form so as to effect production of said receptor activity in vivo.
- 14. A method for the treatment of a subject having need to inhibit activity or expression of the receptor polypeptide of claim 10 comprising:
  - (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
  - (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
- 20 (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.
  - 15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the receptor polypeptide of claim 10 in a subject comprising:
  - (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said receptor polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the receptor polypeptide
   30 expression in a sample derived from said subject.
  - 16. A method for identifying agonists to the receptor polypeptide of claim 10 comprising:
    - (a) contacting cells produced by claim 9 with a candidate compound;
- (b) determining whether the candidate compound effects a signal generated by activation of the receptor polypeptide.

35

and

15

- 17. An agonist identified by the method of claim 16.
- 18. The method for identifying antagonists to the receptor polypeptide of claim 10 comprising:
  - (a) contacting said cell produced by claim 9 with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.
  - 19. An antagonist identified by the method of claim 18.


- 20. An isolated receptor polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC; and
- (b) a nucleotide sequence complementary to the nucleotide sequence of (a).
- 21. A recombinant host cell produced by a method of Claim 9 or a20 membrane thereof expressing a receptor polypeptide.

7 1	10 M T T T T T T W T W T	2 11 0 3 0	0 0 0	30 30 7. H	\$ <b>4</b>
Ω Σ	PLLLLPLLWG	9 <b>(</b> )	E X A A X E A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S A X S	LOVOKSVTVOEGLOB-1.aa	S M HMACR/0.AA S L OB-1.aa
		- B-	- 9-	07	- 8-
C V E	V R C S F S V P C S F S	YPVDSQTD YPWRSWYS	S D P V H G Y W F R A G N D I S W S P P L Y V Y W F R D G E I P Y Y	K A E V	VAT HMACR70.AA VAT OB-1.aa
		-8-	100	110	120
N N N D D D	NNPAWAVQE NNPDRRVKP	ET QGRFRL ET QGRFRL	POTKNC VOKKNC	TLSIRDARMSDAGR SLSIGDARMEDTGS	SDAGR HMCR70.AA EDTGS OB-1.aa
,		130	140	150	160
Y F F R Y F F R	Х Ж В В В В В В В В В В В В В В В В В В	NIKWNYKYDO DVKYSYOONK	SV N N L E V T A L	TEKPDIHFLEP	HYPCR70.AA L E OB-1.aa
		170	180	190	200
S 1 G 1 R 1	RPTRLSC	SLPGSCEAG	P P L T F S W T G N A	ALSPLDPETTRS	HMACR70.AA 8 S OB-1.aa

FIG. IA

210 220 230 240 142NVTYPHHMACR70.AA 196 SELTLTPRPEDHGTNLTCQMKRQGAQVTTERTVQLNVSYA OB-1.aa	250 270 280 280 147 PQNLTVTVFQGEGTASTALGNSSSLSVLEGQSLRLVCAVD HVACR70.AA 236 PQ TITIFR-NGIALEILQNTSYLPVLEGQALRLCDAPPOB-1.aa	290 310 320 187 SNPPARLSWTWRSLTLYPSOPSNPLVLELQ-VHLGDEGEFHMACR70.AA 273 SNPPAHLSWFQGSBALNATPISNTGILELRRVRSAEEGGFOB-1.aa	226 TCRAONSLGSQHVSLNLSHMACR70.AA 313 TCRAOHPLGFLOLFLNLSVYSLPOLLGPSCSWEAEGLHCROB-1.aa	244
ने ने	5. 17	27.	3.1	35

FIG.1B



F16.10




FIG.2A





FIG.2B

MRC-0X44.aa ·HTEDK48.aa PETA-3.aa

245 218 252 FIG.2C



SUBSTITUTE SHEET (RULE 26)

160 C. HPWAE25.aa C NAG-2.aa C TALLA-1.aa	200 TK HPWAE25.aa SL NAG-2.aa	40 G HPWAE25.aa T NAG-2.aa G TALLA-1.aa	280 L HPWAE25.aa - NAG-2.aa - TALLA-1.aa	HPWAE25.aa NAG-2.aa TALLA-1.aa
WNTTMKGLKC WSIIOTDFRC VDHVQRSLSC	190 2 V T N T ± N E T C T T C T C T C T C T C T C T C T C	230 A V T V 3	270 I S P L P L P L L	
140 G S Q E D F T Q V G T Q G N V G L T N A N G N D E R S R A	180 S A F P P F C C N D D R V - P D S C C L E F G I - P P S C C M N E	220 F N Q L L Y D I R T N Y E T V K V W L Q E N Y D L V T S F M E T N	260 V H V S V L Q S T S	300 DR-I. TY-CA QYEMV
130 V P A I K K D Y Q O D L K K G L H L Y L R T Y T D A M Q T Y	170 - F E D S P Y F K E N W F E V Y N A T - W S T S P Y F L E H	210 V E G C K A P C V A A T K V N Q K G C	250 G L E L A - A M N C - G L T F A M T M Y C - S Q L I G M L L A C C	290 A P W Q A A V I G G G Q V V K A D R F I T A N
116 HFLTLLV 113 RY A 113 DT F	151 G F T N Y T D 149 G V S N Y T D 147 G V Q N Y T N	190 QKAHDQK 180 HAPGTWW 179 QDLHNLT	218 V A A G I G - 213 A L V Q I L - 214 V A F G I A F	253 P H G N C E E

SUBSTITUTE SHEET (RULE 26)




FIG.4A




FIG. 4B




FIG.5A

11/27 HSBBF02.aa TALLA-1.aa Y C L HSBBF02.aa 240 FOLIGIES SOLIGME 230 AGISFGVAC AGVAFGIAF 220 日 で 日 日 NEGCFIKVMTII OKGCYDLVTSFM NNOYEI ANOYEM 210 250 S R R

FIG 5B

234 233




FIG. 6A




FIG.6B

		10	20	30	40
₩ ₩	MARACLQAV MARACLQAV	K Y L M F A K Y L M F A	WLGGCGVLG	VGIWLAATO( VGIWLAATO(	HIPBA27.aa
44	FATLSSSFI	50 PSLSAANLLII PSLSAANLLII	NLLITGAFVMAIGFVGCLGAIKENK	70 VGCLGAIKEI VGCLGAIKEI	1 80 1 K HTPBA27.aa 1 K NAG-2.aa
ᄧᄧ	CLLTTFFLLCCLLTFFLL	) LLLVFLLEAT LLLVFLLEAT	TLFFA	110 Y T D K I D R Y A Q Q D L K Y T D K I D R Y A Q Q D L K	120   K HTPBA27.aa   K NAG-2.aa
22 23	KGLHLYGKGLHLYG	TOGNVGLTNAWSI	IOTDI	150 160 OTDFRCCGVSNYTDWFEVY OTDFRCCGVSNYTDWFEVY	160   Y HTPBA27.aa   Y NAG-2.aa
161	NATRVPDSC NATRVPDSC	70 CLEFS	180 190 200 ESCGLHAPGTWWKAPCYETVKVWLQE ESCGLHAPGTWWKAPCYETVKVWLQE	190 CYETVKVWLO CYETVKVWLO	200 E HTPBA27.aa

FIG.7A

HTPBA27.aa NAG-2.aa NLLAVGIFGLCTALVQILGLTFAMTMYCQVVKADTY NLLAVGIFGLCTALVQILGLTFAMTMYCQVVKADTY 230 201

FIG. 7B

HAIDO59.aa CD9 antigen.aa	HAIDQ59.aa CD9 antigen.aa	HAIDQ59.aa CD9 antigen.aa
40 FGLWFRFGG IGLWLRFDS	MAVGFFGCC	120 FIGKGVATR YSHKDEVIK
20 40 LGENLLFWLAGSAVIAFGLWFRFGG HAIDQ59.aa FGFNFIFWLAGIAVLAIGLWLRFDS CD9 antigen.aa	FYVGLYVLVGAGALMMAVGFFGCC HAIDD59.aa FYTGVYILIGAGALMMLVGFFGCC CD9 antigen.aa	CLLVIFAAEVTTGVFAFIGKGVATR HADDS9.aa FLLVIFAIEIAAAIWGYSHKDEVIK CD9 antigen.aa
20 IKYLLLGFNLLF IKYLLFGFNFIF	SPEYFYVGLY NNSSFYTGVY	TE TC L'LVIFA
ORFRGGLRCIK PV-KGGTKCIK	KEL SSEDK KSIFEQETNN	GAMRESOCVLGS GAVQESOCMLG
E E	41 A I 40 Q T	79 G A 80 G A

FIG.84




FIG. 8B




FIG. 3A




FIG. 9B

+ +	<del>,</del> ←	MCTGKCARCVGLSLITLCLVANALLLLVPNGETSWTNT		.aa
, V	4 44	LOVWLMGGFIGGGLMVLCPGIAAVRAGGKGCCGAG	1 Lon.aa 1 80 C HGBGV89.aa C L6H.aa	.aa
	ස් ස්	MLRSVFSSAFGVLGAIYCLSVSGAGLRNGPRCL		ge
<del>, , , , , , , , , , , , , , , , , , , </del>	121	130 140 NGEWGYHFEDTAGAYLLNRTLWDRCEAPPRVVPWNVTLF NGEWGYHFEDTAGAYLLNRTLWDRCEAPPRVVPWNVTLF	160 S HCBGV89.aa S L6H.aa	g
त्न त्न	161	LLVAASCLEIVLCGIQLVNATIGVFCGDCRKKQDTPH. LLVAASCLEIVLCGIQLVNATIGVFCGDCRKKQDTPH	HGBGV89.aa L6H.aa	gg

F. ()




FIG. 11A




FIG. 11E




FIG. 12A

r 50 L - HJACE54.aa - rGalectin-5.aa ? hGalectin-8.aa		HIACES4.aa rGalectin-5.aa hGalectin-8.aa	0 HIACES4.aa rGalectin-5.aa MGalectin-8.aa	HJACE54.aa rGalectin-5.aa hGalectin-8.aa
190 200 	230 240 HAPVTL DIAFHL DLLAGKSKDIALHL	SAPFLFYPQRFFEV PGSMPFSRGQRFEV ITSEPFSPGMYFEM	ALEQLREDRISGSV NLPDINTEEVAGDI ELSSIDTEENGDI	
180   ELTEISRENVP	220 1  EVNANAKSFNV	260 RWG-QKKLI QINNSWGPEERSL FLQESWG-EEERN	300 NGQGLGATSMNQQA DGQHICEYSHRLMN NGVHSLEYKHRFKE	FIG. 12B
170 	Z	RASFADRTLAWIS NPRFDENAVVRNT NPRENIKAFVRNS	LLLFQEGGLKLAU WILCEGHCFKVAV IIYCDVREFKVAV	330 QLYCVHS. QLTHVET HLLEVRSW
46 53 152	46 53 192	52 232 232	83 87 271	127 139 311

	11.	0	20	30	40
न् न	MRTALLLLAALAVAT MRTALLLLLAALAVAT		GPALTLRCHVCTSSSNCKHSVVCPA GPALTLRCHVCTSSSNCKHSVVCPA	SNCKHSVVCSNCKHSVVC	P A HROAD63.aa P A E48 antigen.aa
	- S	C	- 99	07.	<b> -</b> &-
44	SSRFCKTTNTVSSRFCKTTNTV	E P L R E P L R	ASPKV GNLVKKDCAESCTPSYTL	W D O V	OVG HROAD63.aa SSG E48 antigen.aa
	- & -		100	110	120
88 평	MEC. TSSTQCCQED	LCNEKL	HNAAPTRTALAHSALSLGLALSLL	ALSLGLALS	HROAD63.aa L L E48 antigen.aa
r 2	AVILAPSL				HROAD63.aa E48 antigen.aa

FIG. 13

HMMCS46.aa B-cell Recpt Associated Pr.aa	HMGS46.aa B-cell Recpt Associated Pr.aa	HMMSS46.aa B-cell Recpt Associated Pr.aa	HMWS46.aa B-cell Recpt Associated Pr.aa	HMMGS46.aa B-cell Recpt Associated Pr.aa
10 20 30 PULKDLAGRLPAGPRGMGTALKLLLGAG HWGS46.aa	AVAYGVRESVFTVEGGHRAIFFNRIGGVQQ HMMCS46.aa	70 80 90 DTILAEGLHFRIPWFQYPIIYDIRARPRKI DTILAEGLHFRIPWFQYPIIYDIRARPRKI	PTGSKDLQMVNISLRVLSRPNAQELPSM HMCS46.aa	RIGLDYEERVLPSIVNEVLKSVVAKFNARIGLDYEERVLPSIVNEVLKSVVAKFNA
1 M A Q 1	31 A V 31 A V	ម ១ ១	91 8 8 8	121 Y Q 121 Y Q

FIG. 14A

sociated Pr.aa	sociated Pr.aa	sociated Pr.aa	sociated Pr.aa	sociated Pr.aa
HWGS46.aa B-cell Recpt Associated Pr.aa	HMMCS46.aa B-cell Recpt Associated Pr.aa	HMWIS46.aa   B-cell Recpt Associated Pr.aa	HWKS46.aa B-cell Recpt Associated Pr.aa	HWMSS46.aa B-cell Recpt Associated Pr.aa
170 180 ELTERAKDFSLILD ELTERAKDFSLILD	200 210 Y T A A V E A K Q V A Q Q E A Q R A HWGS46.aa Y T A A V E A K Q V A Q Q E A Q R A B-cell Req	230 240 2 K I V Q A E G E A É A A K M L G E 2 K I V Q A E G E A E A A K M L G E	KIRAAQNISKTIATSQNR HMCS46.aa KIRAAQNISKTIATSQNR B-cell Rec	TRGSDSLIKGKK.
LIRR	ம். ம	230 QKIVQAEG OKIVQAEG	260 RKIRAAQNI RKIRAAQNI	ODESF ODESF
160 SQLITQRAQVSL SQLITQRAQVSL	190 DVAITELSFSR DVAITELSFSR	220 1 QFLVEKAKQEQR QFLVEKAKQEQR	250 ALSKNPGYIKLR ALSKNPGYIKLR	280 LYLTADNLVLNL IYLTADNLVLNL
151 151	181	212	241 P	271 271

FIG. 14B

	10 20 30
	MDCEIKGRPCCIGTKGSCEITTREYCEFMH HNEGWO6.aa MDCVITGRPCCIGTKGRCEITSREYCDFMR EGFR related-protein.aa
# #	GYFHEEATLCSOVR RGRPGVVEERTLGMAA HWFGWO6.aa GYFHEEATLCSOVH EGFR related-protein.aa
요 상	CWGRGSRTPSHVGASDSGCFWGAEHHMPIP HNFGWD6.aa
12 45	100 120 120 120 120 120 120 120 1 1 1 1
. 171 88	130 140 150 SYMLGKRSSMPPNPTPVMDTQADPWGKVPG HNFGWO6.aa
151 69	PGYGRSNLPKTTAPEVSGLSLFLHAGILHC FIG.15



Interpolation No PC1765 98/00959

4 01 100		<del> </del>	
IPC 6	C12N15/12 C12N15/85 C12N5/1 C12Q1/68 A61K38/17 A61K48/		C07K16/28
According to	international Patent Classification (IPC) or to both national classific	ation and IPC	
	SEARCHED	,	
	cumentation searched (classification system followed by classificati	on symbols)	
IPC 6	C12N C07K C12Q A61K		
Documentat	ion searched other than minimum documentation to the extent that s	uch documents are included in the	fields searched
Electronia d	ata base consulted during the international search (name of data ba	se and, where practical, search te	rms used)
C. DOCUME	ENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the rela	evant passages	Relevant to claim No.
X	DATABASE GENBANK Accession No. T48852, 8 February "EST70319 from Stratagene place XP002066898 see the whole document		1-12,20
X	DATABASE GENBANK Accession No. G14442, 4 January "STS SHGC-10055" XP000206689 see the whole document	1996	1-3,20
		-/	
X Furth	ner documents are listed in the continuation of box C.	Patent family members a	ure tisted in annex.
"A" docume consid "E" earlier of filing d "L" docume which citation "O" docume other n "P" docume later th	nt which may throw doubts on priority claim(s) or is cited to establish the publication date of another in or other special reason (as specified) and the publication or other special reason (as specified).	cited to understand the princinvention  "X" document of particular releval cannot be considered novel involve an inventive step wh  "Y" document of particular releval cannot be considered to invodocument is combined with	inflict with the application but siple or theory underlying the siple or theory underlying the siple or theory underlying the siple of the claimed invention siple of the claimed invention siple or more other such document one or more other such document one or more other such document one or more other such document of the siple
	June 1998	•	9. 1998
	nailing address of the ISA  European Patent Office, P.B. 5818 Patentiaan 2  NL - 2280 HV Rijawijk  Tel. (+31-70) 340-2040, Tx. 31 651 epo ni,	Authorized officer  Galli, I	

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

PCT/US 98/00959

		PC1/02 98/00959
	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Α	ZANNETTINO A.C.W. ET AL.: "A powerful technique for isolating genes encoding surface antigens using retroviral expression cloning" J. IMMUNOL., vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 see abstract	1-21
Α	ADAMS M D ET AL: "COMPEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 cited in the application see abstract	1-21
A	SIMMONS D. & SEED B.: "Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells" J. IMMUNOLOGY, vol. 141, no. 8, 15 October 1988, pages 2797-2800, XP002066894 see abstract see figure 4	1-21
	CROCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 3 October 1994, pages 4490-4503, XP002066895 see abstract see figure 2	1-21
A	KELM S. ET AL.: "The Sialoadhesins: a family of sialic acid-dependent cellular recognition molecules within the immunoglobulin superfamily" GLYCOCONJ. J., vol. 13, no. 6, December 1996, pages 913-926, XP002066896 see the whole document	1-21
A	DATABASE GENBANK Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document	1-21
	•••	

		PCT/05 9	8/00959
	ation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication; where appropriate, of the relevant passages		Relevant to claim No.
P,X	DATABASE GENBANK Accession No. AA344713, 21 April 1997 "EST50650, H. sapiens cDNA similar to CD33 antigen" XP002066900 see the whole document		1-12,20
P,A	TAKEI Y. ET AL.: "Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta" CYTOGENET. CELL GENET., vol. 78, 1997, pages 295-300, XP002066897 see abstract see figure 1		1-21
•			



inter onal application No.

PCT/US 98/00959

Box I	Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)							
This Inte	emational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:							
1. X 2.	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, and claim 15 (as far as in vivo methods are envisaged) is directed to a diagnostic method, the search has been carried out and based on the alleged effects of the compound/composition.  Claims Nos.:  Decause they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:							
3	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).  Observations where unity of invention is lacking (Continuation of item 2 of first sheet)							
	ernational Searching Authority found multiple inventions in this international application, as follows:							
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.							
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee							
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:							
4. X	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  see further information sheet, subject 1.							
Remark	The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.							

#### INTERNATIONAL SEARCH REPORT

International Application No. PCT/ US 98/00959

#### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-21) - partial

An isolated polynucleotide encoding the HMACR70 cell surface receptor (Seq. IDs 1,18), polynucleotides complementary thereto, fragments thereof, homologs thereof.

An expression system comprising said polynucleotides, a process to produce a cell harbouring said expression system. Said cell or membrane thereof expressing said receptor.

A process for the production of the HMACR70 polypeptide comprising culturing said cell.

Said HMACR70 polypeptide or homologs. An antibody directed against said polypeptide.

Therapeutic and diagnostic applications.

A method to identify agonists or antagonists to said receptors. Said agonists and antagonists.

2. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 2,3,19

3. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 4,20

4. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 5,21

5. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 6,22

6. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 7,23

7. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 8,24

## INTERNATIONAL SEARCH REPORT


International Application No. PCT/ US 98/00959

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

- 8. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 9,10,25
- 9. Clayms: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 11,26
- 10. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 12,27
- 11. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 13,28
- 12. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 14,29
- 13. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 15,30
- 14. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 16,31
- 15. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 17,21



## **PCT**

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶:
C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 31/70, 38/17, 48/00

A2

(11) International Publication Number:

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

60/034,204 60/034,205 21 January 1997 (21.01.97) 21 January 1997 (21.01.97)

US US

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). GENTZ, Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US).
- (74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

Without international search report and to be republished upon receipt of that report.

- (54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS
- (57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВЈ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Сапада	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		
ı							

PCT/US98/00959

5

10

15

20

25

30

35

# POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

#### FIELD OF INVENTION

This invention relates to newly identified polynucleotides and the polypeptides encoded by them, the use of such polynucleotides and polypeptides, and their production. More particularly, the polynucleotides and polypeptides of the present invention relate to specific receptor families described in the specification and known in the art. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

### BACKGROUND OF THE INVENTION

Receptor proteins are found on the membrane of the cells and are generally involved in signal transduction. There are many types of receptor proteins, and for convenience, these proteins are grouped in families based on similarity in structure and function.

For example, the TM4SF superfamily of cell surface proteins, also known as the tetraspan receptor superfamily, is comprised of at least seventeen individual gene products (these include CD9, CD20, CD37, CD53, CD63, CD81, CD82, A15, CO-029, Sm23, RDS, Uro B, Uro A, SAS, Rom-1, PETA3, and YKK8). The TM4SF superfamily is the second largest group in the CD antigen superfamily. Each member of the TM4SF superfamily can be characterized by several putative physical features including four highly conserved transmembrane domains, two divergent extracellular loops, and two short and highly divergent cytoplasmic tails. Expression patterns for members of the TM4SF superfamily tend to be rather broad and can vary widely between members. The functional roles of TM4SF superfamily members are primarily associated with signal transduction events and pathways, but also include cell adhesion in platelets and other lymphocytic and non-lymphocytic cell lines, as well as cell motility, proliferation, and metastasis. In addition, recent evidence suggests that a subset of the members of the TM4SF superfamily may function as potassium channel molecules.

One member of the TM4SF family, CD20, is a four membrane spanning domain cell surface phosphoprotein expressed exclusively on B lymphocytes. Although the precise functional role of CD20 has yet to be determined, it is thought to function primarily as a receptor during B-cell activation. Furthermore, a large number of experimental observations suggest several additional speculative roles for the CD20 molecule. For example, CD20-specific immunoprecipitation of biochemically cross-linked plasma membrane proteins suggests that CD20 assumes a multimeric structural

conformation characteristic of other previously described membrane channel proteins. Further experimentation has revealed that expression of exogenous CD20 on the cell surface specifically increases Ca²⁺ conductance across the plasma membrane. Together, these results suggest that CD20 complexes may function as B-cell specific Ca²⁺ ion channels. In addition, monoclonal antibodies raised against CD20 have been used to stimulate resting B-cells to transition out of the G0/G1 segment of the cell cycle. It has also been demonstrated that CD20 is associated with both serine and tyrosine kinases and, more specifically, that CD20 is associated, although not directly, with the Src family of tyrosine kinases including p56/53lyn, p56lck, and p59fyn.

A second example of a receptor subfamily, called sialoadhesin molecules, belongs to the Ig superfamily of receptor-like molecules. The more than 100 members of the Ig superfamily are generally considered to engage in specific cell-cell interactions through which intercellular communication may occur. In addition to classical protein-protein interactions, intercellular communication may also be mediated through protein-carbohydrate interactions. In fact, all members of the sialoadhesin family of the Ig superfamily are capable of mediating protein-sialic acid binding interactions. To date, only a small number of proteins have been assigned to the sialoadhesin family including sialoadhesin, CD33, CD22, the myelin-associated glycoprotein (MAG), and the Schwann cell myelin protein (SMP). Each of these proteins is expressed in a restricted subset of cell types. For example, CD22 and CD33 are expressed exclusively by B-lymphocytes and cells of the myelomonocytic lineage, respectively.

Similarly, galectins are a family of the lectin superfamily of carbohydrate-binding proteins which have a high affinity for b-galactoside sugars. Although a large number of glycoproteins containing b-galactoside sugars are produced by the cell, only a few will bind to known galectins *in vitro*. Such apparent binding specificity suggests a highly specific functional role for the galectins. Galectin 1 (conventionally termed *LGALS1* for lectin, galactoside-binding, soluble -1) is thought to specifically bind laminin, a highly polylactosaminated cellular glycoprotein, as well as the highly polylactosaminated lysosome-associated membrane proteins (LAMPs). Galectin 1 has also been shown to bind specifically to a lactosamine-containing glycolipid found on olfactory neurons and to integrin  $a_7b_1$  on skeletal muscle cells. Galectin 3 has also been observed to bind specifically to laminin, immunoglobulin E and its receptor, and bacterial lipopolysaccharides.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin 1 appears to inhibit cell adhesion of skeletal muscle presumably by galectin 1-mediated disruption of laminin-integrin  $a_7b_1$  interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined

5

10

15

20

25

30

presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively.

A relatively new family of cell-surface proteins has been identified and termed the Ly6 superfamily. The members of this family include murine and human SCA-2, rat Ly-6 (also termed ThB), human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)], and E48 antigen. The determination of an initial functional role for SCA-2 may lie in an analysis of its expression profile with regard to the complex process of hematopoiesis. SCA-2 is highly expressed in early thymic precusor cells. In turn, progeny of the intrathymic precusor population continue to express SCA-2, but only until the point of transition occurs from blast cell to small cell. Further experimental evidence demonstrates that mature thymocytes and peripheral T-cells do not express detectable levels of SCA-2, whereas mature, peripheral B-cells do continue to express SCA-2. As a result, it seems very likely that SCA-2 plays an important role in thymocyte maturation and differentiation. A plausible explanation for this functional hypothesis is that SCA-2 may act as a receptor for a unknown cytokine which regulates thymocyte maturation and differentiation.

In addition, CD59 is a recently identified integral membrane protein which appears to be involved in the regulation of complement. Recent studies show that the CD59 antigen may prevent damage from complement C5b-9 and protect astrocytes during inflammatory and infectious disorders of the nervous system. Expression of recombinant human CD59 on porcine donor organs have been shown to prevent complement-mediated lysis and activation of endothelial cells that leads to hyperacute rejection. Recently, researchers at Alexion Pharmaceuticals (New Haven, CT) reported on the production of transgenic pigs which expressed human CD59. In these animals, xenogeneic organs were resistant to hyperacute rejection. (Fodor, et al., "Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection," Proc. Natl. Acad. Sci., 91:1153-11157 (1994).) The same company also reported that expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria (PNH) B-

5

10

15

20

25

30

WO 98/31799 PCT/US98/00959

cells confers resistance to human complement. (Rother et al., "Expression of recombinant transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B-cells confers resistance to human complement," Blood, 84:2604-2611 (1994).) PNH is an acquired hematopoietic disorder characterized by complement-mediated hemolytic anemia, pancytopenia, and venous thrombosis. It is thought that retroviral gene therapy with this molecule could provide a treatment for PNH patients.

A final Ly6 superfamily member, the E48 antigen, is involved in intercellular adhesion between keratinocyte cells of the squamous epithelium. Such keratinocytes are attached to adjoining cells by large numbers of desmosomes, which are thought to play a role in the transition of transformed keratinocytes to metastatic tumor cells. Treatment with a monoclonal antibody raised against the E48 antigen has been successful in the eradication of residual, postoperative squamous cell carcinoma cells of the upper aerodigestive tract in several *in vivo* models and, to some degree, in humans. (van Dongen, et al., "Progress in radioimmunotherapy of head and neck cancer," Oncol. Rep. 1:259-264 (1994).) The gene encoding the E48 antigen has been mapped to the q24-qter region of human chromosome 8. Interestingly, a number of human diseases have been mapped to this region of chromosome 8 including Langer-Giedion syndrome, brachio-otorhinolaryngeal syndrome, trichorhinolaryngeal syndrome, and epidermolysis bullosa simplex.

A further example of a receptor family includes the prohibitin receptors. The prohibitin gene product is expressed in a wide variety of tissues and has been implicated as a component of a number of anti-proliferative mechanisms. The prohibitin gene encodes a 30 kD postsynthetically modified polypeptide located primarily in the mitochondria, but also may be associated with the IgM receptor on the B-cell plasma membrane. The protein functionally inhibits DNA synthesis and entry into S phase of the cell cycle by an unknown mechanism. Interestingly, although the prohibitin gene product is hypothesized to be involved in the maintenance of senescence and the prevention of cancer, one study found that, although somatic mutations in the prohibitin gene were present in a small number of breast cancers, no mutations were identified in any other breast, ovary, liver, and lung cancers examined. (Sato et al., Genomics 17:762-764 (1993).) However, the prohibitin gene has been mapped to human chromosome 17q12-21, the same region thought to contain the gene involved in sporadic breast cancer. Furthermore, DNA sequence analysis of the prohibitin gene identified somatic mutation in 4 of 23 cases of sporadic breast cancer examined. Thus, prohibitin family members may be involved in the development of cancer.

Moreover, the EGFR family of plasma membrane proteins are an integral component of normal cellular proliferation and in the pathogenesis of the cancerous state. The family is relatively small and includes the EGFR, c-erbB-2, c-erbB-3, and others. Various cancers are correlated with aberrant expression of one or more of these

5

10

15

20

25

**30** 

5

10

15

20

25

30

35

genes. A number of ligands have been identified which bind to the EGFR-like receptors listed above including TGF-a, heparin-binding EGF, amphiregulin, criptoregulin, heregulin, and others. A large fraction of adenocarcinomas examined to date, especially those of the breast, colon, and pancreas, are typified by the amplification or overexpression of the c-erbB-2 gene. EGF, or an analogous ligand, initiates the cellular growth factor response by binding to the EGFR, or EGFR-related, receptor. Following the binding event, the receptor molecule dimerizes activating its intracellular tyrosine kinase domain. This event results in the phosphorylation of specific tyrosine residues near the carboxy terminus of the receptor. The diversity of signals able to be transduced through the relatively small number of EGFR-related receptor molecules is amplified considerably by the recent finding that EGFR-like receptor molecules can function when dimerized with other EGFR family members forming heterodimers.

Members of the EGFR-related family of integral membrane proteins have been implicated in the pathogenesis of a number of human disease-states. For example, a mutation in the EGFR itself appears to play an important role in the development of glioblastomas. (Sang et al., J. Neurosurg 82:841-846 (1995).) The EGFR gene is amplified or overexpressed in the majority of primary human glioblastomas. Although not conferring a distinct advantage on cell growth, an increase in EGFR expression was found to confer an increase in the ability of glioma cells to maintain anchorage-independent growth in soft agar especially in response to EGF and retinoic acid. Anchorage-independent growth *in vitro* correlates highly with tumorigenicity *in vivo*, therefore, it is likely that cells which express abnormally high levels of EGFR in human glioblastoma cells may be involved in the high potential for these cells to cause tumors *in vivo*.

Moreover, overexpression or amplification of c-erbB-2 has been reported to be involved in a high number adenocarcinomas, particularly of the breast, colon, and pancreas, and in a small proportion of ovarian carcinomas.

Thus, there is a clear need for identifying and exploiting novel members of the receptor families, such as those described above. Although structurally related, these receptors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such receptors may prove useful as therapeutics in an antitumor, diagnostic, or other capacity. Furthermore, receptors described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

5

10

15

20

25

30

35

#### SUMMARY OF THE INVENTION

In one aspect, the invention relates to receptor polypeptides and polynucleotides, as well as the methods for their production. Another aspect of the invention relates to methods for using such receptor polypeptides and polynucleotides. Such uses include the treatment of the specified diseases, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with receptor imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate receptor activity or levels.

# **DESCRIPTION OF THE INVENTION Definitions**

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Receptor" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:Y, or an allelic variant thereof.

"Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said receptor including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said receptor.

"Receptor gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO:X or allelic variants thereof and/or their complements.

"SEQ ID NO:X" comprises all or a substantial portion of the polynucleotide encoding each receptor of the invention. The value X for the nucleotide sequence is an integer specified in Table 1. This nucleotide sequence was translated into the receptor polypeptide identified in Table 1 as "SEQ ID NO:Y," where the value of Y for each receptor polypeptide is an integer defined in Table 1.

The invention further provides a composition of matter comprising a nucleic acid molecule which comprises a human cDNA clone identified by a cDNA Clone ID (Identifier) in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection ("ATCC") and given the ATCC Deposit Number shown in Table 1 for that cDNA clone. The ATCC is located at American Type Culture Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852, USA. The deposit has been made under the terms of the Budapest Treaty on the international recognition of the deposit of micro-organisms for purposes of patent procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that

WO 98/31799 PCT/US98/00959

required under 35 U.S.C. §112. The nucleotide sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can

5

10

15

20

25

30

10

15

20

25

30

35

occur anywhere in a polypeptide, including the peptide backbone, the amino acid sidechains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphotidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY Acad Sci (1992) 663:48-62.)

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a

naturally occurring such as an allelic variant, or it may be a variant that is not known to

10

15

20

25

30

35

J Molec Biol (1990) 215:403.)

occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" per se has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991.) While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans. (Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073.) Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., SIAM J Applied Math (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., et al., Nucleic Acids Research (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. et al.,

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO:X is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: X. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 ' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO: Y is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO:Y. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

20

25

10

5

### Polypeptides of the Invention

In one aspect, the present invention relates to receptor polypeptides (or receptor proteins). The receptor polypeptides include the polypeptide of SEQ ID NO:Y; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:Y; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% identity to SEQ ID NO:Y are highly preferred. Also included within receptor polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:Y over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO:Y. Furthermore, those with at least 97-99% are highly preferred. Preferably receptor polypeptides exhibit at least one biological activity of the receptor.

30

The receptor polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

35

Fragments of the receptor polypeptides are also included in the invention. A "fragment" is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned receptor polypeptides. As with receptor polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most

10

15

20

25

30

35

preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of receptor polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of receptor polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus.

Also preferred are fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. The "domains" of each receptor polypeptide are illustrated in the Figures. The Figures compare SEQ ID NO:Y to the closest know homologue. Identical amino acids shared between the two polypeptides are shaded, while conservative amino acid changes are boxed. By examining the regions or amino acids shaded and/or boxed, the skilled artisan can readily identify conserved domains between the two polypeptides. The amino acids sequences of SEQ ID NO:Y falling within these conserved domains are "fragments" and are specifically contemplated by the present invention. Especially preferred is the extracellular domains of a receptor of the invention. Soluble extracellular domains have antagonist activity mediated by competition with a receptor ligand.

Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain a biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala. Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

10

15

20

25

30

35

The receptor polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

## Polynucleotides of the Invention

Another aspect of the invention relates to receptor polynucleotides. Receptor polynucleotides include isolated polynucleotides which encode the receptor polypeptides and fragments, and polynucleotides closely related thereto. More specifically, a receptor polynucleotide of the invention includes a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:X encoding a receptor polypeptide of SEQ ID NO:Y, and polynucleotide having the particular sequence of SEQ ID NO:X.

Receptor polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under receptor polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:X, or contained in the cDNA insert in the plasmid deposited with ATCC, to hybridize under conditions useable for amplification or for use as a probe or marker. Moreover, the receptor polynucleotide includes a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC, and a nucleotide sequence comprising at least 15 contiguous nucleotides of such cDNA insert. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. The invention also provides polynucleotides which are complementary to all the above receptor polynucleotides.

The receptors of the invention are structurally related to other proteins of specified receptor families, as shown by the results in the Figures. The cDNA sequence of SEQ ID NO:X encodes a polypeptide as described in Table 1 as SEQ ID NO:Y. Because the receptor polypeptides contain domains similar in structure to other receptor family members, the receptors of the present invention are expected to have,

inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1

5

Clone ID Name	SEQ ID NO:X	SEQ ID NO:Y	ATCC Deposit No.	ATCC Deposit Date	Receptor Family	Homology
TD 44 CD 70	NO:A	NO: 1	209054	05/16/97	I a	Sialoadhesin
HMACR70	,	1.0	1		Ig	OB-1
1100001110	1	18	#####	01/21/98	T) (40)	
HTEDK48			209054	05/16/97	TM4SF	MRC-OX44
						PETA-3
1-1849 bp	2					
160-900 bp	3	19				
HTPED39			209054	05/16/97	TM4SF	NAG-2
HPWAE25	4	20	#####	1/21/98		TALLA-1
HTPEF86	5	21	209053	05/16/97	TM4SF	CD20
		_				B1 Antigen
HSBBF02	6	22	209054	05/16/97	TM4SF	TALLA-1
HLTAH80	7	23	97242	08/02/95	TM4SF	TALLA-1
			209054	05/16/97		
HTPBA27	8	24	97242	08/02/95	TM4SF	NAG-2
			209054	05/16/97		
HAIDQ59			209054	05/16/97	TM4SF	CD9
				ļ		Antigen
5' Sequence	9	25				
3' Sequence	10					
HHFEK40	11	26	209054	05/16/97	TM4SF	PETA-3
HGBGV89	12	27	209125	06/09/97	TM4SF	L6H
			209054	05/16/97	: -	
HUVBB80	13	28	209054	05/16/97	TM4SF	L6
HJACE54	14	29	209053	05/16/97	Lectin	Galectin-3
	-					Galectin-5
,	ļ					Galectin-8
HROAD63	15	30	209053	05/16/97	Ly6	E48 splice
		1				variant
HMWGS46	16	31	209053	05/16/97	Prohibitin	BAP-37
HNFGW06	17	32	209053	05/16/97	EGFR	EGFR

The novel full-length cDNA clone designated HMACR70 may be a member of the sialoadhesin family of the Ig superfamily of receptor-like molecules and a CD33 homologue. HMACR70 contains a 1497 nucleotide cDNA insert encoding a 315 amino acid ORF and was cloned from a GM-CSF-treated human macrophage cDNA library. The only additional cDNA libraries in the HGS database which include this clone are human eosinophils and possibly human gall bladder. A BLAST analysis of the amino acid sequence of HMACR70 demonstrates that this clone exhibits approximately 50% identity and 69% similarity over a 300 amino acids stretch of a gene termed human

10

10

15

20

25

30

35

differentiation antigen, and 38% identity and 62% similarity of the human myelin-associated glycoprotein precursor CD33 gene.

A more recent BLAST analysis confirms HMACR70's designation as a sialoadhesin family member. HMACR70 is homologous to two recently identified sialoadhesin family members, human OB binding protein (OB) 1 and 2. (See, Genbank Accession No. U71382; see Figure 1.) It is thought that OB-1 and OB-2 may bind leptin. Thus, HMACR70, as a sialoadhesin family member, may act to attenuate or even amplify intercellular routes of communication, including binding to leptin or modulating the activity of immune cells, such as macrophages. Clearly, any diseases affected by these processes could be treated by the polypeptide or fragment of HMACR70.

The full-length nucleotide sequences of ten novel human cDNA clones which potentially belong to the TM4SF superfamily are disclosed in the table above and will be addressed sequentially.

The cDNA clone HTEDK48 contains a 1849 nucleotide cDNA insert encoding a 245 amino acid ORF that was cloned from a human testes cDNA library. The coding sequence of HTEDK48 (SEQ ID NO: 3) may be fused to other human proteins, such as 3-hydroxyacyl-CoA dehydrogenase. BLAST analysis of the amino acid sequence of HTEDK48 demonstrates that this clone exhibits approximately 30% identity and 51% similarity over a 245 amino acid stretch of the CD82 molecule. Recent studies have shown that CD82 can associate with CD4 or CD8 and deliver costimulatory signals for the TCR/CD3 pathway. CD82 has also been found to be involved in syncytium formation in HTLV-I-infected T-cells. And finally, in a recently published study in which the expression of the CD82 gene by tumors of the lung was examined retrospectively, it was reported that CD82 may be linked to the suppression of tumor metastasis of prostate cancer. The study also reported that decreased CD82 expression may be involved in malignant progression of such cancers. Thus, HTEDK48 may also be involved in the development of cancer.

A more recent BLAST analysis shows that HTEDK48 is homologous the rat leukocyte antigen, MRC OX-44, and the platelet endothelial tetraspan antigen -3 (PETA-3). (See Figure 2X.) MRC OX-44, a member of a new family of cell surface proteins, appears to be involved in growth regulation. (See, Bellacosa, A., et al., "The Rat Leukocyte antigen MRC OX-44 is a Member of a New Family of Cell Surface Proteins which Appear to be Involved in Growth Regulation," Mol. Cell. Bio. 11: 2864-2872 (1991).) Similarly, PETA-3 has been located to platelet endothelial cells, and an anti-PETA-3 antigen monoclonal antibody can stimulate platelet aggregation and mediator release. (See, Fitter, S., "Molecular Cloning of cDNA Encoding a Novel Platelet-Endothelial Cell Tetra-Span Antigen, PETA-3," Blood, 86(4):1348-1355 (1995).) Thus, HTEDK48 may function similar to MRC OX-44 or PETA-3 to affect

growth of blood cells. Administering polypeptides or fragments of HTEDK48 may be an effective treatment of blood disorders.

PCT/US98/00959

The cDNA clone **HPWAE25** contains a 1288 nucleotide cDNA insert encoding a 273 amino acid ORF that was cloned from a human pancreas tumor cDNA library, while clone **HTPED39** represents a truncated cDNA sequence. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including keratinocytes, ulcerative colitis, striatum depression, lymph node breast cancer, ovarian cancer, stage B2 prostate cancer, kidney medulla, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human cell lines including U937, MM96, WM115, and MDAMB231. A BLAST analysis of the amino acid sequence of HTPED39 demonstrates that this clone exhibits approximately 35% identity and 50% similarity over the entire length of the CD37 molecule. The CD37 antigen is expressed on B cells and on a subpopulation of T cells, but not on pre-B or plasma cells. It has been reported that CD37 expression is downregulated in conjunction with B-cell activation, suggesting that CD37 may be involved in the processes which dictate the activation state of the B-cell.

Moreover, HPWAE25 is also homologous to recently identified TM4SF members, NAG-2 and TALLA-1. (See Figure 3.) NAG-2 is thought to complex with integrins and other TM4SF proteins, while TALLA-1 is a highly specific marker of T-cell acute lymphoblastic leukemia and neuroblastoma. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that Complexs with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997); Takagi, S., "Identification of a Higly Specific Surface Marker of T-cell Acute Lymphoblastic Leukemia and Neuroblastoma as a New Member of the Transmembrane 4 Superfamily," Int. J. Cancer 61(5):706-715 (1995).) Thus, HPWAE25 may be involved the development of cancer, particularly leukemia, lymphoma, and neuroblastoma. HPWAE25 may be used as an effective treatment of these cancers, as well as a diagnostic marker.

A subfamily of TM4SF receptors include CD20 proteins. A CD20-like cDNA clone was obtained from a human pancreas tumor cDNA library and contains a 1236 nucleotide insert which encodes a 250 amino acid ORF. A BLAST analysis of the deduced amino acid sequence of HTPEF86 exhibits approximately 41% identity and 61% similarity to the CD20 gene, also known as B1 antigen. (See Figure 4.) Expression of this gene is detected in only two additional HGS human cDNA libraries; amygdala depression and 9 week early stage human. Although the precise functional role of CD20 has yet to be determined, it is clear that CD20 plays a key role in the regulation of B-cell activation. Based primarily on sequence identity, the novel CD20-like molecule presented herein may also be involved in cell cycle activation. Potential therapeutic and/or diagnostic applications for HTPEF86 may include such clinical

. >

5

10

15

20

25

30

10

15

20

25

30

35

presentations as juvenile rheumatoid arthritis, Graves' Disease, and a number of B-cell lymphomas or other lymphoid tumors.

The clone **HSBBF02** contains a 1115 nucleotide cDNA insert encoding a 245 amino acid ORF and was cloned from an HSC 172 cell line cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including brain amygdala depression, endothelial cells, fetal liver and heart, osteoblasts, testes, and others. A BLAST analysis of the amino acid sequence of HSBBF02 demonstrates that this clone exhibits approximately 64% identity and 80% similarity with the A15 molecule over a 131 amino acid stretch (A15 is composed of 244 amino acids). A more recent BLAST search shows that HSBBF02 is similar to the TALLA-1 protein and may in fact be a closely related family member. (See Figure 5.)

In addition, a second cDNA clone, designated **HLTAH80**, exhibits sequence similarity to the A15 molecule and TALLA-1. (See Figure 6.) This clone contains a 1662 nucleotide cDNA insert encoding a 253 amino acid ORF and was cloned from a human T-cell lymphoma cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including B-cell lymphoma, corpus collosum, endometrial tumor, osteosarcoma, testes, and others. Northern blot analysis of HLTAH80 also shows expression in a variety of human tissues including spleen, lymph node, thymus, PBLs, heart, and a particularly strong signal in skeletal muscle and pancreas. A BLAST analysis of the amino acid sequence of HLTAH80 demonstrates that this clone exhibits approximately 35% identity and 55% similarity over the entire length of the A15 molecule.

Since expression of A15 drops to undetectable levels when comparing immature T-cells to peripheral blood lymphocytes, it is thought that A15 may play a role in the development of T-cells. Furthermore, the MXS1(CCG-B7) gene which codes for A15 contains a number of triplet nucleotide repeats which have been associated with neuropsychiatric diseases such as Huntington's chorea, fragile X syndrome, and myotonic dystrophy. In addition, A15 appears to be expressed exclusively on T-cell acute lymphoblastic leukemia cell lines, including several derived from adult T-cell leukemia and those established by immortalization with human T-cell leukemia virus type 1 or Herpesvirus saimiri. Thus, clones HLTAH80 and/or HSBBF02 may also be involved in diseases caused by the expansion of repeats or chromosomal instability.

The cDNA clone HTPBA27 contains a 1345 nucleotide cDNA insert encoding a 238 amino acid ORF and was cloned from a human tumor pancreas cDNA library. This clone also appears in a number of other cDNA libraries constructed from a variety of human cell and tissue types including cerebellum, breast lymph node, osteosarcoma, adult testes, RS4;11 bone marrow cell line, microvascular endothelial cells, and others. A BLAST analysis of the amino acid sequence of HTPBA27 demonstrates that this

WO 98/31799 PCT/US98/00959

clone exhibits approximately 40% identity and 64% similarity with a glycoprotein termed CD53 over its entire length. CD53 is thought to be involved in thymopoiesis, since rat CD53 can be detected on immature CD4-8-thymocytes and the functionally mature single-positive subset, but cannot be detected on the intermediate CD4+8+ thymocytic subset of cells. The CD53 molecule has also been implicated as a component of signal transduction pathways in B cells, monocytes and granulocytes, rat macrophages; NK, and T cells. Moreover, as illustrated in Figure 7, HTPBA27 was recently confirmed as a TM4SF receptor. (See, Tachibana, I., et al., "NAG-2, A Novel Transmembrane-4 Superfamily (TM4SF) Protein that with Integrins and Other TM4SF Proteins," J. Biol. Chem., 272:29181-29189 (1997).) Calling the HTPBA27 polypeptide NAG-2, this group confirmed HTPBA27's status as a TM4SF receptor by showing that NAG-2 complexes with integrin and other TM4SF receptors. diseases caused by the failure of HTPBA27 to complex with integrin and other TM4SF receptors can be treated by administering HTPBA27. HTPBA27 can also be used to diagnose these diseases.

The cDNA clone **HAIDQ59** contains cDNA insert encoding a 221 amino acid ORF that was cloned from a human epithelial cell induced with TNFa and INF cDNA library. The 5' end of HAIDQ59 is represented by the SEQ ID NO: 9, while the 3' end is represented by SEQ ID NO: 10. This clone appears in only two additional cDNA libraries in the HGS database. These two libraries were constructed from the human Jurkat T-cell line and human microvascular endothelial cells. A BLAST analysis of the amino acid sequence of HAIDQ59 demonstrates that this clone exhibits approximately 53% identity and 69% similarity over 226 amino acids of the CD9 TM4SF molecule. (See Figure 8.) It has been demonstrated that the CD9 molecule is involved in signal transduction pathways in platelets, as well as in cell adhesion in both platelets and pre-B-cell lines. Intriguingly, a monoclonal antibody (vpg15), which recognizes the feline homologue of CD9, has been shown to block infection by feline immunodeficiency virus (FIV). Furthermore, a recent study shows that cells expressing high levels of CD9 exhibited suppressed cell motility. Thus, HAIDQ59 may also be involved in signal transduction of blood cells.

The cDNA clone **HHFEK40** contains a 936 nucleotide cDNA insert encoding a 252 amino acid ORF and was cloned from a human fetal heart cDNA library. This clone appears once in the human fetal heart cDNA library and possibly in a hemangiopericytoma cDNA library. A BLAST analysis of the amino acid sequence of HHFEK40 demonstrated that this clone exhibits approximately 60% identity and 75% similarity over the entire length of a molecule designated PETA-3. (See Figure 9.) PETA-3 was originally identified as a novel human platelet surface glycoprotein termed gp27. Although PETA-3 is present in low abundance on the platelet surface, an anti-PETA-3 monoclonal antibody can stimulate platelet aggregation and mediator release.

5

10

15

20

25

30

10

15

20

25

30

35

Thus, HHFEK40 may function similar to PETA-3 to affect growth of blood cells. Administering polypeptides or fragments of HHFEK40 may be an effective treatment of blood disorders.

The cDNA clone HGBGV89 contains a 738 nucleotide cDNA insert encoding a 197 amino acid ORF and was cloned from a human gall bladder cDNA library. The only two additional appearances of this clone in the HGS database are in a normalized fetal liver cDNA library and in a fetal liver/spleen cDNA library. The cDNA clone HUVBB80 contains a 1071 nucleotide cDNA insert encoding a 201 amino acid ORF and was cloned from a human umbilical vein cDNA library. This clone appears in several additional cDNA libraries in the HGS database including prostate BPH, thyroid, and fetal liver/spleen. BLAST analyses of the amino acid sequences of HGBGV89 and HUVBB80 demonstrate that these clones exhibit approximately 49% identity and 65% similarity and 47% identity and 68% similarity, respectively, over the entire length of a molecule designated L6 surface protein or human tumor-associated antigen L6. (See Figures 10 & 11.) Moreover, another group has confirmed the TM4SF receptor homology of HGBGV89 by describing the protein as a putative transmembrane protein L6H. (See Genbank Accession No 2587054; see Figure 10.) The L6 cell surface antigen is highly expressed on lung, breast, colon, and ovarian carcinomas. Promising results of phase 1 clinical studies have been reported with an anti-L6 monoclonal antibody, or its humanized counterpart, suggesting that the L6 antigen may be an attractive target for monoclonal antibody-based cancer therapy.

In summary, there is a clear need for identifying and exploiting novel members of the TM4SF superfamily such as those described herein. Although structurally related, these factors will likely possess diverse and multifaceted functions in a variety of cell and tissue types. Receptor type molecules, such as the novel potential members of the TM4SF superfamily detailed here, should prove useful in target based screens for small molecules and other such pharmacologically valuable factors. Monoclonal antibodies raised against such factors may prove useful as therapeutics in an anti-tumor, diagnostic, or other capacity. Furthermore, factors such as the nine novel TM4SF superfamily-like molecules described here may prove useful in an active or passive immunotherapeutical role in patients with cancer or other immunocompromised disease states.

Besides TM4SF receptors, receptors from other families are also described. For example, clone HJACE54, also called galectin 11, exhibits significant sequence identity to the rat galectin 5, the chicken galectin 3 gene, and the human galectin 8 genes. (See Figure 12.) The galectin 11 cDNA clone contains an 865 nucleotide insert which encodes a 133 amino acid ORF. The clone was obtained from a Jurkat T-cell G1 phase cDNA library. A BLAST analysis of the deduced amino acid sequence of HJACE54 demonstrates approximately 35% identity and 57% similarity to the amino

WO 98/31799 PCT/US98/00959

acid sequence of the rat galectin 5 gene. Expression of galectin 11 is quite limited in the HGS database. In fact, the only two additional ESTs in the HGS database which contain the HJACE54 sequence were found in human neutrophil and human infant adrenal gland cDNA libraries. Northern blot analyses have not been performed to examine expression patterns of the galectin 11 gene.

Various galectins have been shown to function in the mechanisms of intercellular communication. For example, depending on cell type, galectin 1 has been observed to modulate cell adhesion either positively or negatively. More specifically, galectin I appears to inhibit cell adhesion of skeletal muscle presumably by galectin Imediated disruption of laminin-integrin a₇b₁ interactions. Alternatively, galectin 1 appears to promote cell adhesion in several non-skeletal muscle cell types examined presumably by a glycoconjugate cross-linking mechanism. Galectin 3 has also been observed to function in modulating cell-adhesion, as well as in the activation of certain immune cells by cross-linking IgE and IgE receptors. In addition, galectins have been observed to be involved in the regulation of immune cell activity, as well as in such diverse processes as cell adhesion, proliferation, inflammation, autoimmunity, and metastasis of tumor cells. Furthermore, a galectin-like antigen designated HOM-HD-21 was recently found to be highly expressed in a Hodgkin's Disease cDNA library. Very recently, a novel galectin, termed PCTA-1, was identified as a specific cell surface marker on human prostate cancer cell lines and patient-derived carcinomas. Galectins have also been found to function intracellularly as a component of ribonucleoprotein complexes. Finally, galectins 1 and 3 have each been found to modulate T-cell growth and apoptosis by interaction with CD45 and possibly Bcl2, respectively. As a result, the discovery of a novel galectin, such as that encoded by HJACE54, is likely to be a valuable asset both diagnostically and therapeutically.

Additionally, a full-length nucleotide sequence of a novel human cDNA clone which encodes an apparent splice variant of the previously described human E48 antigen has recently been determined. (See Figure 13.) Clone HROAD63 contains a 441 nucleotide cDNA which encodes a 70 amino acid polypeptide. This novel clone exhibits significant sequence identity to several members of a relatively new family of cell-surface proteins termed the Ly6 superfamily. These members include murine and human SCA-2, rat Ly-6 (also termed ThB), and human CD59 [also known as protectin or membrane attack complex inhibition factor (MACIF)]. The novel E48 splice variant was obtained from the HGS human stomach cDNA library. The clone is present in only a limited number of other HGS cDNA libraries including kidney cancer, keratinocyte, and tongue. An alignment of the nucleotide sequences of the human E48 and HROAD63 cDNAs demonstrates that the initial 168 and 178 nucleotides of E48 and HROAD63, respectively, are identical, with the exception of an additional 10 nucleotides of sequence at the extreme 5' end of the HROAD63 sequence. The

5

10

15

20

25

30

10

15

20

25

30

35

sequence of the two clones is also identical for an additional 229 nucleotides including the 3' end of the coding sequences and the entire 3' untranslated regions. The only divergence of nucleotide sequence in this region of the clones is the deletion of a single thymidine residue in the 3' UTR of the E48 cDNA. The major difference between the two nucleotide sequences is a 329 nucleotide deletion from the HROAD63 sequence. This deletion causes a shift in the HROAD63 reading frame and encompasses the translational stop signal used in the E48 clone. As a result, the carboxy terminal sequence of HROAD63 is radically altered with regard to that of E48 (as illustrated in Figure 13 by the obvious differences between amino acids 56-128 of E48 and 56-70 of HROAD63 in the amino acid alignment). The clinical presentation of disorders, including abnormal skin and hair phenotypes, may be attributed, at least in part, to a non-functional Ly6 superfamily member such as E48 or HROAD63. HROAD63 may also be involved in blood disorders, as seen with its homologues SCA-2 and CD59.

A novel prohibitin cDNA clone presented herein was originally identified in a human bone marrow cell line (RS4;11) cDNA library. The clone contains a 1066 nucleotide insert which encodes a 299 amino acid polypeptide. BLAST and BestFit analyses of the predicted amino acid sequence of HMWGS46 demonstrate a highly significant sequence identity to a murine protein termed IgM B-cell receptor associated protein (BAP)-37 (Genbank accession number X78683). The HMWGS46 amino acid sequence exhibits nearly perfect identity and similarity over the entire length of the murine BAP-37 sequence. (See Figure 14.) In addition, the full-length nucleotide sequences of HMWGS46 and BAP-37 exhibit at least 87% identical. The HMWGS46 clone also exhibits approximately 49% sequence identity and 85% sequence similarity to a human gene designated prohibitin. Finally, the HMWGS46 cDNA appears in a substantial number of HGS human cDNA libraries in addition to the bone marrow cell line cDNA library from which it was cloned. Some of the cDNA libraries in which this clone appears include keratinocytes, induced endothelial cells, activated neutrophils, synovial sarcoma, colon carcinoma cell line, Jurkat cell line membrane bound polysomes, epileptic frontal cortex, primary dendritic cells, and a number of others. The novel gene related to prohibitin and BAP-37 may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event. Thus, although the precise functional role of the prohibitin family members are less than clear, it is quite likely that such homologues are involved in such complex processes as development, senescence, and tumor suppression. Therefore a novel gene, such as HMWGS46, may prove quite useful as a diagnostic for tumorigenesis, as well as a target for therapeutic intervention of such an event.

A human cDNA clone encoding a novel epidermal growth factor receptor (EGFR)-like molecule is also disclosed. The novel EGFR-like cDNA clone presented herein was originally identified in an activated human neutrophil cDNA library. The

WO 98/31799

clone contains a 704 nucleotide insert which encodes a 168 amino acid polypeptide. A BLAST analysis of the predicted amino acid sequence of **HNFGW06** demonstrates that this novel clone exhibits approximately 85% identity and 90% similarity to a protein designated epidermal growth factor receptor-related protein [Homo sapiens]. (See Figure 15.) The expression profile of the HNFGW06 clone in the HGS database indicates the existence of a fairly highly restricted expression pattern. In addition to the activated neutrophil library from which this clone was obtained, it also appears in the following HGS human cDNA libraries: synovial sarcoma, smooth muscle, placenta, and possibly primary dendritic cells.

PCT/US98/00959

The novel EGFR-like cDNA clone HNFGW06 may lead to a number of exciting possibilities for therapeutic and/or diagnostic treatments or reagents. For example, HNFGW06 may be involved in the onset of human breast cancers as well. In addition, due to the fact that TGF-a acts through binding to the EGFR, it is possible that HNFGW06 may also play a role in a variety of gastric processes including regulation of acid secretion, regulation of mucous cell growth, and protection against ethanol- and aspirin-induced injury to gastric tissues.

### GENERATING POLYNUCLEOTIDES

Polynucleotides of the present invention encoding a receptor may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells specified in Table I using the expressed sequence tag (EST) analysis (Adams, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et al., Nature (1995) 377 Supp:3-174.) Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding a receptor polypeptide of SEQ ID NO:Y may be identical to the polynucleotide encoding SEQ ID NO:Y, or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:Y.

When the polynucleotides of the invention are used for the recombinant production of a receptor polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in

5

10

15

20

25

30

10

15

20

25

30

35

Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding receptor variants comprising the amino acid sequence of receptor polypeptide of Table 1 (SEQ ID NO:Y) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:X or a fragment thereof, or to the cDNA insert in the plasmid deposited at the ATCC, or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding the receptor and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs) that have a high sequence similarity to the receptor gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding the receptor polypeptide, including homologs and orthologs from other species, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO:X or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

WO 98/31799 PCT/US98/00959

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

# 5 Vectors, Host Cells, Expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., BASIC METHODS IN MOLECULAR BIOLOGY (1986) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., MOLECULAR CLONING, A LABORATORY MANUAL (supra).

10

15

20

25

30

10

15

25

30

35

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the receptor polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the receptor polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Receptor polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

### 20 Diagnostic Assays

This invention also relates to the use of receptor polynucleotides or polypeptides for use as diagnostic reagents. Detection of a mutated form of the receptor gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from underexpression, over-expression or altered expression of the receptor. Individuals carrying mutations in the receptor gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled receptor nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. (See, e.g., Myers et al., Science (1985) 230:1242.) Sequence changes at specific locations may also be revealed by

10

15

20

25

35

nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. (See Cotton et al., Proc Natl Acad Sci USA (1985) 85: 4397-4401.) In another embodiment, an array of oligonucleotides probes comprising receptor nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996).)

The diagnostic assays offer a process for diagnosing or determining a susceptibility to specific diseases through detection of mutation in the receptor gene by the methods described.

In addition, specific diseases can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of receptor polypeptide or receptor mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease which comprises:

- (a) a receptor polynucleotide, preferably the nucleotide sequence of SEQ ID NO:X, or a fragment thereof;
  - (b) a nucleotide sequence complementary to that of (a);
- (c) a receptor polypeptide, preferably the polypeptide of SEQ ID NO:Y, or a fragment thereof; or
- (d) an antibody to a receptor polypeptide, preferably to the polypeptide of SEQ ID NO: Y.
- It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

# Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the

sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10

15

20

5

#### Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the receptor polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the receptor polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole et al., MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

30

25

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against receptor polypeptides may also be employed to treat diseases.

### Vaccines

35

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a receptor polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from a disease. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which

comprises, delivering a receptor polypeptide via a vector directing expression of the receptor polynucleotide in vivo in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a receptor polypeptide wherein the composition comprises a receptor polypeptide or receptor gene. The vaccine formulation may further comprise a suitable carrier. Since a receptor polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation instonic with the blood of the recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

# Screening Assays

5

10

15

20

25

30

35

The receptor polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).

The receptor polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate the receptor on the one hand and which can inhibit the function of the receptor on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions and diseases. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions and diseases.

In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof.

10

15

20

25

30

Such cells include cells from mammals, yeast, Drosophila or E. coli. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a receptor polypeptide to form a mixture, measuring receptor activity in the mixture, and comparing the receptor activity of the mixture to a standard.

The receptor cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of receptor mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of receptor protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of the receptor (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

Examples of potential receptor antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the receptor, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for receptor polypeptides; or compounds which decrease or enhance the production of receptor, which comprises:

- (a) a receptor polypeptide, preferably that of SEQ ID NO:Y;
- (b) a recombinant cell expressing a receptor polypeptide, preferably that of SEQ35 ID NO:Y;
  - (c) a cell membrane expressing a receptor polypeptide; preferably that of SEQ ID NO: Y; or
    - (d) antibody to a receptor polypeptide, preferably that of SEQ ID NO: Y.

SUBSTITUTE SHEET (RULE 26)

10

15

20

25

30

35

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

# Prophylactic and Therapeutic Methods

This invention provides methods of treating an abnormal conditions related to both an excess of and insufficient amounts of receptor activity.

If the activity of the receptor is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking the binding of ligands to the receptor or by inhibiting a second signal, and thereby alleviating the abnormal condition.

In another approach, soluble forms of the receptor polypeptides still capable of binding the ligand in competition with endogenous receptor may be administered. Typical embodiments of such competitors comprise fragments of the receptor polypeptide.

In still another approach, expression of the gene encoding endogenous receptor can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. (See, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Alternatively, oligonucleotides which form triple helices with the gene can be supplied. (See, for example, Lee et al., Nucleic Acids Res (1979) 6:3073; Cooney et al., Science (1988) 241:456; Dervan et al., Science (1991) 251:1360.) These oligomers can be administered per se or the relevant oligomers can be expressed in vivo.

For treating abnormal conditions related to an under-expression of the receptor and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates the receptor, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition.

Alternatively, gene therapy may be employed to effect the endogenous production of the receptor by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells in vivo and expression of the polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and

10

15

20

25

30

35

other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

# Formulation and Administration

Peptides, such as the soluble form of receptor polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds. Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of  $0.1\text{-}100\,\mu\text{g/kg}$  of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide ex vivo, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

**SUBSTITUTE SHEET (RULE 26)** 

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

SUBSTITUTE SHEET (RULE 26)

																							H	M	A	Z	I	2		
CAG	TTO	ССТ	GAG.	AGA	AGA	ACC	CTG.	AGG	AAÇ	AGA	CGT	TCC	CTC	GCGC	SCC	TG	GCA	CCT	CCA	ACC	CA	GATA	TGC	TG	TG	CTG	TGC	TGC	TG	90
GTC	AA	GGA	CTC	TCI	TCT	TGG	GAC	TCC	TTG	тст	GC A	AGG	GAG	CGC	CGGC	SAC	CGT	GGA	GGT	TGG	GT	CTA	TACO	ACC	SAC	GAC	GACG	SACG		50
					,																		М	L	L	L	L	L	L	
ccc	TG	стс	TGG	GG	SAGG	GAG	AGG	GTG	GAA	GGA	CAG	AAG	AGT	AAC	CGG	AAG	GAT	TAC	TCG	CTG	ACG	ATG	CAGA	GT	rcc	GTG.	ACCO	TGC	AA.	180
GGG	AC	GAG.	ACC	cco	CTCC	CTC	TCC	CAC	CTT	CCT	GTC	TTC	TCA	TTG	GCC.	TTC	CTA	ATG	AGC	GAC.	TGC	TAC	STCI	CA	AGG	CAC	TGGC	ACC	113	
Р	L	L	W	G	R	Ε	R	٧	Ε	G	0	K	s	N	R	K	O	Y	S	L	Т	М	O	S	S	٧	T	٧	a	
SAGO	GC.	ATG	TGT	GTO	CCAT	GTG	CGC	TGC	тсс	TTC	TCC	TAC	CCA	GTG	GAC	AGC	CAG	ACT	GAC	тст	GAC	CCA	ĢTTO	AT	GGC	TAC	TGGI	TTCC	ee	070
CTCC	CG	TAC	ACA	CA	GGTA	CAC	GCG	ACG	AGG	AAG	AGG	ATG	GGT	CAC	CTG	TCG	GTC	TGA	CTG	AGA	CTG	GGT	CAA	ATA	CCG	ATG.	ACCA	AAGO	scc	270
Ε	G	М	С	٧	н	٧	R	С	s	F	s	Y	Р	٧	D	S	a	T	D	S	0	Ρ	٧	н	G	Y	W	F	R	
					AAGC											_		<del></del>											_	360
CGT	ccc	TTA	CTA	TA	TTCC	ACC	TTC	CGA	GGT	CAC	CGG	TGT	TTG	TTG	GGT	CGA	ACC	CGT	CAC	GTC	CTC	CTT	TGA	GCC	CTG	GCT	AAG	STGO	GAG	
Α	G	N	D	ì	S	W	κ	A	Р	٧	Α	T	N	N	Р	A	W	A	٧	Q	Ε	Ε	T	R	D	R	F	Н	L	
CTT	GGG	GAC	CCA	CA	GACC	:AAA	AAT	TGC	ACC	CTG	AGC	ATC	AGA	GAT	GCC	AGA	ATG	ĄGT	GAŢ	GCG	GGG	AGA	TAC	TTC	TTT	CGT	ATG	GAG	AAA	450
GAA	CCC	CTG	GGT	GT	CTGC	111	TTA	ACG	TGG	GAC	TCG	TAG	TCT	CTA	CGG	TCT	TAC	TCA	CTA	CGC	CCC	TCT	ATG.	AAG	AAA	GĊA	TAC	CTC.	TTT	450
L	G	D	Р	a	τ	K	N	С	T	L	s	i	R	D	Α	R	M	s	D	Α	G	R	Y	F	F	R	М	Ε	K	
GGA.	AAT	ATA	AAA	TG	GAA1	ΓŢΑΊ	AAA	TAI	GAC	CAG	стс	TCT	GTG	AAC	GTG	AÇA	TAC	CCT	CCT	CAG	AAC	TTG	ACT	GTG	ACT	GTC	TTC	CAA	GGA	540
ССТ	TTA	TAT	TTT	AC	CTTA	ATA	TTT	ATA	CTO	GTC	GAG	AGA	CAC	TTG	CAC	TĠT	ATG	GGA	GGA	GTC	TTG	AAC	TGA	CAC	TGA	CAG	AAG	GTT	CCT	
G	N	1	ĸ	W	N	Y	κ	Y	D	Q	L	s	٧	N	٧	T	Y	P	Р	a	N	Ł	T	٧	T	V	F	a	G	
GAA	GGC	ACA	GCA	TC	CAC	AGCT	CTG	GGG	SAAC	AGC	TCA	TCT	стт	TCA	GTC	CTA	GAG	GGC	CAG	тст	CTO	CGC	TTG	GTC	TGT	GCT	GTT	GAC.	AGC	630
CTT	CCG	TGT	CGT	AG	GTG	rcg/	GAC	ccc	TTC	TCG	AGT	AGA	GAA	AGT	CAG	GA1	СТС	ccg	GTC	AGA	GAC	GCG	AAC	CAG	ACA	CGA	CAA	CTG	TCG	
ε	G	T,	A	s	Т	A	L	G	N	s	s	s	L	S	٧	L	ε	G	a	s	L	R	L	٧	С	A	٧	0	S	
AAT	ccc	:001	GCC	AG	GCT	GAG	TGG	SAC	TG	SAGG	AGT	сто	ACC	CTG	TAC	cçc	TCA	CAG	ccç	TCA	AAC	CCT	CTG	GTA	сто	GAG	CTG	CAA	GTG	720
TTA	GGG	GGA	CGG	TC	CGA	CTC	SACC	TG	SAC	TCC	TCA	GAC	TGC	GAC	ATG	GGG	SAGI	GTC	GGG	SAGT	TTO	GGA	GAC	CAT	GAC	стс	GAC	GTT	CAC	,20
N	Р	Р	Α	R	L	s	W	Ţ	w	R	s	L	T	L	Y	Р	s	Q	P	s	N	Р	L	٧	L	Ε	L	a	٧	
CAC	CTG	SGGG	GAT	ΓGΑ	AGG	GGĄ	ATTO	CACI	CTG	TCGA	NGC 1	CAC	SAAG	TCT	CTG	GĢI	rtco	CCAG	CAC	GTT	TC	CTG	AAC	стс	TC	CŢC	CAA	CAG	GAG	810
GTG	GAC	ccc	CTA	ACT	TCC	CCT	TAAC	STG	GAC	AGCT	CGA	GTO	TTO	SAGA	GAC	CC/	AAGO	GTO	GTO	CAA	AGO	GAC	TTG	GAG	AG	GAC	GTT	GTC		
н		c	n	F	G	F	F	т	۲	R	А	a	N	s	L	G	s	Q	н	٧	s	L	N	Ł	s	L	a	a	Ε	

HMAC RTC Y T G K M K P V S G V L L G A V G G T G A T A L V F L S F C GTCATCTTCATTGTAGTGAGGTCCTGCAGGAAGAATCGGCAAGACCAGCAGCGGGACGTGGGAGACATAGGCATGAAGGATGCAAACACC CAGTAGAAGTAACATCACTCCAGGACGTCCTTCTTTAGCCGTTCTGGTCGTCGCCTGCACCCTCTGTATCCGTACTTCCTACGTTTGTGG V I F I V V R S C R K K S A R P A A D V G D I G M K D A N T ATTCAGGGGCTCAGCCTCTCAGGGTAACTGGATGAGTCCTGGGCAGATGATAACCCCCGACACCATGGCCTGGCTGCCCACTCCCTCAGG TAAGTCCCCGAGTCGGAGAGTCCCATTGACCTACTCAGGACCCGTCTACTATTGGGGGCTGTGGTACCGGACCGACGGGTGAGGGAGTCC IOGLSLSG (SEQIDNO:18) GGAGGAAAGAGAGATCCCAGTATGCACCCCTCAGCTTTCATAAGGGGGAGCCTCAGGACCTATCCAGGTCAAGAAGCCACCAACAATGAG CCTCCTTCTCTCTAGGGTCATACGTGGGGAGTCGAAAGTATTCCCCCTCGGAGTCCTGGATAGGTCCAGTTCTTCGGTGGTTGTTACTC TACTCAGAGATCAAGATCCCCAAGTAAGAAAATGCAGAGGCTCGGGCTTGTTTGAGGGTTCACGACCCCTCCAGCAAAGGAGTCTGAGGC ATGAGTCTCTAGTTCTAGGGGTTCATTCTTTTACGTCTCCGAGCCCGAACAACTCCCAAGTGCTGGGGAGGTCGTTTCCTCAGACTCCG CCAAACTCTCCCTTTCCCCATCCAATCGGTCCACACTCCCCGCCCTGGCCTCTGTACCCACCATTCTCCTCTGTACTTCTCTAAGGATGA + 1440 A (SEQ ID NO:1)

Page 1

																							<u>H</u>	ΓE	$\mathcal{D}$	Kμ	18			
CGC	CAC	GAG	TGG	ACA	ACC	ATC.	AGG	GAG	CCA	GGA	CAC	AGA	ĢGG	GCA	GAG	CAAI	GTC	AGC	ATT	GGC	SCCC	CT	rcc	TCA	GAT	cc.	TAT	CAT	CTT	
GCC	GTG	CTC	ACC	TGT	TGG	TAG	TCC	CTC	GGT	CCTI	GTG.	TCTI	CCC	CGT	CTC	STT	CAG	TCG	TAA	CGG	GGG	GA,	AGG.	AGT	CTAC	GGG/	ATA	GTA	GAA	90
					,																									
000		CAC	TAC		`` `AGAI	CCT	TCA	C C V	A C A	TCT	T A A I	CTT		тст	TCC	200	TCC	ררר	A C T (	TC	r T C /	ver.	۱.T.C.	oc ta	C A A .	A T C 1	C	ACTI		
			<del></del>										+			<del></del>					-+								+	180
CCC	:111	GTC	ATC	GGG	TCT	CCA	AGT	CCT	TCT	ACA	ATT(	GAA'	7 T T	ACA.	AGCI	CCC.	ACG	GGG	TCA	GAC	AAG	rcg.	FAC	CGA	TT:	FAG	GTG'	TGA	GGC	
																							М	Α	E	I	н	T	Р	
TAT	TTCT	TCC	TTG	AAG	AAA	CTG	TTA	TCT	TTA	CTC	AAT	GGC.	TTC	GTG	GCT	STG	TCT	GGC	ATC	ATC	TAC	STT(	age:	CTG	3GC/	ATT(	GGT	GGT	AAA	270
AT A	AAGA	AGG	ÁAC	TTC	TTT	GAC	AAT.	AGA	AAT	GAG	TTA	CCG	AAG	CAC	CGA	CAC	AGA	CCG.	TAG	TAG	GATO	CAA	CCG	GAC	CCG	TAA	CCA	CCA	TTT	2,0
v	c	c		v	v	ı	,	c	,	,	M	C	c	V	۸	v	c	C	ı	ı		v	C		c	;	C	c	к	
1	3	3	L		Α.	L	L	3	L	L	М	G	Г	٧	^	•	3	u			_	•	3	_	G	i	G	G	Λ.	
TGT	FGGA	GGG	GCC	тст	CTG	ACG.	AAT	GTC	CTC	GGGI	CTG	TCC	TCC	GCA	TAC	CTC	CTT	CAC	GTŢ	GGC	AAC	TG	rgc	CTG	GTGA	ATG	GGA	TGC/	ATC	
AC/	ACCT	ccc	CGG	AGA	GAC	TGC	TTA	CAG	GAG	CCCI	GAC	AGG	AGG	CGT.	ATG	GAG	GAA	GTG	CAA	CC.	TTG	SAC	ACG	GAC	CAC	TAC	CCT	ACG:	TAG	360
C	G	G	A	S	L	T	N	٧	L	G	L	S	S	А	Υ	L	L	Н	٧	G	N	L	С	L	٧	М	G	С	I	
ACC	GGTA	CTG	СТТ	GGC	TGT	GCC	GGG	TGG	TAT	GGA	GCG.	ACT.	ΔΑΑ	GAG.	AGC	AGA	GGC.	ACG	CTC.	TTG	TTT	rgc,	ATC	CTG	TCA	ATG	GTT:	ATT(	GTC	
			<del></del>		SACA	-+-					<del></del>		+			+-		<del></del>					<del></del>						<del>+</del>	450
		J, . J	0,.,,																											
T	٧	L	L	G	С	Α	G	W	Υ	G	A	T	K	Ε	S	R	G	T	L	L	F	C	I	L	S	M	٧	I	٧	
сто	TATO	ATG	GAA	GTT	ACA	GCT	פרנ	ΔΓΔ	GTG	STO	CIT	CIT	TTC	TTT	CCA	ATT	GTT	GGA	GAT	STG	3001	rtge	GAA	CAC	ACC.	TTC	GTG.	ACCI	CTG	
			+	<del></del>	TGT	+-							+			+					<del></del>			+		-		•		540
GAL	3 I A G	IAC	C 1 1	САА	1161	CGA	Cuu	161	LAC	LAGI	SAAI	GAA	ААС	<b>AAA</b>	991	IAA	CAA	C C 1 1	LIA	JAC	. 00,	4MC	- ' ' '		1 00,	4861	JAC	1000	346	
L	I	M	Ε	٧	Т	A	Α	T	٧	V	L	L	F	F	P	I	٧	G	D	٧	Α	L	Ξ	Н	T	F	٧	T	L	
400		A A T	T A C	A C A	GGT	T 4 C	A A C	C 4 C	C C A	CAC	CAC	T A T	тст	A C A	C V C.	tee	A A C	TTC	· TC	4 T C 6	- A C /		7.	A A C	TCC.	TOT	ccc	CTC.	A A T	
			+										+			+-		• • •			<del></del>		<del></del>		-	+			<del>+</del>	630
TCC	:110	TTA	ATG	TCT	CCA	ATG	TTG	CTC	GGT	CTG	CTG	ATA	AGA	TGT	GTC	ACC	HG	AACI	CAG	IAC	. 10	TIC	Αi	HC	ACGA	ACAI	CCCI	LAL	IIA	-
R	K	N	Υ	R	G	Υ	N	Ε	Р	D	D	Υ	s	Т	Q	W	N	L	٧	М	Ε	K	L	K	С	С	G	٧	N	
AAC	TAC	ACA	GAT	TTT	TCT	GGC.	TCT	TCC	TTC	GAA	ATG	AC A	ACG	GGC	CAC	ACC	TAC	CCC	AGG	AGT	rgci	GT	AAA	TCC	ATC	3GA/	AGT	GIG		720
TTO	SATG	TGT	CTA	AAA	AGA	CCG	AGA:	AGG	AAG	CTT	TAC	TGT	TGC	CCG	GTG	TGG.	ATG	GGG.	TCC	TCA	ACGA	ACA	[17.	AGG	ΓAG	CCT	TCA	CAC	AGG	
N	Υ	T	n	F	S	G	S	S	F	F	м	Т	т	G	н	т	Y	Р	R	s	c	c	ĸ	s	ī	G	s	٧	s	
	·	•	•	•		•	•	Ŭ	•	_	••	•	·	_		•	•	•	•	•	_			_	•	_				
TGI	TGAC	GGA	CGC	GAT	GTG	TCT	CCA.	AAC	GTÇ.	ATC	CAC	CAG	AAG	GGC	TGT	TTC	CAT	AAA	CTC	CTA	AAA	TC	ACC.	AAG	ACT(	CAG	AGC	TTC		810
AC/	ACTG	ССТ	GCG	CTA	CAC	AGA	GGT	TTG	CAG	TAG	GTG	GTC	TTC	CCG	ACA	AAG	GTA	TTT	GAG	GAT	ודדו	AG	rgg	TTC	TGA	GTC.	TCG.	AAG	TGG	010
_	_	_		_			_					_		_	_	_		•					_		_	_	_	_	-	
-	_	_		_	٧	•			-	•		_		_	_			•	L	_								F	15	C ID
СТО	SAGT	GGG	AGC	тст	CTG	GGA	GCT	GCA	GTG.	ATA,	CAG	AGG	TGG	GGG	тст	CGC	TAT	GTT	SCG	CAG	SCTO	GT	TT	GAAI	CTG	CTGI	GCC	TAA	AGC	MO::
			, -		GAC								_			_,												-		900
																														~
L	S	G	S	S	L	G	A	A	٧	I	a	R	W	G	S	R	Y	٧	Α	a	Α	G	L	Ε	L	L	A	. (	X	ID
																													N1	1. 17

Page 2

HIEDK 48	
GATCCCCCCCCCTAGGCCTCCCAAAGTGCTGGGTTTACCAGCGTGAGCCACCACGCTGGGCTTCCTGCATCCTTTTAAGGTTCCTGAGGG	990
CTAGGGGGGGGGATCCGGAGGGTTTCACGACCCAAATGGTCGCACTCGGTGGTGCGACCCGAAGGACGTAGGAAAATTCCAAGGACTCCC	990
TCTGCCTGAGAGGÀGCTGTCCCTGAATCTCCATGCAGCCCCACCTGCCACATCACCAAGACATCTTTGCCAGCAACACTTCCTCC	1080
AGACGGACTCTCCTCGACAGGGACTTAGAGGTACGTCGGGGGTGGACGGTGTAGTGGTTCTGTATGTTAGAAACGGTCGTTGTGAAGGAGG	1000
TTGCAGATTACAAGCATAGCTAATGCCACCAGCAGCAGACAAGACCGATTCGCTGGCCTCCATTTCTTCAACCCAGTGCCTGTCATGAAACTT	1170
AACGTCTAATGTTCGTATCGATTACGGTGGTGGTCTGTTCTGGCTAAGCGACCGGAGGTAAAGAAGTTGGGTCACGGACAGTACTTTGAA	
GTGGAGGTCATTAAAACACCAATGACCAGCCAGAAGACATTTGAATCTTTGGTAGACTTTAGCAAAACCCTAGGAAAGCATCCTGTTTCT CACCTCCAGTAATTITGTGGTTACTGGTCGGTCTTCTGTAAACTTAGAAACCATCTGAAATCGTTTTGGGATCCTTTCGTAGGACAAAGA	1260
TGCAAGGACACTCCTGGGTTTATTGTGAACCGCCTCCTGGTTCCATACCTCATGGAAGCAATCAGGCTGTATGAACGAGGGCCTCCTGGC	
ACGTTCCTGTGAGGACCCAAATAACACTTGGCGGAGGACCAAGGTATGGAGTACCTTCGTTAGTCCGACATACTTGCTCCCGGAGGACCG	1350
TTTCCCTGTGGGCTTCTGAGAAAGGTTTCTGGAACTCCCACCACCACCACCACCACCAGCCAG	1440
AAAGGGACACCCGAAGACTCTTTCCAAAGACCTTGAGGGTGGTGGGGGGTGATGTCAGGGTCGGTC	
GATATCCTGGATCTCTGCTTTTGATTAAAAGGTGACGCATCCAAAGAAGACATTGACACTGCTATGAAATTAGGAGCCGGTTACCCCATG	1530
CTATAGGACCTAGAGACGAAAACTAATTTTCCACTGCGTAGGTTTCTTCTGTAACTGTGACGATACTTTAATCCTCGGCCAATGGGGTAC	
GGCCCATTTGAGCTTCTAGATTATGTCGGACTGGATACTACGAAGTTCATCGTGGATGGGTGGCATGAAATGGATGCAGAGAACCCATTA	. 1020
CCGGGTAAACTCGAAGATCTAATACAGCCTATGATGCTTCAAGTAGCACCTACCCACCGTACTTACCTACGTCTCTTGGGTAAC	
CATCAGCCCAGCCCATCCTTAAATAAGCTGGTAGCAGAGAACAAGTTCGGCAAGAAGACTGGAGAAGGATTTTACAAATACAAGTGATGT	1710
GTAGTCGGGTCGGGTAGGAATTTATTCGACCATCGTCTCTTGTTCAAGCCGTTCTTCTGACCTCTTCCTAAAATGTTTATGTTCACTACA	
GCAGCTTCTCCGGTTCTGAGAAGAACACCTGAGAGCGCTTTCCAGCCAG	. 1800
CCTCACACAGTACAGTTTAATAAATGTGCATTTTGATTGTAAAAAAAA	
GGAGTGTCTCATGTCAAATTATTTACACGTAAAACTAACATTTTTTTT	

BNSDOCID: <WO___9831799A2_i_>

																											عبط			
GGC	GTO	CCC	ÇTG	CCT	GCC	CĄC	TCA	GTG	GCA	ACA	CCC	GGG	AGC	TGT	TTT	GTC	CTT	TGT	GGA	AGCC	TCA	GCA	GTT	CCC	TCT	TŢC	AGA	ACT	CAC	0.0
CCG	CAC	GGG	GAC	GGA	CGG	GTG	AGT	CAC	CGT	TGT	GGG	ccc	TCG	ACA	AAA	CAG	GAA	ACA	VC C T	rcgo	AGT	CGT	CAA	GGG	AGA	AAG	TCT	TGA	GTG	90
				1																										
TGC	CAA	4GA(	ccc	TĠĀ	ACA	GGA	GCC	AC C	ATG	CAG	TGC	ттс	AGC	TTC	ATT	AAG	ACC	ATO	ATO	SATO	сто	TTC	AAT	TTG	стс	ATC	777	CTG	TGT	
ACG	GTI	TCTO	GGG	ACT	TGT	CCT	cgg	TGG	TAC	GTC	ACG	AAG	TCG	AAC	TAA	TTC	TGG	TAC	TAC	TAG	GAG	AAG	TTA	AAC	GAG	TAG	AAA	GAC	ACA	180
										_	_		•	_			<b>.</b>	м				_					_		_	
									М	a	L	7	5	,	1	K	,	Γř	rı	ŀ	Ļ	r	N	L	L	ī	۲	Ĺ	Ĺ	
GGT	GC/	AGC	CTG	TTG	GCA	GTG	GGC	ATC	TGG	GTG	TCA	ATC	GAT	GGG	GCA	TCC	TTT	CTO	AAC	GATO	TTC	GGG	CCA	CTG	TCG	TCC	AGT	GCC	ATG	270
CCA	CGI	rcad	GAC	AAC	CGT	CAC	CCG	TAG	ACC	CAC	AGT	TAG	ĊTA	ccc	CGT	AGG	ΑΑΑ	GAC	:110	TAG	AAG	ccc	GGT	GAC	AGC	AGG	TCA	ĊGG	TAC	270
G	Α	А	L	L	А	٧	G	i	W	٧	s	ı	D	G	Α	S	F	L	K	i	F	G	P	Ĺ	S	S	S	Α	М	
			<del></del>	<del></del>		<del>- ; -</del>			+		-		+		GTC	-+-							<del>:</del>			- 1,-		+ -	-+	360
GTC	AAA	ACAC	STIG	CAC	CCG	ATG	AAG	GAG	TAG	CGT	CGG	CCG	CAA	CAC	CAG	AAA	LCGA	KGA/	ACCA	AAAG	GAC	CCG	AUG	AIA	CCA	EGA	TTC	TGA	CTC	
Q	F	٧	Ν	٧	G	Υ	F	Ļ	l	Α	Α	G	٧	٧	٧	F	A	L	G	F	L	G	С	Y	G	Α	Κ	T	Ε	
AGC	AAC	GTG1	GCC	стс	GTG	ACG	TTC	TTC	TTC	ATC	стс	стс	стс	ATO	TTC	ATT	GCT	GA(	GTI	rgc A	.GCT	GCT	GTG	GTC	GCC	TTG	GTG	TAC	ACC	
TCC	TTO	CACA	CGG	GAG	CAC	TGC	AAG	AAG	AAG	TAG	GAG	GAG	<del>;                                    </del>	TAC	SAAG	TAA	CGA	CTO	CAA	ACGT	CGA	CGA	CAC	CAG	CGG	AAC	CAC	ATG	TGG	450
		_			.,	<b>~</b>	_	_	-					,	_	,		_	14				v	.,			.,	v	Т	
3	κ.	·	А	L		ŀ	r	r	г	,	L	L	L	١	г	ı	А	C	٧	A	А	^	٧	٧	А	L	٧	ı		
ACA	ATO	GC	GAG	CAC	TTC	CTG	ACG	TTG	CTG	GTA	GTG	CCT	GCC	ATO	AAG	AAA	GAT	TA	GG	TTCC	CAG	GAA	GAC	TTC	ACT	CAA	GTG	TGG	AAC	540
TGT	TAC	CGA	CTC	GTG	AAG	GAC	TGC	AAC	GAC	CAT	CAC	GGA	CGG	TAG	TTC	TTT	CTA	ATA	ACC/	AAGG	GTC	CTT	CTG	AAG	TGA	GTT	CAC	ACC	TTG	•
Т	.M	Α	ε	Н	F	L	Ţ	Ļ	L	٧	٧	P	A	i	к	K	0	Υ	G	s	Q	Ε	D	F	т	Q	٧	W	N	
۸۲۲	۸۲۱	~ A T (	2	ccc	רדר	AAC	TCC	TCT	eer	TTC	۸۲۲	A A C	TAT	۸۲۵	GAT	TTT	.C.V.C	CAC	· T C /	ירני	. T A C	TIC	A A A	CAC	: A A C	AGT	מכר	TTT	درد	
			-			<del></del>		+					+		CTA	<del></del>							+							630
	,, ,,	3.70			unu	, , ,	ACG.	~~			, , ,	, , ,							,,,,,		,,,,,					., .,			-	
Ţ	T	M	K	G	L	K	С	С	G	F	T	N	Y	T	D	F	Ε	D	S	Р	Υ	F	K	E	N	S	A	F	Р	
CCA	TTC	CTGI	TGC	AAT	GAC	AAC	GTC.	ACC	AAC	ACA	GCC	AAT	GAA	ACC	TGC	AÇC	AAG	CAA	AAC	GGCT	CAC	GAC	ÇAA	AAA	GTA	GAG	GGT	TGC	ттс	700
GGT	AAC	GAC	ACG	TTA	CTG	TTG	CAG	TGG	TTG	TGT	CGG	TTA	CTT	TGC	ACG	TGG	TTC	GT	TTC	CGA	GTO	CTG	GTT	TTT	CAT	стс	CCA	ACG	AAG	720
Р	F	۲	٦	N	n	NI.	v	т	N	т	Δ	N	Ε	т	۲	т	K	a	ĸ	Δ	н	0	Ω	ĸ	v	F	G	c	F	
·	·	Ū	Ū	.,	Ü	.,	•	•		•		••		•	Ū	•				^		,	•	``	·	_		•	-	
	-							+					+		GGT					-	<del></del>		+			-+-				810
TTA	GTO	CGAA	AAC	ATA	CTG	TAG	GCT	TGA	TTA	CGT	CAG	TGG	CAC	CCA	CCA	CAC	CGT	CGA	ACC1	TTAA	CCC	CCG	GAG	CTC	GAC	CGA	CGG	TAC	TTA	
N	۵	L	L	Υ	D	1	R	Ţ	N	A	٧	T	٧	G	G	٧	A	A	G	1	G	G	L	E	L	A	A	M	N	
TGI	GTI		GTA	TCT	GTA	CTG	<b>C A A</b>	TCT	ልሮል	ΔΤΔ	ACT	٤٢٨	רדד	CIC	CCT	ств	CCA	LC T A	AC T	SCTO	:cr4	СДТ	GGG	AAC	TRT	GAA	GAG	GCA	ccc	
			+					+					+		GGA	-		+		<del></del>			<del></del>			-+-		• • •	+	900
					<b>5</b> 71	<b>-</b>	J 1 1						J		.007			<b>U</b> A .	571						CM		_			

HPWAE25

TGGCAAGCAGCAGTGATTGGGGGGAGGGACAGGATCTAACAATGTCACTTGGGCCAGAATGGACCTGCCCTTTCTGCTCCAGACTTGGGGACCTGGGCAGACTGGGGAAAGACCCCCTCCCCTGTCCTAGATTGTTACAGTGAACCCGGTCTTACCTGGACGGGAAAGACGAGGTCTGAACCCC WOAAYIGGGORI. (SOO IDNO: 20) 1080 GCCAGTTCTGTTGCCCATTCCCCCAGTCTATTAAACCCTTGATATGCCCCCTAGGCCTAGTGGTGATCCCAGTGCTCTACTGGGGGATGA

CGGTCAAGACAACGGGTAAGGGGGTCAGATAATTTGGGAACTATACGGGGGGATCACCACTAGGGTCACGAGATGACCCCCTACT

GAGAAAGGCATTTTATAGCCTGGGCATAAGTGAAATCAGCAGAGCCTCTGGGTGGATGTGTAGAAGGCACTTCAAAATGCATAAACCTGT CTCTTTCCGTAAAATATCGGACCCGTATTCACTTTAGTCGTCTCGGAGACCCACCTACACATCTTCCGTGAAGTTTTACGTATTTGGACA

Päge 1

																							t	T	۲	二	- %	صا		
AAAA	AA	ACA	AGG	TC	000	ACAG	CAA	AG	AAA	AGGA	ATA	GGA	TC	AAGA	GAT	ACG	TG	GCTO	CTG	GCA	GAG	SCA/	GCA	TGA	AT	TCG	TGA	CTT	CA	90
TTTT	TT	TG	TCC	AG	GGG	TGTO	GTI	TC	TTT	TCCT	TAT	CCT	AG	TTC	CTA	TGC	CAC	CGA	GAC	CGI	СТО	GT	rcgt	ACT	TA	AGC	TACT	GAA	GŤ	
					į.																			М	N	S	М	T	S	
CAG	rtr	^C60	STGO	CC.	AAT	TCTO	STGT	rTG	GTG	GTG	SCAC	ccc	AC.	AAT(	GTT	AT(	CT	GTG	4000	CAC	GGA	ATT,	ATGT	CTO	CAC	GŢG	2000	TGT	ΑŢ	100
GTC																														180
		_	.,			c	.,		v	V	٨	P	ы	N	G	Y	Р	٧	Т	ρ	G	i	М	S	н	٧	Р	L	Υ	
Α '	٧	Ρ	٧	Д	N	3	٧	_	•	•	^	•	• •	"	J	•														
CAA																														270
GGTT	T G	TEG	GTC	GC	GTT	CAG	GTG	GAT	CAA	GGA	CCC.	rtg	GGT	GGA	TCAA	AACI	CAC	AGC	TTA	CAC	TTA	ECE	Gice	GGA	CAC	GTC	111	CGA	i A C	
ρ	N	s	Q	٩	۵	٧	н	L	٧	Р	G	N	Ρ	P	S	L	٧	S	N	٧	N	G	Q	Ρ	٧	a	K	A	L	
AAAG	AΑ	GGC	AAA	ACC	TTG	GGG	GCC.	ATC	CAG	ATC.	ATC	ATT	GGC	CTG	GCT	CAC	ATC	GGC	CTC	GGC	тсс	ATC	ATG	GCG.	ACG	GTT	CTC	GTA	GGG	360
TTTC																														300
v	F	c	K	т	i	G	А	ı	a	I	į	1	G	L	Α	н	į	G	L	G	s	I	м	A	Т	٧	L	٧	G	
ĸ.	_	u		•	•	-							T 0.0		000	T T C	TCC		4 T.C	A T T	TCA	CCA	TOT	ctr	TCC	.c.r.c	GC A	CC A	GAA	
GAAT CTTA																														450
CTTA	TG	GAC	AGA	IAA	AGI	AAG	AIG	L L !	LLE	ААА	666	AAG	ACC	.661	CCG	AAC	~~					_	_						_	
Ε	Y	L	S	i	S	F	Y	G	G	F	Ρ	F	W	G	G	L	W	F	I	I	5	G	S	L	S	٧	А	А	E	
AATC																														540
TTAG	TC	GGT	ATA	AGA	ATA	ACG	GAC	GAC	CAGA	ACCG	TCA	AAC	ĊC	GAAC	TTG	TAG	CAC	STCA	CGT	TAG	ACG	AGA	CGT	CAA	CCT	TCAG	TAT	GAG	AAG	
N	a	Ρ	Υ	s	Y	C	L	L	S	G	s	L	G	L	N	i	٧	S	A	ı	С	s	A	٧	G	٧	ï	L	F	
ATCA		CAI	· C T A	A G T	ΓΔΤΊ	זררר	· C A C	ירכי	ΔΤΔ.	raco	TAC	ccc	GA	CTAT	TAT	CCT	TAI	cgco	TGG	GGT	GTO	SAAC	CCT	GGA	AT	GGC	SATT	тст	GGÇ	
TAGI																														630
		_				_		_	v		v			v	v	P	¥	Δ	W	c	v	N	Р	G	M	A	ı	s	G	
ì	T	D	L	S	i	Р	н	Р	Ţ	Д	ī	r	U	•	•	•		^	"	J	•	.,	·	Ū	•	•				
GTG																														720
CACC	SAC	CGA	CAG	AA	GAC	GGA	GAC	CT	CAA.	ACC	STAC	CGI	ГАС	GCG	TAGA	AG	GGT	GAA	4000	SAC	GGT	CAAI	CCAE	SAC	SAC.	AGI	IAGI	1166	HA	
٧	L	L	٧	F	С	L	L	Ε	F	G	1	A	С	A	S	S	Н	F	G	С	۵	L	٨	С	С	0	S	5	N	
GTG	AG1	TGT	CATO	:TA	TCC	ΑΑΑ	CATO	TA.	TGC	AGC	444(	cc	AGT	GAT	CACC	cc	AGA	ACC	GGT	GACI	CTC.	ACC.	ACC/	AAG	TTA	TTÇ	CAG	rga(	ATC	810
CAC																														
																		Р												
٧	3	٧	t	1	-	14		1	^		• 4	•	•	•	•	•	-													

39 HTPEF86	Page 2
AAGCAAATAAGTAAGGCTACAGATTCTGGAAGCATCTTTCACTGGGACCAAAAGAAGTCCTCCCTTTCTGGGCTTCCATAACCCAC	
TTCGTTTATTCATTCCGATGTCTAAGACCTTCGTAGAAAGTGACCCTGGTTTTCTTCAGGAGGAGGGAAAGACCCGAAGGTATTGGGT	:
OANK (SEQIDNO: 21)	
TEGTTECTGTTETGACAGETGAGGAAACGTETETECCACTGTTTGTACTETCACCTTCATTETTCAATTCAGTCTAGGAAACCATGETC	3 + 990
AGCAAGGACAAGACTGTCGACTCCTTTGCAGAGAGGGTGACAAACATGAGAGTGGAAGTAAGAAGTTAAGTCAGATCCTTTGGTACGAC	
TTCTCTATCAAGAAGAAGACAGAGATTTTAAACAGATGTTAACCAAGAGGGACTCCCTAGGGCACATGCATCAGCACATATGTGGGGCA	[ oso
AAGAGATAGTTCTTCTGTCTCTAAAATTTGTCTACAATTGGTTCTCCCTGAGGGATCCCGTGTACGTAGTCGTGTATACACCCGT	+ 1080 4
CAGCCTCTGGGGCCTTGGCACACCCATTCGTGTGCTCTGCTGCATGTGAGCTTGTGGGTTAGAGGAACAATATCTAGACATTCAATC	T + 1170
GTCGGAGACCCCGGAACCGTGTGGGTAAGCACACGAGACGACGTACACTCGAACACCCAATCTCCTTGTTTATAGATCTGTAAGTTAG	
CACTETTECAATTGTGCATTCATTTAATAAATAGATACTGAGCATTCAAAAAAAA	
ATT A CALL OF TARCE COLOR OF TARCE AND TARTE A TOTAL COLOR CALCULATION ACCULATION TO THE TARTE A TOTAL COLOR CALCULATION ACCULATION	

Päge 1

																								HS	B	$\mathbb{R}^{2}$	EC	12		
CACG	ACGAGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGACTGCAGACTAAACCAGTCATTACTTGTTTCAAGAGCGTTCTGCTA  TGCTCGTCCCAGAGCCCGATCAGTACCGCAGGGGCAGAGCCTCTGACGTCTGATTTGGTCAGTAATGAACAAAGTTCTCGCAAGACGAT															90														
GTGC	TC	STC	CAG	AGC	CCG	AT	AGI	rac (	GC/	AGGG	GGC	AGAC	SCC.	гст	GAC	STC	rgai	TTO	GTO	CAGT	TAAT	GAA	CAA	AGT	TCT	CGC	CAAG	SACG	AT	
				1				M	A	s	Р	S	R	R	L	0	T	K	Ρ	٧	Į	T	С	F	Κ	S	٧	L	L	
ATCT	ACA	ACT.	TTT/	TTI	гтст	GG	ATC	ACTO	GCC	STT	ATC	стт	TT (	GCA	GTT	GGC.	ATT	rggo	GC,	AAG	STGA	GCC	TGO	GAGA	ATI	rac 1	777	CTC	:TT	180
ATCT TAGA	TG	TGA	AAA1	[AA	AAGA	VCC.	TAG	TGA	CCGC	CAA	TAG	GAA	GAA	CGT	CAA	CCG	TAA	ACCO	CCG	TTC	CACI	CGC	SAC	CTCT	TAA	4 T GA	\AA/	4GA0	SAA	
ı	Y	Ŧ	F	ι	F	W	ľ	T	G	٧	I	L	L	Α	٧	G	I	W	G	Κ	٧	S	L	Ε	N	Y	F	S	L	
TTAA	ATO	GAG.	AAG	SCCA	4CC <i>A</i>	ĄŢ	GTC	ccc.	TTC	GTG	CTC	ATT	GCT.	ACT	GGT.	ACC	GTC.	ATT	ATT	CTT.	TTG	GC/	ACC.	111	GT	TGT	111	GCTA	ACC	270
AATT	TA	CTC	TTC	GG	TGGT	TA	CAG	GGG	AAG	CAC	GAG	TAA	CGA	TGA	CCA	TGG	CAG	TAA	TAA	GAA	AAC	CCG.	TGG.	ΔΑΑΙ	CCA	ACA	AAA	CGAT	rgg	
L	N	ε	κ	Α	T	Ņ	٧	Р	F	٧	L	I	A	T	G	T	٧	1	i	L	L	G	Т	F	G	С	F	A	T	
TGC	CGA	GCT	TCT	GCA	TGG	ATG	CTA.	AAA	CTG	TAT	GCA	ATG	ŢŦŦ	CTG	ACT	стс	GTT	TTT	TTG	GTC	GAA	CTG	GTC	GCT	GCC.	ATC	GTA	GGA	TTT	360
ACG	GC T	CGA	AGA	CGT.	ACC.	TAC	GAT	TTT	GAC	ATA	CGT	TAC	AAA	GAC	TGA	GAG	CAA	AAA.	AAC	CAG	CTT	GAC	CAG	CGA	CGG	TAG	CAT	CCT	AAA	000
С	R	A	S	А	w	М	L	Κ	L	Υ	Α	M	F	L	Т	L	٧	F	L	٧	Ε	L	٧	A	Α	I	٧	G	F	
GTT	TTC	AGA	ÇAT	GAG	ATT	ΑĄG	AAC	AGC	TTT	AAG	AAT	AAT	TAT	GAG	AAG	GCT	TTG	AAG	CAG	TAT	AAC	тст	ACA	GGA	GAT	TAT	AGA	AGC	CAT	450
CAA	AAG	TCT	GTA	стс	TAA	TTC	TTG	TCG	AAA	TTC	TTA	TTA	ATA	CTC	TTC	CGA	AAC	TTC	GTC	ATA	TTG	AGA	TGT	сст	CTA	ATA	TCT	TCG	GTA	
٧	F	R	н	Ε	1	K	N	S	F	K	N	N	Y	E	ĸ	Α	L	κ	a	Υ	N	s	T	G	D	Υ	R	S	н	
GCA	GTA	GAC	AAG	ATC	CAA	AAT	ACG	TTG	CAŢ	TGT	TGT	GGT	GTC	ACC	GAT	TAT	AGA	GAT	TGG	ACA	GAT	ACT	AAT	TAT	TAC	TCA	GAA	AAA	GGA	540
CGT	CAT	стс	TTC	TAG	GTT	TTA	TGC	AAC	GTA	ACA	ACA	ACCA	CAG	TGC	CTA	ATA	TCT	ĊТА	ACC	TGT	CTA	TGA	TTA	ATA	ATG	AGT	CTT	TTT	сст	
A	٧	D	κ	ļ	a	N	τ	L	н	С	С	G	٧	T	D	Y	R	D	W	T	D	T	N	Υ	Υ	s	Ε	K	G	
TTT	сст	AAG	AGT	TGC	TGT	AAA	CTT	GAA	GAT	TGT	TACT	rcc#	ŲÇΑ	AGA	AGAT	rgç	GAC	AAA	GTA	AAC	AAT	GAA	GGT	TGT	TTT	'A T A	AAG	GTG	ATG	630
																													TAC	
F	Р	K	s	С	С	K	L	Ε	D	С	Ţ	Ρ	Q	R	D	A	О	K	٧	N	N	Ε	G	С	F	ı	κ	٧	М	
ACC	ATT	TATA	AGAG	TCA	GAA	АТС	GGA	GTO	GTT	GCA	4 G G A	AATI	гтс	: 7 7 :	FGG/	AGȚ	rgc1	TGC	TTO	CAA	CTG	ATT	GGA	ATC	TTT	·c TC	:GCC	TAC	TGC	720
TGG	TAA	ATA	CTC	AGT	CTT	TAC	CCT	CAG	C A A	ACG.	TCC	TTA	AAG	SAA	4CC1	TCA	AC G	ACC	AAG	GT1	GAC	TAA	cc1	TAC	AAA	\GAG	CGG	ATG	ACG	, 20
Т	I	ī	Ε	s	Ε	М	G	٧	٧	A	G	i	s	F	G	٧	A	С	F	a	L	l	G	Ī	F	L	A	Y	С	
сто	TCI	rcg ⁻	rgco	ATA:	ACA	AAI	ΓΑΑΟ	CAC	STAT	ΓGΑ	GAT.	AGT	STA	ACC	CAA	TGT	ATC1	GTO	GGG	CCTA	ATTO	сто	TC	TAC	TTI	raac	GAC	CATT	TAG	- 810
GAG	AG	AGC/	ACGO	TAT	TGT	TT	ATTO	GGT	CATA	AC T	CTA	TCA	CAT	TGG	GTT	ACA	TAGA	AGAC	CCC	GGA1	TAAG	GAC	AGA	ATGO	SAAA	ATTO	CTO	STAA	AATC	:
L	s	R	A	i	Ţ	N	N	a	Y	Ε	i	٧		65	X	ב ע	D	N	<u>)</u> ;	22	2)									
GGT	cco	CCC	CTGI	rga,	ATTA	AG Ą	AAG	TTG	CTT	GGC	TGG	AGA	ACT	GAC	AAC.	ACŢ	ACT	TAC	TGA	TAG	ACC	AAA	AAA	CTA	CAC	CAG	FAG	STT		900
CCA	ree.	GGG	GAC	AC T	[AA]	rc T	TTC	AACI	GAAI	CCG.	ACC	TCT	TGA	CTG	TTG	TGA	TGA	ATG	ACT	ATC	TGG	TT.	rŤT	GAT	STG	STC	ATC	CAAC	CTAA	١ -

HSBRFOZ

TGTTTGCTATGTTC TAAGTCCACCTTCTATCCCATTCATGTTAGATCGTTGAAACCCTGTATCCCTCTGAAACACTGGAAGAGCTAGTAA
ACAAACGATACAAGATTCAGGTGGAAGATAGGGTAAGTACAATCTAGCAACTTTGGGACATTTGTGACCTTCTCGATCATT

1080

Page 1

ACGAGCATTGCCGCTCTCTCGGTGAGCGCAGCCCCGCTCTCCGGGCCGGGCCTTCGCGGGCCACCGGCGCCATGGGCCAGTGCGG	CATC + 90
STGCTCGTAACGGCGAGAGAGCCACTCGCGTCGGGGGGAGAGGCCCGGCCCGGAAGCGCCCGGTGGCCGCGGTACCCGGTCACGCC	GTAG
меосе	I
ACCTCCTCCAAGACCGTGCTGGTCTTTCTCAACCTCATCTTCTGGGGGGGCAGCTGGCATTTTATGCTATGTGGGAGCCTATGTCTT	CATC
TGGAGGAGGTTCTGGCACGACCAGAAAGAGTTGGAGTAGAAGACCCCCCGTCGACCGTAAAATACGATACACCCTCGGATACAGAA	<del></del>
T S S K T V I V F L N L I F W G A A G I L C Y V G A Y V F	ī
	•
ACTTATGATGACTATGACCACTTCTTTGAAGATGTGTACACGCTCATCCCTGCTGTAGTGATCATAGCTGTAGGAGCCCTGCTTTT	
TGAATACTACTGATACTGGTGAAGAAACTTCTACACATGTGCGAGTAGGGACGACATCACTAGTATCGACATCCTCGGGACGAAAA	GTAG
TYDOYDH FFEDVYTLIPAVVIIAVGALLF	I
ATTGGGCTAATTGGCTGCTGTGCCACAATCCGGGAAAGTCGCTGTGGACTTGCCACGTTTGTCATCATCCTGCTCTTGGTTTTTGT	CACA
TAACCCGATTAACCGACGACACGGTGTTAGGCCCTTTCAGCGACACCTGAACGGTGCAAACAGTAGTAGGACGAGAACCAAAAACA	<del></del>
	<b>+</b>
I G L I G C C A T I R E S R C G L A T F V I I L L L V F V	'
GAAGTTGTTGTAGTGGTTTTGGGATATGTTTACAGAGCAAAGGTGGAAAATGAGGTTGATCGCAGCATTCAGAAAGTGTATAAGAC	CTAC 450
CTTCAACAACATCACCAAAACCCTATACAAATGTCTCGTTTCCACCTTTTACTCCAACTAGCGTCGTAAGTCTTTCACATATTCTG	GATG
E V V V V L G Y V Y R A K V E N E V D R S I Q K V Y K T	Y
AATGGAACCAACCCTGATGCTGCTAGCCGGGCTATTGATTATGTACAGAGACAGCTGCATTGTTGTGGAATTCACAACTACTCAGA	ствв
TTACCTTGGTTGGGACTACGACGATCGGCCCGATAACTAATACATGTCTCTGTCGACGTAACAACACCTTAAGTGTTGATGAGTCT	
THACE FIGGI TGGGACTACGACGATCGGCCCGATAACTACATGTCTCTGTGCGACGTAACAACACCTAAAAAAAA	
NG TNPD A A S R A ! D Y V Q R Q L H C C G I H N Y S D	W
GAAAATACAGATTGGTTCAAAGAAACCAAAAACCAGAGTGTCCCTCTTAGCTGCTGCAGAGAGACTGCCAGCAATTGTAATGGCAG	CCTG
CTTTTATGTCTAACCAAGTTTCTTTGGTTTTTGGTCTCACAGGGAGAATCGACGACGTCTCTCTGACGGTCGTTAACATTACCGTC	GGAC
ENTOWFKETKNQSVPLSCCRETASNCNGS	i L
GCCCACCCTTCCGACCTCTATGCTGAGGGGTGTGAGGCTCTAGTTGTGAAGAAGCTACAAGAAATCATGATGCATGTGATCTGGGC	CGCA
CGGGTGGGAAGGCTGGAGATACGACTCCCCACACTCCGAGATCAACACTTCTTCGATGTTCTTTAGTACTACGTACACTAGACCCG	+ 720 GCGT
	. A
CTGGCATTTGCAGCTATTCAGCTGCTGGGCATGCTGTGTGCTTGCATCGTGTTGTGCAGAAGGAGTAGAGATCCTGCTTACGAGCT	
GACCGTAAACGTCGATAAGTCGACGACCCGTACGACACACGACGTAGCACACACGTCTTCCTCATCTCTAGGACGAATGCTCGA	IGAAG
I A F A A I O I I G M I C A C I V I C R R S R D P A Y E L	. F

Page 1

HTPBA27	
ACGAGCGCAGAGCTTGGGGCTTCCTTGGTCGCACCCACCACCTGCCTG	90
TGCTCGCGTCTCGAACCCCGAAGGAACCAGCGTGGGTGGG	
· ·	
TCTTTCCTGTGGCCAGCCCAGAACTGAAGCGCTGCGGCATGGCGCGCGC	100
AGAAAGGACACCGGTCGGGTCTTGACTTCGCGACGCCGTACCGCGCGGGGGGGG	
M A R A C L Q A V K Y L M F A F N	
TGTTCTTCTGGCTGGGAGGCTGTGGCGTGCTGGGTGTCGGCATCTGGCTGG	
ACAAGAAGACCGACCCTCCGACACCGCACGACCCACAGCCGTAGACCGGCGGTGTGTCCCCTCGAAGCGGTGCGACAGAAGAAGAAGAAGA	
TCCCGTCCCTGTCGGCTGCCAACCTGCTCATCATCACCGGCGCCTTTGTCATGGCCATCGGCTTCGTGGGCTGCCTGGGTGCCATCAAG	360 ·
AGGGCAGGGACAGCCGACGGTTGGACGAGTAGTAGTGGCCGCGGAAACAGTACCGGTAGCCGAAGCACCCGACGGACCCACGGTAGTTC	:
FP S L S A A N L L I I T G A F V M A I G F V G C L G A I K	
AGAACAAGTGCCTCCTGCTCACTTTCTTCCTGCTGCTGCTGCTGGTGTTCCTGCTGGAGGCCACCATCGCCATCCTCTTCTTCGCCTAC	<b>A</b>
TCTTGTTCACGGAGGACGAGTGAAAGAAGGACGACGACGACCACAAGGACGACCTCCGGTGGTAGCGGTAGGAGAAGAAGCGGATG	F 450
	_
CGGACAAGATTGACAGGTATGCCCAGCAAGACCTGAAGAAAGGCTTGCACCTGTACGGCACGCAGGGCAACGTGGGCCTCACCAACGCC	
GCCTGTTCTAACTGTCCATACGGGTCGTTCTGGACTTCTTTCCGAACGTGGACATGCCGTGCGTCCCGTTGCACCCGGAGTGGTTGCGG	ц.
T D K I D R Y A Q Q D L K K G L H L Y G T Q G N V G L T N A	
GGAGCATCATCCAGACCGACTTCCGCTGTGGCGTCTCCAACTACACTGACTG	T 630
CCTCGTAGTAGGTCTGGCTGAAGGCGACGACGCGCAGAGGTTGATGTGACCTGACCAAGCTCCACATGTTGCGGTGCGCCCATGGACTG	
W S T T U T B T K C C C C C C C C C C C C C C C C C C	r
CCTGCTGCTTGGAGTTCAGTGAGAGCTGTGGGCTGCACGCCCCCGGCACCTGGTGGAAGGCGCCGTGCTACGAGACGGTGAAGGTGTGG	
GGACGACGACCTCAAGTCACTCTCGACACCCGACGTGCGGGGGCCGTGGACCACCTTCCGCGGGCACGATGCTCTGCCACTTCCACACC	G
S C C L E F S E S C G L H A P G T W W K A P C Y E T V K V W	
TTCAGGAGAACCTGCTGGCTGTGGGCATCTTTGGGCTGTGCACGGCGCTGGTGCAGATCCTGGGCCTGACCTTCGCCATGACCATGTAC	T + 810
AAGTCCTCTTGGACGACCGCGACACCCGTAGAAACCCGACACGTGCCGCGACCACGTCTAGGACCCGGACTGGAAGCGGTACTGGTACATC	
L Q E N L L A V G I F G L C T A L V Q I L G L T F A M T M Y	
	١٢
GCCAAGTGGTCAAGGCAGACACCTACTGCGCGTAGGCCGCCCACCGCCGGCTTCTCTGCCAAAAGGACGCCCACGGGGAGATGGCCGC	
CGGTTCACCAGTTCCGTCTGTGGATGACGCGCATCCGGCGGTGGCGGCCGAAGAGAGACGGTTTTCCTGCGGGTGCCCCTCTACCGGCG	5
COVVKADTYCA. (SEQID'NO: 24)	

Page 2

CCACAGCTGCTTTTCCCACCACCAGCTTCGGTGTTCTGCCCCATGCTGGGAGGGA	000
GGTGTCGACGAAAAGGGTGGTCGAAGCCACAAGACGGGGTACGACCCTCCTCCCTC	990
CTGTTTCTGGAAGGECCTAGCTCAGGTGGCTTCAGGGCCTCCGGACCCCCCTGGGAGGGGTGGCCACGTGCTGGCTG	1080
GACAAAGACCTTCCGGGATCGAGTCCACCGAAGTCCCGGAGGCCTGGGGGGGACCCTCCCCACCGGTGCACGACCGAC	1080
CAGGGGTGGGAGGGCCTCCAGCACTTTTTATATTTACGTATTCTCCAAAGCAGTGTTCACACGGGAGCCAGCC	1170
GTCCCCACCCTCCCGGAGGTCGTGAAAAATATAAATGCATAAGAGGTTTCGTCACAAGTGTGCCCTCGGTCGG	1170

GACCTTTGTCCAACCGCGACCTCCTCGGCCCAGAACCGTAGGACCTCCACCGGGGTGACCAGGACGACGACGACGGCCCCGGCACCTGG

Page 1

HAIDQ59																														
GCGG	CG	CCG	ĢGC	CGC	AGC	AŢĞ	GGG	CGC	TTC	CGC	GGG	GGC	CTG	cgg	TGC	ATC.	AAG	TAC	CTG	CTG	TT	GGC.	TTC	AACC	:TĠ	CTC	TTC	rgg	CTG	90
cGCC	GC	GGC	CCG	GCG.	TCG	TAC	CCC	GCG	AAG	GCG	CCC	CCG	GAC	GCC	ACG	TAG	TTC	ATG	GAC	GAC	GAA	CCG.	AAG	TTG	SAC	GAG.	AAGA	ACC	GAC	-
				Į,	<u>.</u>	М	G	R	F	R	G	G	L	R	С	i	κ	Y	L	L	L	G	F	N	L	L	F	W	L	
GCT	GGA	TCG	GCC	GTC	ATT	GCT	TTT	GGA	CTA	TGG	TTT	cgg	TTC	GGA	GGT	GCC.	ATA	AAG	GAG	TTA	TCA	TCA	GAG	GAC	AAG	TCC	CCA	GAG	TAT	180
CGA	CCT	AGC	CGG	CAG	TAA	EGA	AAA	CCT	GAT	ACC	AAA	GCC	AAG	CCT	CCA	cĠĠ	TAT	TTC	CTC	AAT.	AGT	AGT	CTC	CTG	TTC	AGG	GGT	CTC	ATA	
A	G	S	A	٧	Ī	Α	F	G	L	W	F	R	F	G	G	A	Į	K	Ε	L	s	S	Ε	D	K	S	Ρ	Ε	Υ	
TTC	TAT	GTG	GGG	CTG	TAT	GTT	CTG	GTT	GGA	GCC	GGG	GCC	CTG	ATG	ATG	GCC	GTG	GGG	TTÇ	TTC	GGA	TGC	TGC	GGA	GCC	ATG	CGG	GAG	TCG	270
AAG	ATA	CAC	CCC	GAC	ATA	CAA	GAC	CAA	CC1	CGG	ccc	CGG	GAC	TAC	TAC	CGG	CAC	ССС	AAG	AAG	ССТ	ACG	ACG	CCT	CGG	TAC	GCC	CTC	AGC	270
F	Y	٧	G	L	Y	V	L	٧	G	A	G	Α	L	M	м	Α	٧	G	F	F	G	С	С	G	Α	М	R	Ε	s	
CAA			1					-										<del></del>	<del></del>				+					+	_	360
GTT	ACA	CAC	GAA	CCT	AGT	AAA	AAA	TGG	SAC	GAG	GAC	CAC	TAT	ΓΑΑΑ	CGA	CGA	CTT	CAT	TGG	TGA	CCT	CAT	AAA	CGA.	AAA	TAT	CCG	TTC	ccc	
Q	С	ν	L	G	S	F	F	Ţ	С	L	L	٧	1	F	Α	A	Ε	٧	T	Τ	G	٧	F	Α	F	ı	G	Κ	G	
GTA	GCT	ATO	CGA	CAT	GTI	rcag	ACC	ATO	ATE	TGAA	GAG	GCT	TAC	CAAT	GAT	TAC	СТТ	AAA	GAC	AGG	GGA	AAA	GGC	AAT	GGG	AÇA	стс	ATC	ACC	450
CAT	CGA	TAG	GCT	GTA	CAA	GTC	TGG	TAC	AT	ACTT	CTC	CGA	ATO	STTA	CTA	ATG	GAA	TTT	CTG	TCC	CC1	TTT	ĊCG	TTA	ccc	TGT	GAG	TAG		,55
٧	A	ı	R	н	V	Q	т	М	Y	Ε	ε	A	Y	N	D	Y	L	κ	D	R	G	κ	C	N	G	T	L	I	7	
ттс	CAC	TCA	ACA	TTT	CAC	STGC	TGI	rgg	444	ĄGAA	AGC	TCC	GA/	ACAC	GTC	CĄA	(CC)	TAC A	TGC	CCA	AAC	GAG	CTT	CTA	GGA	CAC	AAG	AAT	TGC	540
AAG	GTO	AGI	TGI	AAA	GTO	CACC	AC/	100	TTT	TCTI	TCC	AGO	CT	TGT	CAG	GTT	GG	ATGI	ACG	GGT	TTO	стс	GAA	GAT	CC1	GTG	TTC	TTA	ACG	0.0
F	н	s	τ.	F	a	С	С	G	K	Ε	s	s	Ε	Q	٧	a	Р	T	C	Ρ	κ	Ε	L	L	G	н	K	N	С	
ATC	GA ⁻	ΓGA/	AATI	rgag	ACC	CAŢA	AATO	CAG	TGT	TAA	CTO	CAC	SÇT	CAT	TGGA	ATI	GT	CGGT	[TAT	rggA	AT	rgc/	GGT	CTG	AC	ATO	TTT	GGC	ATG	630
TAG	CT	ACT	TTA	AC TO	TG	GTAI	TAC	STC	ACA	ATTO	GAC	GTO	GA	GTA	ACCI	TAA	CA	GCC	ATA	ACCT	TA	ACGI	CCA	GAC	TG	TAC	AAA	CCG	TAC	
1	D	Ε	1	Ε	T	ı	I	S	٧	K	L	a	L	1	G	1	٧	G	ï	G	ì	A	G	L	T	I	F	G	M	
ATA	TT(	CAG	CAT	GGTC	CT	CTG	TG.	TGC	GAT	ACG/	AAA	CTC	AÇG.	AGA	TGTO	SATA	ATG.	AAG	CTA	TTC	TA	CATO	AAA	ATT	GC	AATO	TAA	AGC	777	- 720
TAT	AAI	GTC	GTA(	CCAC	GGA	GAC	GAC	ACG	CTA	TGC	TTT	SAG	TGC	TCT	ACAC	TA	TAC	TTC	GAT(	GAAC	AT	GTA	ודד:	TAA	CG.	T TÁC	TTAG	TCC	AAA	
ł	F	s	М	٧	L	С	С	A	I	R	N	s	R	D	٧	ī		(	EC	ŢĢ		1	VO	1.0	25	)	)			
		GTT	-+-	<del></del>	73	4 (	S	Ģ	J	D	N	0	9													~				

AGTGTTTATGGGACTAAAAAACTTTTAACACCTTTTTAGGGGAAATATTTTGGTCCTATACAAAACATGTAAATATGCTTTATTACTTTC	90
TCACAAATACCCTGATTTTTTGAAAATTGTGGAAAAATCCCCTTTATAAAACCAGGATATGTTTTGTACATTTATACGAAATAATGAAAG	
	180
TAAAAGACTGGGACGACATTTGATGACGTTGGGAGTGTAGGGAGTTTCCCTGAAAATACAGTTTGAGAAGACAAAGAGGTTTATATTCCT	
	270
TTTTTCTGATTTCGTTCTCTAGACCGTCAACTITTAACACCCTTTCTCTTAAACATACCCGTGACATAGATACTTTATGGAGTATGAATG	
GTTTACATGTTTTCCTAACTTTTTGTATTTTCTTGTATAGCCACCTAGAGAATTCTTCATAGATTAAGAACTACAGTTTTCACCACTTA	360
CAAATGTACAAAAGGATTGAAAAAACATAAAAAAGAACATATCGGTGGATCTCTTAAGAAGTATCTAATTCTTGATGTCAAAAAGTGGTGAAT	
ACATAAGTAAAACAAAGTCCTTCATAATTTAACCATTAGCATCTTTGGCCAAACCAAAATAAAGAAAAGCATCTTCTCCTAGTTGTGTG	450
TGTATTCATTTGTTCAGGAAGTATTAAATTGGTAATCGTAGAAACCGGTTTGGTTTTATTTCTTTTCGTAGAAGAGGATCAACACACA	
GGGCAACAGAACAAGTTAAGGAAACAAAAATACTTATATATA	540
CCCGTTGTCTTTGTTCAATTCCTTTGTTTTTATGAATATATGTGTCTTGTTTTTATTACAAGAAAAATACGTTTAGGGGACACTTTTA	
AAAATTTTCAATGTTTAAAAAAAAAAAAAAAAAAAAAAA	

Page 1

HHFEK4	")
TTCGGCACGAGCTGCGGCGGTGGGCGGCTGGGCGGCCCCGGGAGCCGCGCTCTCAGTCTCTCTAGGCGCAGTCCCTTCGCCGCTTCC	GG
AAGCCGTGCTCGACGCCCGCCACCCGCCGACCCGCCGGGGCCCTCGGCGCGAGAGTCAGAGAGATCCGCGTCAGGGAAGCGGCGAAG	<del>}</del> 90
AGCCCCTGGCAGGGCCCAGAAGCCATGGCCCACTATAAGACTGAGCAGGACGACTGGCTGATCATCTACTTGAAGTATTTACTCTTT	TC 180
TCGGGGACCGTCCCGGGTCTTCGGTACCGGGTGATATTCTGACTCGTCCTGCTGACCGACTAGTAGATGAACTTCATAAATGAGAAA	AG
M A H Y K T E Q D D W L I I Y L K Y L L F	v
TTCAACTTCTTCTTCTGGGTCGGGGGAGCAGCCGTCCTGGCTGTGGGCATCTGGACCCTGGTGGAGAAGAGTGGCTACCTCAGCGTC	
AAGTTGAAGAAGAAGACCCAGCCCCCTCGTCGGCAGGACCGACACCCGTAGACCTGGGACCACCTCTTCTCACCGATGGAGTCGCAG	→ 270 AC
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	L
GCCTCCAGCACCTTTGCCGCCTCCGCCTACATCCTCATCTTTGCGGGCGTACTTGTCATGGTGACCGGCTTCCTGGGCTTCGGTGCC	TC -+ 360
CGGAGGTCGTGGAAACGGCGGAGGCGGATGTAGGAGTAGAAACGCCCGCATGAACAGTACCACTGGCCGAAGGACCCGAAGCCACGG	
A S S T F A A S A Y I L I F A G V L V M V T G F L G F G A	l
CTCTGGGAGCGGAAGGGCTGCCTCTCCACGTATTTCTGCCTGTTGCTCGTCATCTTCCTGGTTGAGCTGGTGGCGGGAGTCCTGGCC	AT + 450
GAGACCCTCGCCTTCCCGACGGAGAGGTGCATAAAGACGGACAACGAGCAGTAGAAGGACCAACTCGACCACCGCCCTCAGGACCGG	TA
LWERKGCLSTYFCLLLVIFLVELVAGV _. LA	Н
GTGTATTACCAGAGGCTGAGTGATGAACTGAAGCAGCACTTGAACCGGACTCTGGCTGAGAACTACGGGCAGCCGGAGCACGCAGAT	<del>+</del> 540
CACATAATGGTCTCCGACTCACTACTTGACTTCGTCGTGAACTTGGCCTGAGACCGACTCTTGATGCCCGTCGGCCTCGTGCGTCTA(	TG
V Y Y Q R L S D E L K Q H L N R T L A E. N Y G Q P, E H A D	н
GCCTCAGTGGACCGACTCCAGCAGGATTTCAAGTGCTGCGGAAGCAACAGCTCAGCCGACTGGCAGCACAGCACGTACATCCTGTTG	<del>-+</del> 630
CGGAGTCACCTGGCTGAGGTCGTCCTAAAGTTCACGACGCCTTCGTTGTCGAGTCGGCTGACCGTCGTGTCGTGCATGTAGGACAAC	CC
A S V D R L Q Q D F K C C G S N S S A D W Q H S T Y I L L	R
GAGGCCGAGGGCCGCCAGGTGCCCGACAGCTGCTGCAAGACAGTGGTGGCGCGCTGCGGCCAGCGGGCCCACCCCTCCAACATCTAT	<del>+</del> 720
CTCCGGCTCCCGGCGGTCCACGGGCTGTCGACGACGTTCTGTCACCACCGCGCGACGCCGGTCGCCCGGGTGGGGAGGTTGTAGATA	TC
E A E G R Q V P D S C C K T V V A R C G Q R A H P S N I Y	K
GTGGAGGGAGGCTGCCTCACCAAGCTGGAGCAGTTCCTGGCCGACCACCTGCTGCTTATGGGGGCAGTGGGCATCGGGGTGGCCTGC	TG 810
CACCTCCCTCCGACGGAGTGGTTCGACCTCGTCAAGGACCGGCTGGTGGACGACGAATACCCCCGTCACCCGTAGCCCCACCGGACG	
V E G G C L T K L E Q F L A D H L L L M G A V G I G V A C	L
CAGATCTGCGGGATGGTTCTCACCTGCTGCTTGCACCAGAGGCTCCAGCGGCATTTTTACTAATGGCAACCACCTCCTCTTCCAACT	.cc → 900
GTCTAGACGCCCTACCAAGAGTGGACGACGACGTGGTCTCCGAGGTCGCCGTAAAAATGATTACCGTTGGTGGAGGAGAAGGTTGA	ann
Q I C G M V L T C C L H Q R L Q R H F Y . (SCO.TD NO: 26	

HHFEKHO

Page 2

CCTCAAGACAACATGTGGCACATGCCATCTGCAAGG
GGAGTTCTGTTGTACACCGTGTACGGTAGGCGTTCC

936 SCO IDNO: 11

3NSDOCID: <WO___9831799A2_I_>

																		_						H	G	Be	۷'	89	L	
AGC	TTA	CTT	TCA	CTC	ACC	SCC	TGT	сст	TCC	TGA	CAC	CTC.	ACC	ΑTG	TGT.	ACG	GGA	AAA	TGT	GCC	CGC	TGT	GTG	GGG	стс	TCC	стс	ATT.		90
TCG.	AAT	GAA	AGT	GAG	TGG	ĊĠĠ	ACA	GGA	AGG	ACT	GTG	GAG	TGG	TAC	ACA	TGC	ССТ	TTT	ACA	CGG	GCG	ACA	CAC	ccc	GAG	AGG	GAG	TAA		50
				Ŋ,	-									M	С	T	G	K	С	A	R	С	٧	G	L	s	L	I	τ	
стс	TGC	стс	GTC	TGC	ATT	GTG	GCC	AAC	GCC	CTC	CTG	CTG	GTA	CCT	AAT	GGG	GAG	ACC	TCC	TGG	ACC	AAC	ACC	AAC	CAT	стс	AGC	TTG	CAA	180
GAG.	ACG	GAG	CAG	ACG	TAA	CAC	CGG	TTG	CGG	GAG	GAC	GAC	CAT	GGA	TTA	ССС	стс	TGG	AGG	ACC	TGG	TTG	TGG	TTG	GTA	GAG	TCG	AAC	GTT	
L	С	L	٧	С	ī	٧	A	N	A	L	L	L	٧	Р	N	G	Ε	Ţ	s	W	T	N	T	N	Н	L	s	L	۵	
GTC	TGG	стс	ATG	GGC	GGC.	ттс	ATT	GGC	GGG	GGC	СТА	ATG	GTA	CTG	TGT	CÇA	GGG	ATT	GCA	GCC	GTT	CGG	ĢCA	GGG	GGC	:AAG	GGC	TGC	TGŢ	
CAG	ACC	GAG	TAC	CCG	CCG	AAG	TAA	CCG	CCC	CCG	GAT	TAC	CAT	GAC	ACA	GGT	ССС	TAA	CGT	CGG	CAA	GCC	CGT	ССС	CCG	TTC	ccg	ACG.	ACA	2/0
٧	W	L	М	G	G	F	Į	G	G	G	L	М	٧	L	С	Р	G	i	A	A	٧	R	A	G	G	ĸ	G	С	С	
GGT	GCT	GGG	TGC	TGT	GGA	AAC	CGC	TGC	AGG	SATG	CTG	CGC	TCG	GTC	TTC	тсс	TCG	GCG	TTC	GGG	GTG	CTT	GGT	GCC	ATC	TAC	TG			
CCA	CGA	CCC	ACG	ACA	сст	TTG	GCG	ACG	TCC	TAC	GAC	GCG	AGC	CAG	AAG	AGG	AGC	CGC	AAC	CCC	CAC	GAA	CCA	cgg	TAG	ATG	ACG	e.		
G	A	G	С	С	G	N	R	С	R	M	L	R	s	٧	F	S	S	Α	F	G	٧	L	G	Α	ŧ	Y	С	L		
GTG	TCT	GGA	GCT	GGG	CTC	CGA	AAT	GGA	ccc	AGA	TGC	TTA	ATG	AAC	GGC	GAG	TGG	GGC	TAC	CAC	TTC	GAA	GAC	ACC	GCG	GGA	GCT	TAC	TTG	
CAC	AGA	CCT	CGA	CCC	GAG	GC T	TTA	CCT	GGG	TCT	ACG	AAT	TAC	TTG	CCG	CTC	ACC	CCG	ATO	GTG	AAG	CTT	CTG	TGG	CGC	:ccT	CGA	ATG		450
٧	s	G	A	G	L	R	N	G	Р	R	С	L	M	N	G	ε	W	G	Y	Н	F	Ε	D	т	A	G	A	Y	L	
стс	AAC	cgc	ACT	СТА	TGG	GAT	CGG	TGC	GAG	GCG	ссс	CCT	CGC	GTG	GTC	cçc	TGG	AAT	GTO	ACG	сто	:110	TCG	CTG	CTG	GTG	GCC	GCC	тсс	5"0
GAG	TTG	GCG	TGA	GAT	ACC	CTA	GCC	ACG	СТ	CGC	GGG	GGA	GCG	CAC	CAG	GGG	ACC	TTA	CAC	TGC	GAG	AAG	AGC	GAC	GAC	CAC	CGG	CGG	AGG	540
L	N	R	т	L	W	D	R	С	Ε	A	Ρ	Р	R	٧	٧	P	W	N	V	Ţ	L	F	S	L	L	ν.	A	A	s	
TGC	CTG	GAG	ATA	GTA	CTG	TGT	GGG	ATC	CAC	SCTG	GTG	AAC	GCG	ACC	ATT	GGT	GTC	TTC	TGO	GGC	GAT	TGC	AGG	AAA	AAA	LCAG	GAC	ACA	CCŢ	
ACG	GAC	CTC	TAT	CAT	GAC	AC A	ССС	TAG	GTO	GAC	CAC	TTG	CGC	TGC	TAA	CCA	CAG	AAC	ACC	CCG	CTA	ACG	TCC	TTT	TTI	rgtc	CTG	TGT	GGA	630
С	L	Ε	i	٧	L	С	G	I	a	L	٧	N	A	Т	ŧ	G	٧	F	С	G	D	С	R	ĸ	κ	Q	D	т	Р	
CAC	TGA	GGC	TCC	ACT	GAC	CGC	EGG	GTT	AC/	ACCT	GCT	CCT	TCC	TGG	ACG	стс	ACT	CCC	TTO	стс	GCT	TAGA	ATA	AAC	TGC	: 111	GCG	стс	TCA	
GTG	ACT	CCG	AGG	TGA	CTG	<del>- } -</del>	GCC	CAA	TG	GGA	CGA	GGA	AGG	SACO	TGC	GAG	TGA	GGG	AAC	GAG	CGA	TCT	TAT	TTG	ACC	AAA	CGC	GAG	AGT	720
н	٠ (	(5	eG	) J	D	Ν	0:	2	7	)																				
										ි ඛ්	Tí	1 C	SC	) .	12															
							-			_ `			. –	•		_														

Page 1

GGCACGAGAGATTGTCGGCTGCGGGTATATTCCAATTCCCCGTCTCCTCATGAATATGAAGTGAAGGGCTCTGACCCTGGAAGTGGT	TCT + 90
CCGTGCTCTCTAACAGCCGACGCCCATATAAGGTTAAGGGGCAGAGGAGTACTTATACTTCACTTCCCGAGACTGGGACCTTCACCA	
AAGCAGGGCAAAATGGGGTCTCGGAAGTGTGGAGGCTGCCTAAGTTGTTTGCTGATTCCGCTTGCACTTTGGAGTATAATCGTGAAC	
TTCGTCCCGTTTTACCCCAGAGCCTTCACACCTCCGACGGATTCAACAAACGACTAAGGCGAACGTGAAACCTCATATTAGCACTTG	TAT
M G S R K C G G C L S C L L I P L A L W S I I V N	ī
TTATTGTATTTCCCGAATGGGCAAACTTCCTATGCATCCAGCAATAAACTCACCAACTACGTGTGGTATTTTGAAGGAATCTGTTTC	TCA
AATAACATAAAGGGCTTACCCGTTTGAAGGATACGTAGGTCGTTATTTGAGTGGTTGATGCACACCATAAAACTTCCTTAGACAAAG	<del></del>
AATAACATAAAGGGCTTACCCGTTTGAAGGAAGAAGAAGAAGAAGAAGAAGAAGAAGAAGA	
LLYFPNGQTSYASSNKLINYVWTFEGICF	S
GGCATCATGATGCTTATAGTAACAACAGTTCTTCTGGTACTGGAGAATAATAACAACTATAAATGTTGCCAGAGTGAAAACTGCAGC	AAA + 360
CCGTAGTACTACGAATATCATTGTTGTCAAGAAGACCATGACCTCTTATTATTGTTGATATTTACAACGGTCTCACTTTTGACGTCG	TTT
CIMMII VITVII VIFNNNNY KCCQSENCS	K
AAATATGTGACACTGCTGTCAATTATCTTTTCTTCCCTCGGAATTGCTTTTTCTGGATACTGCCTGGGTCATCTCTGCCTTGGGTCTT	430
TTTATACACTGTGACGACAGTTAATAGAAAAGAAGGGGAGCCTTAACGAAAAAGACCTATGACGGACCAGTAGAGACGGAACCCAGAA	CAG
K Y V T L L S I I F S S L G I A F S G Y C L V I S A L G L	٧
CAAGGGCCATATTGCCGCACCCTTGATGGCTGGGAGTATGCTTTTGAAGGCACTGCTGGACGTTTCCTTACAGATTCTAGCATATGC	ATT
GTTCCCGGTATAACGGCGTGGGAACTACCGACCCTCATACGAAAACTTCCGTGACGACCTGCAAAGGAATGTCTAAGATCGTATACG	770
Q G P Y C R T L D G W E Y A F E G T A G R F L I D S S I W	I
CAGTGCCTGGAACCTGCACATGTTGTGGAGTGGAACATCATTTTATTTTCCATTCTCATAACCCTCAGTGGGCTTCAAGTGATCATC	TGC+ 630
GTCACGGACCTTGGACGTGTACAACACCTCACCTTGTAGTAAAATAAAAGGTAAGAGTATTGGGAGTCACCCGAAGTTCACTAGTAC	
OCIEPA H V V F W N I I L F S I L ! T L S G L Q V I I	С
CTCATCAGAGTAGTCATGCAACTATCCAAGATACTGTGTGGAAGCTATTCAGTGATCTTCCAGCCTGGAATCATTTGAATAAGGAC	
GAGTAGTCTCATCAGTACGTTGATAGGTTCTATGACACACCTTCGATAAGTCACTAGAAGGTCGGACCTTAGTAAACTTATTCCTG	
LIRVVMALSKILCGSYSVIFAPGII.(SCOI)	> 1,0 ! é
TGTTTTCCATTATCAAGACATGGCCATCTATCTAAATATTATATCAACTGTGTTAGACTTGAGGGCAATATTGAAAAATGATGGTGC	TTC
ACAAAAGGTAATAGTTCTGTACCGGTAGATAGATTTATAATATAGTTGACACAATCTGAACTCCCGTTATAACTTTTACTACCACG.	
TGCATTTGGTGTTTATTTGTAAAAAATTTGCAGTCCTCACTGCACATGCAAGTATACCACCCTTCCATTTAGTATGTTTTTTAAGT	AATA
ACGTAAACCACAAATAAACATTTTTTTAAACGTCAGGAGTGACGTGTACGTTCATATGGTGGGAAGGTAAATCATACAAAAAATTCA	<del></del>

HUVBB80

AGAAACCTTTATGCAATATGTATATTGCAACATTATTTAATATTCTGGAAAATTGGAAACCCCCAAAATTCTAACCTCAAA

TCTTTGGAAATACGTTATACATATAACGTTGTAATAAAATTATAAGACCTTTTAACCTTTGTGGGGTTTTAAGATTGAGTTT

1071

HJACE54

Page 1

TTTG	TGGA	GGC	AGC	AGA	GAG	TAC	CCA	GCT	GGA	CAT	ССТ	TTC	CTG	CTG	ATG	AGC	ccċ	AGG	CTG	GAG	GTG	ccc	TGC	TCA	CAT	GCT	стт	ccc	90
AAAC	ACCT	ccs	TCG	TCT	ctc	ATG	GGT	CGA	CCT	GŤA	GGA	AAG	GAC	GAC	TAC	TCG	GGG	TCC	GAC	стс	CAC	GGG	ACG	AGT	GTA	CGA	GAA		30
			Ţ												М	s	Р	R	L	Ε	٧	Ρ	С	s	н	Α	L	Р	
			`																										
	GTCT	+			-+-		+	$\longrightarrow$				+			<del></del>		+	$\overline{}$		-+-		+			<del>+-</del>		+	$-\!\!\!+$	180
GTCC	CAGA	SAGC	GGA	CCC	GTC	CAG	TAG	STAT	CAT	GCC	CCT	GAC	CAG	AAC	GIT	CTC	GGC	TTC	GTA	AAA	TGA	CAC	TCG	GAC	TCC	CTG	GTC	CGA	
0	G L	S	Р	G	Q	٧	i	1	٧	R	G	L	٧	L	Q	Ε	Ρ	K	н	F	T	٧	s	L	R	D	a	A	
GCCC	ATGC	COT	GIG	ACA	.c.r.c	AGG	GCC	TCC	TTC	GCA	GAC	AGA	ACT	CTG	GCC	TGG	ATC	TCC	CGC	TGG	GGG	CAG	AAG	AAA	CTG	ATC	TCA	GCC	
	TACGA	<u> </u>			<del></del> -			+				+					+	+				+			<del></del>		+	+	270
cada	TAC GA	.00/	· • · · · ·		<b>U</b> // <b>U</b>		-																						
Α	н а	Р	٧	T	L	R	Α	S	F	Α	D	R	T	L	Α	W	Ī	S	R	W	G	Q	ĸ	K	L	I	S	Α	
ссст	тсст	TTT	TAC	CCC	CAG	AGA	TTC	TIT	GAG	GTG	CTG	ĊТС	СТĢ	TTC	CAG	GAG	GGA	GGG	CTG	AAG	CTG	GCG	CTC	AAT	GGG	CAG	GGG	CTG	360
GGGA	AGGA	SAAA	ATG	GGG	GTC	TCT	AAG	SAAA	стс	CAC	GAC	GAG	GAC	AAG	GTC	CTC	ССТ	ССС	GAC	TTC	GAC	cgc	GAG	TTA	CCC	GTC	ccc	GAC	300
Р	F L	F	Υ	Р	Q	R	F	F	Ε	٧	L	Ł	L	F	a	Ε	G	G	L	K	L	А	L	N	G	a	G	L	
•					_																• • •	- <del>-</del> -							
	CCAC	+			$\rightarrow$			+							-+-			<del></del> +		$\rightarrow$		+	+				+		450
cccc	GGTG	STCG	TAC	TTG	GTC	GTC	CGG	GGAC	СТС	GTC	GAC	GCC	CTC	GAG	GCC	TAG	TCA	CCT	TCA	CAG	GIC	GAG	AIG	ACA	CAG	GIG	AGG (	SC	OI
G	A T	s	M	N	a	Q	Α	L	Ε	a	L	R	Ε	L	R	ı	S	G	S	٧	0	L	Y	С	٧	н	S	,	NO:2
AGGA	TGGT	ГССА	.GGA	AAT	ACC	GCA	GAA	AAAC	AAG	AGT	CAG	CCA	CTC	ccc	AGG	GCC	CCA	стс	TCC	TCC	сст	CAT	TAA	ACÇ	ATC	CAC	CTG	AAÇ	
	ACCA						+					+			$\overline{}$		+	+		-+		+					<del></del>		540
ACCA	GCAC	ATCA	GGG	сст	GGT	TCA	CCT	rctg	GGG	TCA	CGA	GAC	TGA	GTC	TAC	AGG	AGC	TTT	GGG	CCT	GAG	GGA	AGG	CAC	AAG	AGT	GCA	AAĢ	
TGGT	CGTG	ragt	CCC	GGA	CCA	AGT	GGA	AGAC	CCC	AGT	GCT	CTG	ACT	CAG	ATO	STCC	TCG	AAA	CCC	GGA	стс	CCT	TCC	GTG	TTC	TCA	CGT		630
																	٠												
GTTC	CTCG	AACT	сто	CAC	стт	ССТ	CCA	ACCA	GGA	GCC	TGG	GAT	ATG	GCT	CCA	TCT	GCC	TTC	AGG	GCC	TGG	ACT	GCA	CTC	AÇA	GAG	GCA	AGŢ	700
CAAG	GAGC	<del>-i-</del> ΓΤGΑ	GAC	GTG	GAA	GGA	GGT	rggt	CCT	CGG	ACC	CTA	TAC	CGA	GGT	AGA	CGG	AAG	TCC	CGG	ACC	TGA	CGT	GAG	TGT	стс	CGT		720
GTTG	TAGA	CTAA	CAA	AGA	TAC	TCC	AAA	ATA	CAA	TGG	CTT	AAA	GAA	TGT	GGT	CAT	TTA	TTC	ттт	ATT	ATT	ŢAT	TTA	TTT	GTG	GTC	AAA	TAĄ	010
CAAC	ATCT	SATT	GII	TCT	ATG	AGG	111	TAT	GTT	ACC	GAA	<del> </del>	CTT	ACA	CCA	GTA	AAT	AAG	AAA	TAA	TAA	ATA	AAT	AAA	CAC	CAG	TTT		810
ATAA	ATAA	GGTT	ATT	TAT	TTA	AAA	AAA	<b>AAA</b>	AAA	AAA	AAA	AAA	AAA	AAA	AAA	AAA	À		1-	`aC-	. 7	^	N iz	<b>つ</b> ・	14	1			
TATT	TATT	CAA	TAA	ATA	AAT	TTT	TTI	1111	ттт	TTT	111	111	TTT	TTT	T T 1	TTT	► 8 T	65		eG	L	-17	110	ノ·	• '	• J			

HROAD63

Päge 1

GCACGAGAGACGACATCAGAGATGAGGACAGCATTGCTGCTCCTTGCAGCCCTGGCTGTGGCTACAGGGCCAGGCCCTTACCCTGCGCTGC

CGTGCTCTCTGCTGTAGTCTCTACTCCTGTCGTAACGACGAGGAACGTCGGGACCGACACCGATGTCCCGGTCGGGAATGGGACGCGACG

MRTALLLLAAALAVATGPAALTTLRC

CACGTGTGCACCAGCTCCAGCAACTGCAAGCATTCTGTGGTCTGCCCGGCCAGGCCTCTCGCTTCTGCAAGACCACGAACACAGTGGAGCCT

GTGCACCACGTGGTCGAGGTCGTTGACGTTCGTAAGACACCAGACCAGACCAGAGCCTCTCGCTTCTGCAAGACCACGAACACAGTGGAGCCT

GTGCACCACGTGGTCGAGGTCGTTGACGTTCGTAAGACACCAGACCAGAGCGCCGGAGAGACGTTCTGGTGCTTGTGTCACCTCGGA

HVCTSSSNCKHSVVCPAASSTCTGGGACCAGGTCCAGGTGGGCATGGAATGCTGATGACTTGGAGCAGGCCCCACAGACCCCACAGAGGAT

CTGAGGGCTTCCCCGAAAGTCTGGGACCAGGTCCAGGTGGGCATGGAATGCTGATGACTTGGAGCAGGCCCCACAGACCCCACAGAGGAT

GACTCCCGAAGGGGCTTTCAGACCCTGGTCCAGGTCCACCCGTACCTTACGACCTCGACCTGGAGCCTCCGGGGGTGTCTGGGGGTGTCTCCTA

LRASPKVWDQVGMEC.CCGTACCTTACGACCTGGACCTCGGGGGTGTCTGGGGGTGTCTCCTA

GAAGCCACCCCACAGAGGATGCAGCCCCCAGCTGCATGGAAGGTGGAGGACAGAAGCCCTGTGGATCCCCGGATTTCACACTCCTTCTGT

CTTCGGTGGGGTGTCTCCTACGTCGGGGGTCGACGTACCTTCCACCTCCTGTCTTCGGGACACCTAGGGGCCTAAAGTGTGAGGAAGACA

HMW6546

Page 1

AGCGGGCCCGAACCCTCGTGTGAAGGGTGCAGTACCTAAGCCGGAGCGGGGTAGAGGCGGGCCGGCACCCCCTTCTGACCTC	CAGTGCCG	- 90
TCGCCCGGGCTTGGGAGCACACTTCCCACGTCATGGATTCGGCCTCGCCCCATCTCCGCCCGGCCGTGGGGGAAGACTGGAC	GTCACGGC	-
CCGGCCTCAAGATCAGACATGGCCCAGAACTTGAAGGACTTGGCGGGACGGCTGCCCGCGGGCCCCGGGGCATGGGCACGG	CCCTGAAG	. 18
GGCCGGAGTTCTAGTCTGTACCGGGTCTTGAACTTCCTGAACCGCCCTGCCGACGGGCGGCCCGGGGCCCCGTACCCGTGC	GGGACTTC	
M A Q N L K D L A G R L P A G P R G M G T	A L K	
CTGTTGCTGGGGGCCGGCGCCGTGGCCTACGGTGTGCGCGAATCTGTGTTCACCGTGGAAGGCGGGCACAGAGCCATCTTCT	TTCAATCGG	
GACAACGACCCCCGGCCGCGCACCGGATGCCACACGCGCTTAGACACAAGTGGCACCTTCCGCCCGTGTCTCGGTAGAAGA	AGTTAGCC	27
L L G A G A V A Y G V R E S V F T V E G G H R A I F	F N R	
ATCGGTGGAGTGCAGCAGGACACTATCCTGGCCGAGGGCCTTCACTTCAGGATCCCTTGGTTCCAGTACCCCATTATCTATC	GACATTCGG	
TAGCCACCTCACGTCGTCCTGTGATAGGACCGGCTCCCGGAAGTGAAGTCCTAGGGAACCAAGGTCATGGGGTAATAGATAC	TGTAAGCC	• 36
I G G V O Q D T I L A E G L H F R I P W F O Y P I I Y	D I R	
GCCAGACCTCGAAAAATCTCCTCCCCTACAGGCTCCAAAGACCTACAGATGGTGAATATCTCCCTGCGAGTGTTGTCTCGAG	CCAATGCT	
CGGTCTGGAGCTTTTTAGAGGAGGGGATGTCCGAGGTTTCTGGATGTCTACCACTTATAGAGGGACGCTCACAACAGAGCTC	<del></del>	45
ARPRKISSPIGSKOLOMVNISLRVLSR	PNA	
CAGGAGCTTCCTAGCATGTACCAGCGCCTAGGGCTGGACTACGAGGAACGAGTGTTGCCGTCCATTGTCAACGAGGTGCTCA	<del>+</del>	- 54
GTCCTCGAAGGATCGTACATGGTCGCGGATCCCGACCTGATGCTCCTTGCTCACACGGCAGGTAACAGTTGCTCCACGAGT	TETEALAL	
Q E L P S M Y Q R L G L D Y E E R V L P S I V N E V L	K S V	
GTGGCCAAGTTCAATGCCTCACAGCTGATCACCCAGCGGGCCCAGGTATCCCTGTTGATCCGCCGGGAGCTGACAGAGAGG	CCAAGGAC	63
CACCGGTTCAAGTTACGGAGTGTCGACTAGTGGGTCGCCCGGGTCCATAGGGACAACTAGGCGGCCCTCGACTGTCTCTCCC	GGTTCCTG	
V A K F N A S Q L I T Q R A Q V S L L ! R R E L T E R	A K D	
TTCAGCCTCATCCTGGATGATGTGGCCATCACAGAGCTGAGCTTTAGCCGAGAGTACACAGCTGCTGTAGAAGCCAAACAAC	STGGCCCAG	:
AAGTCGGAGTAGGACCTACTACACCGGTAGTGTCTCGACTCGAAATCGGCTCTCATGTGTCGACGACATCTTCGGTTTGTTC	<del></del>	· 72
	v A O	
FSEILDUVALIELSFSKETIAAVEAKU	V A U	
CAGGAGGCCCAGCGGGCCCAATTCTTGGTAGAAAAAGCAAAGCAGGGAACAGCGGCAGAAAATTGTGCAGGCCGAGGGTGAGG	CCGAGGCT	. 81
GTCCTCCGGGTCGCCCGGGTTAAGAACCATCTTTTTCGTTTCGTCCTTGTCGCCGTCTTTTAACACGTCCGGCTCCCACTC	GGCTCCGA	
Q E A Q R A Q F L V E K A K Q E Q R Q K ! V Q A E G E	A E A	
GCCAAGATGCTTGGAGAAGCACTGAGCAAGAACCCTGGCTACATCAAACTTCGCAAGATTCGAGCAGCCCAGAATATCTCCA	AAGACGATC	. 0/
CGGTTCTACGAACCTCTTCGTGACTCGTTCTTGGGACCGATGTAGTTTGAAGCGTTCTAAGCTCGTCGGGTCTTATAGAGG		
	K T I	

HMWG546

^oage 2

GCCACATCACAGAATCGTATCTATCTCACAGCTGACAACCTTGTGCTGAACCTACAGGATGAAAGTTTCACCAGGGGAAGTGACAGCCTC
CGGTGTAGTGTCTTAGCATAGATAGATGTGTCGACTGTTGGAACACGACTTGGATGTCCTACTTTCAAAGTGGTCCCCTTCACTGTCGGAG

A T S Q N R I Y L T A D N L V L N L Q D E S F T R G S D S L

ATCAAGGGTAAGAAATGAGCCTAGTCACCAAGAACTCCACCCCCACAAGAAGTGGATCTGCTTCTCCAGTTTTTGA
TAGTTCCCATTCTTTACTCGGATCAGTGGTTCTTGAGGTGGGGGTGTTCTTCACCTAGACGAAGAGGTCAAAAACT

IKGKK (SCQIDNO: 31)

HIF. WOL

Page 1

GCA	CGA	GAT	GAC	ATC	ACT	AAG	TGG	ccc	SATE	TGC	ACA	GAG	CAG	GCC	AGG	AGC	AAC	CAC	AC A	GGC	110	CTO	CAC	ATG	GAC	TGC	GAG:	ATC	AA	90
CGT	GCT	CTA	CTG	TAG	TGA	T.T.C	CACC	GGC	TAG	ACG	TGT	CTC	GTC	CGG	STCC	TCG	TTG	GTG	TGT	CCG	AAG	GAC	GTG	TAC	CTG	ACG	СТС	TAG	TT	
				, v																				M	D	С	Ε	I	K	
GGG	CGC	ccç	TGC	TGC	ATC	ĢGC	ACC	AAG							ACC				TĢT	GAG	TTC	ATO	CAC	GGC	TAT	TTC	CAT	GAG	GA	•
CCG	GCG	GGG	ACG	ACG	TAG	ccc	TGG	TTC	ccc	TCG	ACA	стс	TAG	TGC	TGG	GCC	стт	ATG	ACA	CTC	AAG	TAC	GTG	CCG	ATA	AAG	GTA	стс	СТ	18
G	R	Р	С	С	1	G	Т	κ	G	s	С	Ε	I	Ţ	T	R	Ε	Y	C	Ε	F	M	н	G	Y	F	н	E	Ε	
AGCA	ACA	CTO	TGC	TCC	CAG	GTG	SAGG	CG/	AGGC	AGG	CCT	GGA	GTA	GTO	GAG	GAG	AGG	ACG	CTG	GGC	ATG	GCA	GCC	TGC	TGG	GGC	CGG	GGC	TC	
CGT	TGT	GAG	ACG	AGG	GTC	CAC	TCC	GCT	CCG	TCC	GGA	CCT	CAT	CAC	стс	CTC	TCC	TGC	GAC	CCG	TAC	CGI	CGG	ACG	ACC	CCG	GCC	CCG	AG	2
A	T	L	С	s	a	٧	R	R	G	R	Р	G	٧	٧	E	Ε	R	T	L	G	М	A	A	С	W	G	R	G	s	
	+					+			-+-		+				GGC	+			<del></del>		+					+			-+	3
GCG	TGA	GGG	AGG	GTA	CAG	CCI	CGG	AGI	CTG	AGT	CCG	ACG	AAG	ACC	CCG	CGA	CTC	GTG	GTA	TAC	GGG	TAA	GGG	TCC	ACG	TGA	CAA	AAC	CT	
R	Ŧ	Ρ	\$	Н	٧	G	A	S	D	S	G	С	F	W	G	A	Ε	Н	Н	М	P	I	Ρ	R	С	T	٧	L	Đ	
CAAG	GTG	TGT	TGG	GCT	GCT	ĢCC	: T T C	сто	AĄC	сст	GAG	GTC	CCA	GA1	CAG	ŢŢŢ	TAC	AGG	тст	GGC	TGT	СТС	TTT	тсс	TAC	ĄTG	TTG	GGT	AA	"
TTC	CAC	ACA	ACC	CGA	CGA	ccc	AAG	GAC	STTG	GGA	CTC	CAG	GGT	CTA	AGTC	AAA	ATG	TCC	AGA	CCG	ACA	GAG	AAA	AGG	ATG	TAC	AAC	CCA	TT	4
K	٧	С	W	A	A	Α	F	L	N	Р	Ε	٧	P	D	Q	F	Y	R	s	G	С	L	F	s	Y	м	L	G	κ	
GAGG	TCC	TCA	ATG	ccc	CCG	AAC	cce	SACO	CCT	GTG	ATG	GAC	ACC	CAC	SGEG	GAC	ccc	TGG	GGA	AAG	GTT	CCI	GGG	CCA	GGG	TAT	GGT	CGG	TC	
TCC	AGG	AGT	TAC	GGG	GGC	TTG	GGC	TGC	GGA	CAC	TAC	CTG	TGG	GTO	CGC	CTG	GGG	ACC	CCT	TTC	CAA	GG A		GGT	CCC	ATA	CCA	GCC	AG	5
R	s	s	м	Ρ	Р	N	Р	т	Р	٧	м	D	T	a	A	0	Р	W	G	K	٧	Ρ	G	Р	G	Υ	G	R	s	
	CTG	CCG	AAG	ACT	ACT	GCT	rcct	rga/	GTG	тст	GGA	TGA	AGG	cco	SCTG	сст	GGT	GTG	TCC	CTC	ccc	CAG	TGT	GGG	TGC	ACT	GCC	стс	GG	
	+					+			<del></del>		<del></del>	+			GAC	+			-+-		+					+	+		-+	6
N		0	v	т	т	٨	P	_	v	c	c				IC				_			٠								
14	Ĺ	r	٨	,	'	^	F	E-	٧	J	u							_	•						,	/ Ç	~	I!	` 1	VI/
TGTC	TTG	TGG	GTC	TIT	TCA	AAT	GAC	ATO	CCT	GAA	GGG	GAC	CTG	GAG	GAA	GGT	GGT	CCG	GCT	GGC	ACC	CTA	TCG	:C → 7	'04	ـدـ	9	٠.	١	Ac

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 13 , line Table 1										
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet									
Name of depositary institution  American Type Culture Collection										
Address of depositary institution (including postal code and count	אמ									
12301 Parklawn Drive Rockville, Maryland 20852 United States of America										
Date of deposit May 16, 1997	Accession Number 209053									
C. ADDITIONAL INDICATIONS (leave blank if not applicable	e) This information is continued on an additional sheet									
D. DESIGNATED STATES FOR WHICH INDICATIONS A	RE MADE (if the indications are not for all designated States)									
E. SEPARATE FURNISHING OF INDICATIONS (leave blank)										
The indications listed below will be submitted to the International   Number of Deposit')	Bureau later (specify the general nature of the indications e.g., "Accession									
For receiving Office use only	For International Bureau use only									
This sheet was received with the international application	This sheet was received by the International Bureau on:									
Authorized officer Yvstts Sinning	Authorized officer									
Paralegal Specialis: IAPD-PCT Operation:										
IN OF OTAC										

Form PCT/RO/134 (July 1992)

#### INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

REC'D 23 FER 1998 (PCT Rule 13bis) POT WIPO A. The indications made below relate to the microorganism referred to in the description on page Table 1 **B. IDENTIFICATION OF DEPOSIT** Further deposits are identified on an additional sheet Name of depositary institution American Type Culture Collection Address of depositary institution (including postal code and country) 12301 Parklawn Drive Rockville, Maryland 20852 United States of America Accession Number Date of deposit May 16, 1997 209053 This information is continued on an additional sheet C. ADDITIONAL INDICATIONS (leave blank if not applicable) EUROPE In respect of those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused of withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC). D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") For International Bureau use only For receiving Office use only This sheet was received by the International Bureau on: This sheet was received with the international application Authorized office Notes Authorized officer Particuel Specialist เครียะคือT Operations

Form PCT/RO/134 (July 1992)

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

and the state of t	
A. The indications made below relate to the microorganism refer on page 13 ,/ih// Tab	
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet
Name of depositary institution  American Type Culture Collection	
Address of depositary institution (including postal code and coun 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	try)
Date of deposit  May 16, 1997	Accession Number 209054
C. ADDITIONAL INDICATIONS (leave blank if not applicab	le) This information is continued on an additional sheet
D. DESIGNATED STATES FOR WHICH INDICATIONS A	RE MADE (if the indications are not for all designated States)
E. SEPARATE FURNISHING OF INDICATIONS (leave bla	nk if not applicable)
The indications listed below will be submitted to the International Number of Deposit")	Bureau later (specify the general nature of the indications e.g., "Accession
For receiving Office use only	For International Bureau use only
This sheet was received with the international application	This sheet was received by the International Bureau on:
Authorized officer Control Specialist Farebook Specialist FARD-PCT Operations	Authorized officer

Form PCT/RO/134 (July 1992)

# INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

REC'D	2 3	FEB	1998
WIPO	PC	T	

A. The indications made below relate to the microorgan	rism referred to in the description				
on page, <del>line</del> ,					
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet				
Name of depositary institution					
American Type Culture Collection					
Address of depositary institution (including postal code and country)  12301 Parklawn Drive  Rockville, Maryland 20852  United States of America					
	The second secon				
Date of deposit May 16, 1997	Accession Number 209054				
C. ADDITIONAL INDICATIONS (leave blank if not	applicable) This information is continued on an additional sheet				
application has been refused or wi by the issue of such a sample to a the sample (Rule 28(4) EPC).	iropean patent or until the date on which thdrawn or is deemed to be withdrawn, only an expert nominated by the person requesting				
E. SEPARATE FURNISHING OF INDICATIONS (I	leave blank if not applicable) national Bureau later (specify the general nature of the indications e.g., "Accession				
Number of Deposit")	iational Bureau later (specify the general nature of the translations e.g., Accession				
For receiving Office use only	For International Bureau use only				
This sheet was received with the international applic	11,				
Authorized officer  Water Smiths  Haralegal Specialist  IAPD-PCT Operations	Authorized officer				

10

15

20

25

35

#### What is claimed is:

- 1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the receptor polypeptide of SEQ ID NO:Y; or a nucleotide sequence complementary to said isolated polynucleotide.
- 2. The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:X encoding the receptor polypeptide of SEQ ID NO:Y.
- 3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80% identical to that of SEQ ID NO:X over its entire length.
- 4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: X.
  - 5. The polynucleotide of claim 1 which is DNA or RNA.
- 6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a receptor polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:Y when said expression system is present in a compatible host cell.
  - 7. A host cell comprising the expression system of claim 6.
- 8. A process for producing a receptor polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
  - 9. A process for producing a cell which produces a receptor polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a receptor polypeptide.
  - 10. A receptor polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:Y over its entire length.

10

25

- 11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:Y.
  - 12. An antibody immunospecific for the receptor polypeptide of claim 10.
- 13. A method for the treatment of a subject in need of enhanced activity or expression of receptor polypeptide of claim 10 comprising:
- (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
  - (b) providing to the subject polynucleotide of claim 1 in a form so as to effect production of said receptor activity in vivo.
- 14. A method for the treatment of a subject having need to inhibit activity or expression of the receptor polypeptide of claim 10 comprising:
  - (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
  - (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
- 20 (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.
  - 15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the receptor polypeptide of claim 10 in a subject comprising:
  - (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said receptor polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the receptor polypeptide30 expression in a sample derived from said subject.
  - 16. A method for identifying agonists to the receptor polypeptide of claim 10 comprising:
    - (a) contacting cells produced by claim 9 with a candidate compound;
- 35 and
  - (b) determining whether the candidate compound effects a signal generated by activation of the receptor polypeptide.

- 17. An agonist identified by the method of claim 16.
- 18. The method for identifying antagonists to the receptor polypeptide of claim 10 comprising:
  - (a) contacting said cell produced by claim 9 with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.
  - 19. An antagonist identified by the method of claim 18.

15

5

- 20. An isolated receptor polynucleotide comprising a nucleotide sequence selected from the group consisting of:
- (a) a nucleotide sequence having at least 80% identity to a nucleotide sequence encoding the receptor polypeptide expressed by the cDNA insert deposited at the ATCC; and
- (b) a nucleotide sequence complementary to the nucleotide sequence of (a).
- 21. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a receptor polypeptide.

FIGURE 1

				1
	10 MLLILL PLLW GRERV	20 	30 VIST T M 60 S 9 W 15	40 L V C E C M HMACR70.AA
1 1	MLLILLPLENG MLPLLLBLENG	- G S L Q E K P V	YELOVOKSVI	VOEGL OB-1.aa
	50	60	70	80
41 36	CWHVRCSFSYFVDSQ7	DSDPVHGXW SSPELYVYW	RRAGNDISWK RRDGEIPYYA	APVAT HMACK/U.AA EVVAT OB-1.aa
20		100	110	120
81.	THEAN A TO E BY ROBE TO MAR PORRUK PET QGRE		C TISIRDARM	SDAGR HMACR70.AA EDITGS OB-1.aa
76	DRR NKP 取割 Q G R 和		<del></del>	160
-01	130 YEERMEKE - NIKWNY	140 KYDOBSV	150	HMACR70.AA
121 116	Y F F R M R K G - N I K W N Y Y F F R V B R G R D V K Y S Y	QONKINLEVT	ALIEKPDIHE	LEPLE OB-1.aa
	170	180	190	200 
142 156	SGRPTRLSCSLPGSC	EAGPPLTFSW	TGNALSPLDI	
	210	220	230	240
142		LTCQMKRQGA	QVTTERTVQ	- NVTYPHMACR70.AA LVVSYA OB-1.aa
196	250	260	270	280
147	250 PONLEVIV QGEGTA		SVIEGOSLE	VCAVD HMACR70.AA LCDAP OB-1.aa
236	PONLTVIV PQGEGTA	LELINOR INCL.	i v initiation	320
100	290 7 SMPRARLSWITWRSLIT	300 1 1 Y P S Q P S N P I	310 V L E L Q - V H L	GID E G E R HMACR70.AA
18° 27.		INATPLENT C	SIBBBRRM RS	
	330	340	350	360 
22 31	6 TCRAQNSLGSQHVSI 3 TCRAQHPLGFLQLFI	M T S A A S F b O I	LLGPSCSWEA	EGLHCR OB-1.aa
	370	380	390	400
24	4	EEKPLEGNS	S Q G S F K V N S S	SAGPWA OB-1.aa
J.		420	430	440
24	410 4 QQEYTGKM	98	G S O S G S V I. I. I.	OGRSNL OB-1.aa
39	4 TOQEYTGKM 3 NSSLITHGGLSSDL	<u>EVISCKAWNII</u>	G 5 Q 5 C 6 7 2 2 2	~




FIGURE 2

			,	
	10	20	30	40
1 1	MAE			SSLKKLL HIEDK48.aa SSLKLL MRC-0X44.aa
1	MGEFNEKKTTCGTVC	LKYLLFTYNC	CFWL	AGBAVM PETA-3.aa
	50	60	70	80
15 10	KYVLFFFNFLFWVC	CCTLGFGI	. <b>–</b> – на цур – –	NTYGIL MRC-OX44.aa
36			; IWTEALK	SDYISL PETA-3.aa
	90	100	110	120
29 44	LVGLGIGGKCGGASI FRNLPFLTI	, GIN V파 VIIVIGIS J	L) I M	V V FIX OX44.aa
52	LASGTY LA	AVII VVAGT	/ <u>[V M</u> ]	<u>V</u> T PETA-3.aa
	130	140	150	160
69 67	VLEGEAGWYGATEE AFLGEMGSIEE	*! **	' IT T T. T. T. L. Δ 3893 \	TITLE LIEUCEUMANIA
75	AFLECEMGSIKE	RIR NIL II R L YIF II I	<u> </u>	TAG TELA TELA 5.82
	170	180	190	200 1 DIVIN PIETVAR 33
109 10 <b>4</b>	LLFFPIVGDVALEH'	citi N DISIT ~ (	THEMHSDNS	TRMAINID MICTORALIZE
112	YAYYQQLNTELKE-		- TOPGREE	
	210	220	230	240 FEMTTGH HIEDK48.aa
146 137	LVMEKLKCCCGVMNY FIQSQLQCCGVMGS	s®owlISIGIP		MRC-OX44.aa
148	QLQQEFHCCGSMNS	OBW R DISEWIR	SQEAGGRVV	
	250 TYPRSCCKSIGSVS	260	270	280 1. 1. K T T K HIELK48.aa
17 <b>4</b> 158	491949 924142404241	CDGRDVSPNV	ADVQG - CYKI	GQAWFH MRC-OX44.aa
181	的 D S 例 C K T V V A L -	C G Q R D H A S M I	I R V Eigi Gigi I I ii	
	290	300	310	320
213 180	TQSFTLSGSSLGAA SNFLYIGIVTICVC EHLRVIGAVGIGIA	VIORWSSRYV VIOVLGMSFA	LTLNCQI	OKTSQAL MRC-OX44.aa
218	EHLRVIGAVGIGIA	C[V]関V[F]線MIFTT		
				нгелк48.аа
218	A. GL			MRC-OX44.aa PETA-3.aa
252	нү			<b>-</b>

# FIGURE 3

	10	20	30	40
1		TMMILFNLLI	FLCGAALLAVG	IWVSID HFWAE25.aa IWLAAT NAG-2.aa
1			W L G G C G V H G V E W I T G V I L H A V E	
1	M ETK PV IT C D	<u> </u>		
	50	60	70	80
36	GASFLKIFGPLS			L G F L G C HPWAF25.aa I G F V G C NAG-2.aa
38 39	QGSFATLSSSFP LGTYISLIAE-N	TNAPY	V I I G T G T T I V V	FGLFGC TALLA-1.aa
		100	110	120
	YGAKTESKCAL V		A HE WALLAND ALL V	YTTMAE HEWAE25.aa
76 73	Y G A K T E S K C A L V L G A I K E N K C L L L	TFFLLLLIVEL	LEATIAILFFA	YTDKID NAG-2.aa FRHEIK TALLA-1.aa
73	LGAIKENKCLTL FATCRGSPWMIK	L Y A M F B S E V E L	A E D VA GLI 5 GL V	
	130	140	150	160
116	HFLTLLVVPAIK	KDEGSQED	FTQVWNTTM	KGLKCC HFWAE25.aa TDFRCC NAG-2.aa
113 113	RYAQQDLK DTFLRTYT	KGLHLYGTQGN DAMQTYNGNDE	RSRAVDHVQ	R S L S C C TALLA-1.aa
				200
	170	180	190	200 IN E T T K HFWAE25.aa
151	170 170 1 CFTNYTD-FEDS	180 PYFKENSAF PF YNATRV - F	190 FCCNDNVTNTA	200 NETCTK HFWAE25.aa SESCGL NAG-2.aa
151	170	180 PYFKENSAF PF YNATRV - F	190 FEGNDNVTNTA	200 IN E T T K HFWAE25.aa
151	170 170 1 CFTNYTD-FEDS	180  PYFKENSAFPFYNATRVY BEHGL FE	190 PFCCNDNVTNTA DSCCLEF PSCMNE	200  NETCTK HFWAE25.aa SESCGL NAG-2.aa TD- NP TALLA-1.aa
151	170  170  GFTN XTD - FEDS  GVSN XTD W FEV - W STS  210  QKAHDQKVE	PYFKENSAFFFF YNATRV-FL PYFLEHGIJ-FF	190 PFCCNDNVTNTA SCCLEF SCCMNE 230	200  N E T C T K HFWAE25.aa S E S C G L NAC-2.aa T D - C N P TALLA-1.aa  240
151 149 147 190 180	170  GFTNYTD-FEDS  GVSNYTDWFEV-  GVONYTN-WSTS  210	PYFKENSAFPEYNATRV FL PYFLEHGI 220	190 PFCCNDNVTNTA SCCLEF SCCMNE 230 LYDIRTNAVTVC	200  N E T C T K HFWAE25.aa S E S C G L NAG-2.aa T D - K N P TALLA-1.aa  240  G G HFWAE25.aa G I F G L C T NAG-2.aa
151 149 147	170  GFTNYTD-FEDS GVSNYTDWFEV- GVONYTN-WSTS  210  QKAHDQKVE HAPGTWWKA	180  P Y F K E N S A F P F Y N A T R V - F L P Y F L E H G L - F F  220 G C F N Q L I P C Y E T V F V N Q K G C Y D L V	190  FCCNDNVTNTA  SCCMNE  230  YDIRTHAVTV  VWLQENLLAV  SFMETNMGIII	200  N E T C T K HFWAE25.aa S E S C G L NAG-2.aa T D - N P TALLA-1.aa  240 G G HFWAE25.aa G I F G L C T NAG-2.aa A G TALLA-1.aa
151 149 147 190 180	170  GFTNYTD-FEDS  GVSNYTDWFEV-  GVONYTN-WSTS  210  QKAHDQKVE OHAPGTWWKA QDLHNLTVAATK	180  P Y F K E N S A F P F F - Y N A T R V - P L P F L E H G L - P F P F L E H G L - P F P T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T V F T T T T	190 PFCCNDNVTNTA SCCLEF SCCMNE 230 LYDIRTNAVTVC CVWLQENLLAVC FSFMETNMGIII	200  N E T C T K HFWAE25.aa S E S C G L NAG-2.aa T D - K N P TALLA-1.aa  240 G G HFWAE25.aa A G TALLA-1.aa
151 149 147 190 180 179	170  GFTN T D - FEDS  GVSN T D W FEV -  VON T N - W S T S  210  QKAHDQKVE  OHAPGTWWKA  QDLHNLTVAATK  250  BVAAGIG-GLELA  ALVOIL-GLTFA	180  PYFKENSAFPFYNATRVYNATRV PYFLEHGI	190  FCCNDNVTNTA  SCCMNE  230  YDIRTHAVTV  VWLQENLLAV  SFMETNMGIII	200  A N E T C T K HFWAE25.aa  S E S C G L NAC-2.aa  T D - K N P TALLA-1.aa  240  G G HFWAE25.aa  A G TALLA-1.aa  280  L P L P L L L HFWAE25.aa  NAG-2.aa
151 149 147 190 180 179	170  GFTN T D - FEDS  GVSN T D W FEV -  VON T N - W S T S  210  QKAHDQKVE  OHAPGTWWKA  QDLHNLTVAATK  250  BVAAGIG-GLELA  ALVOIL-GLTFA	180  P Y F K E N S A F P F F F F F F F F F F F F F F F F F	190 PFCCNDNVTNTA SCCLEF SCCMNE 230 LYDIRTNAVTVC CVWLQENLLAVC FSFMETNMGIII	200  N E T C T K HFWAE25.aa S E S C G L NAC-2.aa T D - N P TALLA-1.aa  240 G G HFWAE25.aa A G TALLA-1.aa  280 L P L P L L L HFWAE25.aa
151 149 147 190 180 179	170  G F T N X T D - F E D S  G V S N Y T D W F E V -  V O N Y T N - W S T S  210  Q K A H D Q K V E  O H A P G T W W K A  Q D L H N L T V A A T K  250  B V A A G I G - G L E L A  A L V Q I L - G L T F A	180  PYFKENSAFPFYNATRVYNATRV PYFLEHGI	190 PFCCNDNVTNTA SCCLEF SCCMNE 230 LYDIRTNAVTVC CVWLQENLLAVC FSFMETNMGIII	200  A N E T C T K HFWAE25.aa S E S C G L NAC-2.aa T D - C N P TALLA-1.aa  240  G G HFWAE25.aa A G TALLA-1.aa  280  L P L P L L L HFWAE25.aa NAG-2.aa TALLA-1.aa  TALLA-1.aa
151 149 147 190 180 179	170  GFTNYTD-FEDS GVSNYTDWFEV- GVONYTN-WSTS  210  QKAHDQKVE DHAPGTWWKA QDLHNLTVAATK  250 BVAAGIG-GLELA ALVQIL-GLTFA VAFGIAFSQLIG	180  PYFKENSAFPE YNATRVYNATRVE  220	190  PFC NDNVTNTA  SCC LEF 230  LYDIRTNAVTV  (VWLQENLLAV)  FSFMETNMGILL  270  VSVLQSTISPL	200  A N E T C T K HFWAE25.aa  S E S C G L NAC-2.aa  T D - K N P TALLA-1.aa  240  G G HFWAE25.aa  A G TALLA-1.aa  280  L P L P L L L HFWAE25.aa  NAG-2.aa

FIGURE 4

_	10	20	30	40
1	MNSMTSAVPVANSVL MTTPRNSVNGTF	- <b>.</b>	TPGIMSHVP	LYPNSQ HIPEF86.aa
41. 23	PQVHLVPGNPPSEVS MQSGPKPEFR	60 NVNGQPVQKA RMSSLVGPT	70 1 1 L K E G K T 2 S F F M R E S K T	80 LGAIQI HIPEF86.aa LGAVQI B1.aa
78 58	MNGLFEIALGGLLMI	100 VLVSEYLSI P-ASIYAPI	110 5 F Y G G F P F W G C V T V W Y F L W G	120 G L W F I I HIPEF86.aa G I M Y I I B1.aa
118 97	130 S G S L S V E A E N Q P Y S Y S G S I L A G T E K N S R K -	140 GILSGSLGL GLVKGKMIM	150 IVSAICSAV SLSLFAAIS	160 S V I G F I HIPEF86.aa G M I I S I B1.aa
158 136	170  T D L S I P H	180 	190 A PDY PYAW PYINIYNCE-	2000 GV N P G - HIPEF86.aa PAN P S E B1.aa
182 175	210 KNSPSTQYCYSIQSL	220 - MAISGVLU FLGILSVM	230 V F C E L E I P A F F Q E E V I	FGI HIPEF86.aa AGIVEN B1.aa
199 215	250 ACASSHFGCQL EWKRTCSRPKSNIVL	260 V C C Q S S N V S L S A E E K K E Q	270 V I Y P N I Y A A N T I E I K E	PWITPE HTPEF86.aa EWVGLT Bl.aa
235 251	PVTS PPSYSETQ ETSSQPKNEEDIEII	300  PIQEEEEEE	310 TETNFPEPPQ	320 
248 291	ANK. IENDSSP			HIPEF86.aa Bl.aa

#### FIGURE 5





FIGURE 6

		~	20	40
1	10 MG-QCGITSSKTVLV METKPVITCLKTLL	20 / FLNLIEWGA A I I Y S F V E W I T (	30 AGIICYVGAYV GVIILAVSVWG	FIRYD HLTAH80.aa
40 41	D Y D H F F E D V Y T L I P A	O O O O V V I I A V C A L I Y V L I G T G T T	70 LFIIGLIGCCA IVVFGLFGCFA	80 TIRES HLTAH80.aa FCRGS TALLA-1.aa
80 80	RCGLATFVIILLU PWMLKLYAMFUSLV	100 VTEVVVVVL LASLVAGIS	110 Y V Y R A K V E N E F V F R H E I K D T	120 V D & S I HLTAH80.aa F L & T Y TALLA-1.aa
120 120	130 QKVYKTYHSTNPDAA TDAMQTYTSNDEF	140 10 1 S R A I D Y V Q R ( 1 S R A V D H V Q R S	150 QLHCCGIHNYS 5LSCCGVQNYT	D W E N T HLTAH80.aa N W S T S TALLA-1.aa
160 158		180 RETASNEM CMNET-DCN	190 G S L A - H P P Q D L H N L T V A A	200 S D L Y A HLTAH80.aa T K V N Q TALLA-1.aa
195 194		220 1 1 M H V I W M A L A I I M G I I – A G V A	230 AAIQLLGM GIAFSQLIGM	240 LACI HLTAH80.aa LACC TALLA-1.aa
233 232		260 TGGTYA. TANOYEMV		HLTAH80.aa TALLA-1.aa

#### FIGURE 7



#### FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11

			30	40
	10	20		TO STREET
1	MGSRKCGGCLSCLLI	PLALWSIIVN	ILLYFPNGQ	rky & S.F. I.6 aa
1	MGSRECGGCLSCLUI MCYGECARCIGHSIV	JG BARLC AAW	(被)(被)(被)(被)(死)(死)(死)(死)(死)(死)( <b>2</b> (2)	A TOTAL
			70	<del></del>
	50	60		
41	MKLTNYWYFEGICF	SSIMMLIVIT	ALTAIRNAN	NYKCCQ HUVBB80.aa
41	M K L T N Y V W Y F E G I C F M H L S R F V W F F S G I V C	G G L L M B L P A F	<u> </u>	
		100	110	120
	90	100		_,
81	SENCSKKYVTLLSI	FSSLGTAFSC AALIGTAGSC	YCLVISABG	A F O P I I I A A A
81	HEDCGKRCAMISSVI	JAA LII 编编和GAN	HANNE A T A) WINNERSON	
		140	150	160
	130	140		1
121	C - RTLDGWEXAFEG	TAGRELTOSSI	MIQCLLEAAN	T V E W N V L6.aa
121	C L D S I G Q W N Y T P A S	EGQY		
		180	190	200
	170			YSVIF HUVBB80.aa
160	ILFSILITLSGLQV SLFSILLALGGIEF	IICLIRVYMQI	JSKILL GFCC	HOOOY L6.aa
161	SIRSILLAIGGIEF	新丁! (機能能	The German Co	(m) LEC X X
				HLVBB80.aa
197	QPGII.			L6.aa
201	D.C.			

FIGURE 12

	10	20	30 4	0
1	10 M S P R L E M S S F S T Q T P Y P N L A M M L S L N N L	VPC	SHALFQGUSPGQV FTSIFNGUYPSKS VGTIPDQUDFGT	L / HJACE54.aa 5 rGalectin-5.aa
	50	60	70 8	0
	IIVRG LVLQEPKH BTV IVISG VVLSDAKREQI IVIRGHVPSDADRFOV	INDED C = C C = = = =		- HJACE54.aa - rGalectin-5.aa < hGalectin-8.aa
	90	100	110 11	20
46 53 72	RAGCIVCNTLINEKWG	REEITYDTPFQ	KEKKSFEIVIMV	- HJACE54.aa - rGalectin-5.aa L hGalectin-8.aa
	130	140	150 1	<del></del> 60
46				- HJACE54.aa - rGalectin-5.aa
53 112	KAKFQVAVNGKHTLLY	GHRIGPEKIDT	LGIYGKVNIHSI	G hGalectin-8.aa
	170	180	190 2	<del>1-</del> 00
46				- HJACE54.aa - rGalectin-5.âa
53 152	F S F S S D L Q S T Q A S S L F	LTEISRENVPK	SGTPQLRLPFAA	
		220		<del>1 -</del> 40
46	210			HJACES4.aa
53 192	LNTPMGPGRTVVVKGI	EVNANAKSFNVD	LLAGKSKDIALH	rGalectin-5.aa Malectin-8.aa
22				<del>T</del> 280
	250 RASFADRTLAWIS	260 R W 😂 - OKKLIS	A PETENPORFEE	U HJACE54.aa
52 59	NPRFDENAVVRNTQII NPRLNIKAFVRNSFL	T NT C:347:65: D F F F R S I. F	プログログログ はんしゅん はんしゅん はんしゅん はんしゅん しゅんしゅん しゅん	A TOTTCOOT
232	NIPRIINI KAIF VIKNOJE I.	Z E O BRIDGE B D D A A A		┰
	290	300		320 L V HTACE54 aa
87 99 27 <b>1</b>	LLLFQEGGLELALNG WILCEGHCFEVAVDG IIYCDVREFEVAVNG			
	330			
	Q Y C V H S .			HJACE54.aa rGalectin-5.aa
139 31 <b>1</b>	Q			hGalectin-8.aa

#### FIGURE 13






FIGURE 14



#### FIGURE 15



### PCT

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



# INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/12, 15/85, 5/10, C07K 14/705, 16/28, C12Q 1/68, A61K 38/17, 48/00

**A3** 

US

US

(11) International Publication Number:

WO 98/31799

(43) International Publication Date:

23 July 1998 (23.07.98)

(21) International Application Number:

PCT/US98/00959

(22) International Filing Date:

21 January 1998 (21.01.98)

(30) Priority Data:

60/034,204 21 January 1997 (21.01.97) 60/034,205 21 January 1997 (21.01.97)

(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): NI, Jian [CN/US]; 5502
Manorfield Road, Rockville, MD 20853 (US). GENTZ,
Reiner, L. [DE/US]; 13404 Fairland Park Drive, Silver
Spring, MD 20904 (US). ROSEN, Craig, A. [US/US]; 22400
Rolling Hill Road, Laytonsville, MD 20882 (US).

(74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report:
5 November 1998 (05.11.98)

(54) Title: POLYNUCLEOTIDES AND POLYPEPTIDES ENCODING RECEPTORS

#### (57) Abstract

Receptor polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing receptor polypeptides and polynucleotides in the design of protocols for the treatment of diseases and diagnostic assays for such conditions.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Моласо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	ТJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
B.J	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	ΪL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA.	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		•
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# INTERNATIONAL SEARCH REPORT

ternational Application No PCT/US 98/00959

A CLASS	SIFICATION OF SUBJECT MATTER		
ÎPC 6	C12N15/12 C12N15/85 C12N5, C12Q1/68 A61K38/17 A61K48		C07K16/28
According	to International Patent Classification (IPC) or to both national class	ification and IPC	
B. FIELDS	SEARCHED	٠.	
IPC 6	ocumentation searched (classification system followed by classific C12N C07K C12Q A61K	. ,	
	tion searched other than minimum documentation to the extent the	·	
	lata base consulted during the international search (name of data	base and, where practical, search ten	ms used)
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the re	elevant passages	Relevant to claim No.
X	DATABASE GENBANK Accession No. T48852, 8 Februar "EST70319 from Stratagene plac XP002066898 see the whole document		1-12,20
X	DATABASE GENBANK Accession No. G14442, 4 January "STS SHGC-10055" XP000206689 see the whole document	1996 -/	1-3,20
X Furth	er documents are listed in the continuation of box C.	Patent family members are	listed in annex.
A' document consider E' earlier do filing dai L' document which is citation of the me comment of the me comment of the me comment of the comment of the action of the acti	t which may throw doubts on priority claim(s) or cited to establish the publication date of another or other special reason (as specified) t referring to an oral disclosure, use, exhibition or	T later document published after the or priority date and not in conflicited to understand the principal invention  "X" document of particular relevance cannot be considered novel or involve an inventive step when "Y" document of particular relevance cannot be considered to involve document is combined with one ments, such combination being in the art.  "&" document member of the same published with the art.  Date of mailing of the internation	ct with the application but e or theory underlying the e; the claimed invention cannot be considered to the document is taken alone e; the claimed invention e an inventive step when the e or more other such docu- cobvious to a person skilled patent family al search report
lame and ma	ifing address of the ISA  European Patent Office, P.B. 5818 Patentiaan 2  NL - 2280 HV Rijswijk  Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  Fax: (+31-70) 340-3016	Authorized officer  Galli, I	

# INTERNATIONAL SEARCH REPORT

PC17US 98/00959

		PC1/03 98/00959
C.(Continua	tion) DOCUMENTS CONSIDERED TO BE RELEVANT	Relevant to claim No.
Category °	Citation of document, with indication, where appropriate, of the relevant passages	New and to claim to
А	ZANNETTINO A.C.W. ET AL.: "A powerful technique for isolating genes encoding surface antigens using retroviral expression cloning" J. IMMUNOL., vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 see abstract	1-21
Α	ADAMS M D ET AL: "COMPEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 cited in the application see abstract	1-21
А	SIMMONS D. & SEED B.: "Isolation of a cDNA encoding CD33, a differentiation antigen of myeloid progenitor cells" J. IMMUNOLOGY, vol. 141, no. 8, 15 October 1988, pages 2797-2800, XP002066894 see abstract see figure 4	1-21
A	CROCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 3 October 1994, pages 4490-4503, XP002066895 see abstract see figure 2	1-21
A	KELM S. ET AL.: "The Sialoadhesins: a family of sialic acid-dependent cellular recognition molecules within the immunoglobulin superfamily" GLYCOCONJ. J., vol. 13, no. 6, December 1996, pages 913-926, XP002066896 see the whole document	1-21
Α	DATABASE GENBANK Accession No. U71382, 12 November 1996 PATEL N. ET AL.: "OB binding protein - 1" XP002066901 cited in the application see the whole document	1-21

0.10	Continues and DOCUMENTO CONTINUES		PC1/05 98/00959	
C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT  Category *   Citation of document, with indication; where appropriate, of the relevant passages   Relevant to claim No.				
P,X	DATABASE GENBANK Accession No. AA344713, 21 April 1997 "EST50650, H. sapiens cDNA similar to CD33 antigen" XP002066900 see the whole document		1-12,20	
, A	TAKEI Y. ET AL.: "Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta" CYTOGENET. CELL GENET., vol. 78, 1997, pages 295-300, XP002066897 see abstract see figure 1		1-21	

PCT/US 98/00959

Box I	Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)			
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:				
1. X 2.	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:  Remark: Although claims 13 and 14 are directed to a method of treatment of the human/animal body, and claim 15 (as far as in vivo methods are envisaged) is directed to a diagnostic method, the search has been carried out and based on the alleged effects of the compound/composition.  Claims Nos.:  Claims Nos.:  because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:			
з. 📗	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).			
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)			
	ernational Searching Authority found multiple inventions in this international application, as follows:  e further information sheet			
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.			
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.			
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:			
4. X	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  see further information sheet, subject 1.			
Remari	The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.			

#### INTERNATIONAL SEARCH REPORT

International Application No. PCT/ US 98/00959

#### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-21) - partial

An isolated polynucleotide encoding the HMACR70 cell surface receptor (Seq. IDs 1,18), polynucleotides complementary thereto, fragments thereof, homologs thereof.

An expression system comprising said polynucleotides, a process to produce a cell harbouring said expression system. Said cell or membrane thereof expressing said receptor.

A process for the production of the HMACR70 polypeptide comprising culturing said cell.

Said HMACR70 polypeptide or homologs. An antibody directed against said polypeptide.

Therapeutic and diagnostic applications.

A method to identify agonists or antagonists to said receptors. Said agonists and antagonists.

2. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 2,3,19

3. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 4,20

4. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 5,21

5. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 6,22

6. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 7,23

7. Claims: (1-21) - partial

Idem as subject matter 1 but limited to Seq. IDs 8,24

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/ US 98/00959

### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

- 8. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 9,10,25
- 9. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 11,26
- 11. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 13,28
- 12. Claims: (1-21) partial
  Idem as subject matter 1 but limited to Seq. IDs 14,29
- 13. Claims: (1-21) partial
  Idem as subject matter 1 but limited to Seq. IDs 15,30
- 14. Claims: (1-21) partial
   Idem as subject matter 1 but limited to Seq. IDs 16,31
- 15. Claims: (1-21) partial

  Idem as subject matter 1 but limited to Seq. IDs 17,21